

CELL COMPLEXES, POSET TOPOLOGY AND THE REPRESENTATION THEORY OF ALGEBRAS ARISING IN ALGEBRAIC COMBINATORICS AND DISCRETE GEOMETRY

STUART MARGOLIS, FRANCO SALIOLA, AND BENJAMIN STEINBERG

ABSTRACT. In recent years it has been noted that a number of combinatorial structures such as real and complex hyperplane arrangements, interval greedoids, matroids and oriented matroids have the structure of a finite monoid called a left regular band. Random walks on the monoid model a number of interesting Markov chains such as the Tsetlin library and riffle shuffle. The representation theory of left regular bands then comes into play and has had a major influence on both the combinatorics and the probability theory associated to such structures. In a recent paper, the authors established a close connection between algebraic and combinatorial invariants of a left regular band by showing that certain homological invariants of the algebra of a left regular band coincide with the cohomology of order complexes of posets naturally associated to the left regular band.

The purpose of the present monograph is to further develop and deepen the connection between left regular bands and poset topology. This allows us to compute finite projective resolutions of all simple modules of left regular band algebras over fields and much more. In the process, we are led to define the class of CW left regular bands as the class of left regular bands whose associated posets are the face posets of regular CW complexes. Most of the examples that have arisen in the literature belong to this class. A new and important class of examples is a left regular band structure on the face poset of a CAT(0) cube complex. Also, the recently introduced notion of a COM (complex of oriented matroids or conditional oriented matroid) fits nicely into our setting and includes CAT(0) cube complexes and certain more general CAT(0) zonotopal complexes. A fairly complete picture of the representation theory for CW left regular bands is obtained.

Date: August 25, 2015.

2010 *Mathematics Subject Classification.* 20M30, 16G10, 05E10, 52C35, 52C40, 16S37, 20M25, 52B05, 16E10.

Key words and phrases. regular cell complexes, quivers, finite dimensional algebras, hyperplane arrangements, zonotopes, oriented matroids, CAT(0) cube complexes, CAT(0) zonotopal complexes, left regular bands.

This work was partially supported by a grant from the Simons Foundation (#245268 to Benjamin Steinberg) and the Binational Science Foundation of Israel and the US (#2012080 to Stuart Margolis and Benjamin Steinberg). Also Steinberg was supported in part by a CUNY Collaborative Research Incentive Grant. The second author was supported by an NSERC grant.

CONTENTS

1. Introduction	3
2. Left regular bands, hyperplane arrangements, oriented matroids and generalizations	10
2.1. Free left regular bands and matroids	14
2.2. Free partially commutative left regular bands	17
2.3. Hyperplane arrangements and oriented matroids	17
2.4. Strong elimination systems, lopsided systems and COMs	24
2.5. Complex hyperplane arrangements	27
3. Regular CW complexes and CW posets	29
3.1. Simplicial complexes	29
3.2. Regular CW complexes and CW posets	31
3.3. Oriented interval greedoids	36
3.4. The topology of left regular bands	44
3.5. CAT(0) cube complexes	48
3.6. CAT(0) zonotopal complexes	57
4. Algebras	58
4.1. Finite dimensional algebras	58
4.2. Gradings, quadratic algebras and Koszul algebras	64
4.3. The algebra of a left regular band	65
4.4. Existence of identity elements in left regular band algebras	75
4.5. Cartan invariants	77
5. Projective resolutions and global dimension	78
5.1. Actions of left regular bands on CW posets	78
5.2. Projective resolutions	82
5.3. Ext and global dimension	87
5.4. Minimal projective resolutions	93
6. Quiver presentations	95
6.1. A general result	95
6.2. A quiver presentation for CW left regular bands	97
7. Quadratic and Koszul duals	98
7.1. Quadratic duals	98
7.2. Koszul duals	103
8. Injective envelopes for hyperplane arrangements, oriented matroids, CAT(0) cube complexes and COMs	104
8.1. Hyperplane arrangements	106
8.2. Oriented matroids	110
9. Enumeration of cells for CW left regular bands	113
9.1. Flag vectors	113
9.2. Cartan invariants	117
10. Cohomological dimension	118
References	120
Index	126

1. INTRODUCTION

We begin with a brief history of the background and motivation for the topics of this monograph before delving into the new results and the organization of the paper. An announcement of the results of this paper appears in [74].

Application to Markov chains. In a highly influential paper [17], Bidigare, Hanlon and Rockmore showed that a number of popular Markov chains, including the Tsetlin library and the riffle shuffle, are random walks on the faces of a hyperplane arrangement (the braid arrangement for these two examples). More importantly, they showed that the representation theory of the monoid of faces, where the monoid structure on the faces of a central hyperplane arrangement is given by the Tits projections [114], could be used to analyze these Markov chains and, in particular, to compute the spectrum of their transition operators.

The face monoid of a hyperplane arrangement satisfies the semigroup identities $x^2 = x$ and $xyx = xy$. Semigroups satisfying these identities are known in the literature as left regular bands (although they were studied early on by Schützenberger [102] under the more descriptive name “*treillis gauches*,” translated by G. Birkhoff in his Math Review as skew lattices, a term which nowadays has a different meaning). Brown developed [34, 35] a theory of random walks on finite left regular bands. He gave numerous examples that do not come from hyperplane arrangements, as well as examples of hyperplane walks that could more easily be modeled on simpler left regular bands. For example, Brown considered random walks on bases of matroids. Brown used the representation theory of left regular bands to extend the spectral results of Bidigare, Hanlon and Rockmore [17] and gave an algebraic proof of the diagonalizability of random walks on left regular bands.

Brown’s theory has since been used and further developed by numerous authors. Diaconis highlighted hyperplane face monoid and left regular band walks in his 1998 ICM lecture [49]. Björner used it to develop the theory of random walks on complex hyperplane arrangements and interval greedoids [23, 24]. Athanasiadis and Diaconis revisited random walks on hyperplane face monoids and left regular bands in [7]. Chung and Graham considered further left regular band random walks associated to graphs in [43]. Saliola and Thomas proposed a definition of oriented interval greedoids by generalizing the left regular bands associated to oriented matroids and antimatroids [100]. A detailed study of symmetrized versions of random walks on hyperplane face monoids was initiated in recent work of Reiner, Saliola and Welker [91]. Left regular bands have also appeared in Lawvere’s work in topos theory [69, 70] under the name “graphic monoids.”

The representation theory of hyperplane face monoids is also closely connected to Solomon’s descent algebra [105]. Bidigare showed in his thesis [18] (see also [35]) that if W is a finite Coxeter group and $\mathcal{A}_W$ is the associated

reflection arrangement, then the descent algebra of W is the algebra of invariants for the action of W on the algebra of the face monoid of $\mathcal{A}_W$. This, together with his study of the representation theory of hyperplane face monoids [98], allowed the second author to compute the quiver of the descent algebra in types A and B [97] (see also [101]), and to compute the Loewy length of the descent algebra of type D [99].

To further emphasize the scope and applicability of the theory of left regular bands, we remark that the original version of Brown's book on buildings [31] makes no mention of the face monoid of a hyperplane arrangement, whereas it plays a prominent role in the new edition [1]. Hyperplane face monoids also have a salient position in the work of Aguiar and Mahajan on combinatorial Hopf algebras and Hopf monoids in species [4, 5].

Techniques from the topology of hyperplane arrangements. Using the topology of hyperplane arrangements, Brown and Diaconis [36] found resolutions of the simple modules for the face monoid that were later shown by the second author to be the minimal projective resolutions [98]. Brown and Diaconis used these resolutions to prove diagonalizability of the transition operator of the hyperplane chamber random walk. Bounds on rates of convergence to stationarity were obtained in [17, 36]. They observed, moreover, that one can replace the faces of a hyperplane arrangement by the covectors of an oriented matroid [26] and the theory carries through.

The resolutions of Brown and Diaconis coming from the topology of hyperplane arrangements also played an important role in work of the second author. They were used to compute the Ext-spaces between simple modules for face monoids of hyperplane arrangements [98]. He also computed a quiver presentation of such algebras. Consequently, he computed the global dimension of these algebras. (Left regular band algebras have acyclic quivers and hence finite global dimension, which makes computing their global dimension a natural question.)

In [96] the second author computed the projective indecomposable modules for arbitrary left regular band algebras and also the quiver. The topological arguments to compute minimal projective resolutions and Ext-spaces did not immediately extend to this setting because one does not have any obvious topology available. Our previous paper [75] provided the appropriate topological tools and will be expanded upon, and simplified, in this work.

The paper [96] also contained an intriguing unpublished result of Ken Brown stating that the algebra of a free left regular band is hereditary. The proof is via a computation of the quiver. A theorem of Gabriel [6] states that a split basic finite dimensional algebra is hereditary if and only if it is isomorphic to the path algebra of an acyclic quiver. Since left regular band algebras are split basic (over any field), Brown's proof amounts to proving that the dimension of the path algebra of the quiver of the free left regular band is the cardinality of this monoid.

The third author (unpublished) showed that Brown’s counting argument could be modified to prove that the algebra of any left regular band whose poset of principal right ideals has a Hasse diagram which is a tree is hereditary. Brown’s result is a direct consequence because the Hasse diagram of the poset of principal right ideals of a free left regular band is a tree. On the other hand, the poset of principal right ideals of a hyperplane monoid is dually isomorphic to the poset of faces of the arrangement. This offers a topological explanation for the computations of Brown and Diaconis and of Saliola cited in the previous paragraph.

These two examples strongly suggested that there was a deep connection between the representation theory and cohomology of the algebra of a left regular band B and the topology of the poset of principal right ideals of B . There seem to be only a handful of results in the finite dimensional algebra literature that use topological techniques to compute homological invariants of algebras. The primary examples seem to be in the setting of incidence algebras, where the order complex of the poset plays a key role [44, 55, 62]. A more general setting is considered in [38].

In our previous paper [75], we clarified this connection by using topological techniques to compute all the Ext-spaces between simple modules of the algebra of a left regular band monoid. In particular, we used order complexes of posets and classifying spaces of small categories (in the sense of Segal [103]) to achieve this. A fundamental role was played by the celebrated Quillen’s Theorem A [90], which gives a sufficient condition for a functor between categories to induce a homotopy equivalence of classifying spaces. Somewhat surprisingly to us, a combinatorial invariant of simplicial complexes, the Leray number [63, 64], played an important part in [75]. The Leray number is tied to the Castelnuovo-Mumford regularity of Stanley-Reisner rings. In particular, that paper gives a new, non-commutative interpretation of the regularity of the Stanley-Reisner ring of a flag complex.

CW left regular bands. Despite the use of this heavy machinery, we were not able to directly compute finite projective resolutions for the simple modules of the algebra of a left regular band. One of the main contributions of the current work is to do just that. Our starting point is the well-known observation that the face poset of a central hyperplane arrangement in $\mathbb{R}^n$ is dually isomorphic to the face poset of a regular CW complex decomposition of a ball—in fact, the face poset of an important type of polytope known as a zonotope [122]—and that the same is true for every contraction of the arrangement. Moreover, the face poset of the zonotope is isomorphic to the poset of principal right ideals of the face monoid. These observations motivate the following important definition in this paper.

We define a left regular band B to be a CW left regular band if every “contraction” (certain upper sets of a poset, corresponding to contractions in the case of the face poset of a central hyperplane arrangement or oriented matroid) of the poset of principal right ideals of B is isomorphic to the

face poset of a connected regular CW complex. Most of the examples that have been studied in the literature have this property. We study actions of left regular bands B on CW posets P , that is, face posets of regular CW complexes. We prove that under a technical restriction (that we term semi-freeness of the action) and when the (order complex of the) poset P is acyclic, then the augmented chain complex of the associated CW complex is a projective resolution of the trivial $\mathbb{k}B$ module, where $\mathbb{k}$ is any commutative ring with unit.

In particular, this allows us to show that if B contains a two-sided identity (that is, if B is a left regular band *monoid*) then the augmented chain complex of the order complex of B is a projective resolution of the trivial $\mathbb{k}B$ -module. More generally, this result holds for any connected left regular band (these are defined in terms of a topological condition that turns out to be equivalent to the semigroup having a unital algebra over any ground ring). This allows us to compute all Ext-spaces between simple $\mathbb{k}B$ -modules and thus the global dimension of the algebra.

In the case that B is a connected CW left regular band, we also compute a quiver presentation of $\mathbb{k}B$, which is an exact generalization of the second author's results in the case of real central hyperplane arrangement monoid [98]. However, even when restricted to the case of hyperplane arrangements, our approach here is fundamentally different from the original approach of [98] in that Brown and Diaconis [36] had to carefully choose orientations for the cells of the zonotope (or incidence numbers) in order to ensure that the augmented cellular chain complex of the zonotope was a chain complex of modules rather than just vector spaces. This would have been virtually impossible to extend to the level of generality we are considering here. Instead, we construct actions of the left regular bands on their associated cell complexes by cellular maps. Functoriality of the cellular chain complex then comes into play to turn it into a chain complex of modules. To the best of our knowledge it was not even previously known that hyperplane face monoids act on their associated zonotopes by cellular mappings.

Outline. We provide an outline of the paper here. We have tried to keep the paper as self-contained as possible by providing background material in a number of fields because we are targeting the paper at a variety of audiences, in particular, toward specialists in algebraic combinatorics, representation theory, poset topology and semigroup theory.

In Section 2 we recall the definitions and basic properties of left regular bands. In particular, left regular bands are also posets and left multiplication induces an action on the corresponding poset and on the order complex of this poset. This observation plays a crucial role in the paper.

We define a number of new operations on left regular bands such as suspension and join that model the corresponding operations for posets and simplicial complexes. Also the fundamental notion of a contraction of a

left regular band is introduced. We give examples of left regular bands that have arisen in the literature and that play an important part in this work. We pay particular attention to the left regular band structure on various combinatorial structures such as real and complex hyperplane arrangements [17, 23], matroids [34, 117], oriented matroids [26], COMs [13] and interval greedoids [23]. We include the necessary background material for these structures as well.

Section 3 begins with a survey of cell decompositions of various types. All are examples of regular CW complexes. We look at the corresponding notion of CW posets, which by definition are posets isomorphic to face posets of regular CW complexes. Novel to this text is that we promote the standard construction [21] of a regular CW complex from a CW poset to a functor by establishing the functorial nature of the construction with respect to appropriate mappings of posets that we term “cellular.” This is crucial in order to transform an action of a semigroup on a CW poset into an action on the corresponding CW complex by regular cellular maps. This development could be of interest in its own right for workers in poset topology.

With this background out of the way, we are able to define the notion of CW left regular band as a left regular band all of whose contractions (as defined in Section 2) are CW posets. Most of the left regular bands that are associated to combinatorial structures are CW left regular bands. Oriented interval greedoids from [100] are presented as another naturally arising family of CW left regular bands.

An important new example of CW left regular bands is detailed in the penultimate subsection of Section 3. We recall the important notion of a cube complex and, in particular, that of a CAT(0) cube complex. There is a natural notion of hyperplane in such complexes and this leads us to define the structure of a left regular band associated to a CAT(0) cube complex by associating covectors to the faces of a CAT(0) cube complex. CAT(0) cube complexes have played an important role in geometric group theory and three-dimensional manifold theory in recent years, in particular, with respect to the work of Agol and Wise on Thurston’s virtual Haken and virtual fibering conjectures [3, 118]. In combinatorics, the Billera-Holmes-Vogtman phylogenetic tree space [19] has been a popular CAT(0) cube complex to study, particularly because of its connections with tropical geometry, where it appears in the guise of the tropical Grassmanian [106]. We provide all the necessary background on cube complexes and CAT(0) spaces. The left regular band structure for CAT(0) cube complexes was discovered independently in [13], where it is shown that there is, in fact, a left regular band structure on the face poset of any CAT(0) zonotopal complex whose cells are Coxeter zonotopes. We discuss this development briefly in the last subsection of Section 3.

In Section 4 we review several elements of the representation theory of finite dimensional algebras that may not be familiar to all our target audience. We give the necessary background on the Gabriel quiver, quiver

presentations and the connections with split basic algebras. We recall the definition of a Koszul algebra and its Koszul dual. We then review known results on the algebra of a left regular band monoid and its representation theory. We note that the algebra of any left regular band semigroup has a right identity element and show that the algebra has an identity element if and only if every contraction of the semigroup (in the sense noted above) is connected. By extending the setting from monoids to semigroups with unital algebras, we are able to treat a number of important examples—such as affine hyperplane arrangements, affine oriented matroids, $\text{CAT}(0)$ cube complexes and, more generally, COMs—whose associated left regular bands are not monoids but do have unital algebras.

Section 5 is devoted to one of the central results of this work. We introduce the notion of a semi-free action of a left regular band on a poset. We show that if a left regular band admits a semi-free action on an acyclic CW poset P , then the augmented cellular chain complex of the associated CW complex is a projective resolution of the trivial $\mathbb{k}B$ -module. In particular, this applies to the poset of faces of the order complex of B if B is a monoid, or more generally if B is connected (so that its algebra is unital), and so the augmented simplicial chain complex of the order complex of B is a projective resolution of the trivial $\mathbb{k}B$ -module. We use these projective resolutions to compute all Ext-spaces between all simple modules for monoid (and connected) left regular band algebras over a field, providing a more direct and enlightening proof of the main results of our previous paper [75].

We show that the global dimension of a CW left regular band B is equal to the dimension of the CW complex associated to B . For CW left regular bands, the chain complexes we discuss in this section are actually the minimal projective resolutions of every simple module over a field of coefficients. In particular, this applies to the face monoid of a real or complex hyperplane arrangement, the monoid of covectors of an oriented matroid or oriented interval greedoid, the face semigroup of an affine hyperplane arrangement or $\text{CAT}(0)$ cube complex, or the semigroup of covectors associated to a COM.

We turn to quiver presentations in Section 6. We abstract the argument used by the second author in the case of real hyperplane monoids [98] to prove that if B is a connected CW left regular band (so, in particular, if B is a monoid), then over a field of coefficients $\mathbb{k}$, the quiver of $\mathbb{k}B$ is isomorphic to the Hasse diagram of the maximal semilattice image (or support semilattice) $\Lambda(B)$ of B , which is a graded semilattice. Furthermore, $\mathbb{k}B$ is the quotient of the path algebra of its quiver by adding one relation for each interval of length 2 in the Hasse diagram of $\Lambda(B)$, which is just the sum of all paths that begin at the bottom point of the interval and end at the top point. Thus $\mathbb{k}B$ is completely determined by $\Lambda(B)$.

It follows from the results of Section 6 that the algebra $\mathbb{k}B$ of a CW left regular band is a quadratic algebra—it has a basis of quiver relations of degree 2. We prove in Section 7 that the algebra of CW left regular band is, in fact, a Koszul algebra. We show that the Koszul dual of $\mathbb{k}B$ is the opposite

of the incidence algebra of the algebra of the support semilattice $\Lambda(B)$. It follows that the Ext-algebra of $\mathbb{k}B$ is isomorphic to the incidence algebra of $\Lambda(B)$. Results proved independently by Polo [87] and Woodcock [119] then yield that every open interval of the semilattice $\Lambda(B)$ is a Cohen-Macaulay poset. This is particularly important because it has consequences for the Möbius function of $\Lambda(B)$. To the best of our knowledge, the results of Polo and Woodcock have never before been used to prove a poset is Cohen-Macaulay, but rather have been used to prove that certain incidence algebras are Koszul.

Section 8 studies injective envelopes. In particular, we give an explicit geometric construction of the injective envelopes of the simple $\mathbb{k}B$ -modules when B is a CW left regular band whose cells are zonotopes (or, more generally, dual to face posets of oriented matroids). In particular, the result applies to COMs. The construction is based on the idea of performing a line shelling of the zonotope in order to construct a visual hemisphere whose faces in turn span a projective right ideal in the algebra.

There are many famous enumerative results that count cells in hyperplane arrangements and, more generally, in oriented matroids. In particular, the theorem of Las Vergnas and Zaslavsky [26] computes the number of chambers in a real hyperplane arrangement (or topes in an oriented matroid) in terms of the Möbius function of its lattice of flats (intersection lattice). From our perspective, the number of chambers is equal to the number of elements in the minimal ideal of the hyperplane face monoid. This leads to a natural direction for generalizing these results. The main results of Section 9 do just that for connected CW left regular bands: we count the number of cells in each dimension in terms of the Möbius function of the support semilattice. When our result is restricted to hyperplane arrangements and oriented matroids we recover the Las Vergnas-Zaslavsky theorem discussed above. In the case of complex hyperplane arrangements we recover a result of Björner [23] and in the case of CAT(0) cube complexes we recover an enumerative result of Dress *et. al* [51], originally expressed in the equivalent language of median graphs. We use these results to generalize the second author's computation of the Cartan matrix of hyperplane face monoids to that of all connected CW left regular bands.

In the last section of the paper we compute the cohomological dimension of left regular band monoid algebras. We show that there is a surprising connection to Leray numbers. The cohomological dimension of the free partially commutative left regular band on a graph Γ is the Leray number of the clique complex of Γ . We show how to use our result to construct easily a finite monoid (in fact, a regular band, that is a submonoid of a direct product of a left regular band and right regular band) that has left cohomological dimension m and right cohomological dimension n for any pair of natural numbers m and n . (The reader is invited to compare with the construction in [60].)

2. LEFT REGULAR BANDS, HYPERPLANE ARRANGEMENTS, ORIENTED MATROIDS AND GENERALIZATIONS

In this paper, all semigroups and monoids are assumed finite unless otherwise stated. Standard references on semigroup theory are [46, 92]. In this paper we shall make some use of *Green's relations* [58]. Two elements $s, t \in S$ of a semigroup S are said to be $\mathcal{J}$ -equivalent (respectively, $\mathcal{R}$ -equivalent, respectively $\mathcal{L}$ -equivalent) if they generate the same principal two-sided (respectively, right, respectively, left) ideal. An element $e \in S$ is *idempotent* if $e^2 = e$. In this case, notice that the principal left ideal generated by e is Se and that $a \in Se$ if and only if $a = ae$. Indeed, if $a \in Se$, then $a = xe$ with $x \in S$ and so $ae = xee = xe = a$. Conversely, if $ae = a$, then trivially $a \in Se$. In particular, if $e, f \in S$ are idempotents, then they are $\mathcal{L}$ -equivalent if and only if $ef = e$ and $fe = f$. Dual statements hold for the principal right ideal generated by an idempotent.

A *left regular band* is a semigroup B satisfying the identities $x^2 = x$ and $xyx = xy$. More generally, a semigroup is called a *band* if all its elements are idempotent. By a *left regular band monoid*, we mean a left regular band with identity. Left regular band monoids have also been studied by Lawvere in the context of topos theory under the name *graphic monoids* [69–71]. If B is a left regular band, then B^1 denotes the monoid obtained by adjoining an external identity element to B (even if it already had one). Notice that B^1 is a left regular band whenever B is one.

In this section we review some basic facts about left regular bands and provide a miscellany of examples coming from combinatorics and topology.

In any semigroup, there is a natural partial order on the set of idempotents defined by putting $e \leq f$ if $ef = fe = e$. In a left regular band, $fe = e$ implies $ef = efe = e$ and so $e \leq f$ if and only if $fe = e$. Said differently, Green's relation $\mathcal{R}$ is trivial on a left regular band B and the order on idempotents is Green's $\mathcal{R}$ -order, that is, $e \leq f$ if and only if $eB \subseteq fB$.

If P is a (meet) semilattice, that is, a partially ordered set with binary meets, then P is a commutative left regular band with respect to the meet operation. Moreover, $e \leq f$ if and only if $e = f \wedge e$ and so the natural partial order on P is its original ordering. Conversely, if P is a commutative left regular band, then we claim that P is a semilattice with respect to the natural partial order and the meet is the product in P . Indeed, if $e, f \in P$ then $e(e f) = e f$ and $f(e f) = (e f) f = e f$ and so $e f \leq e, f$. If $a \leq e, f$, then $ea = a$ and $fa = a$, whence $efa = ea = a$. Therefore, $a \leq e f$ and we conclude that $e f = e \wedge f$. Let us record this as a proposition.

Proposition 2.1. *Commutative left regular bands are precisely meet semilattices.*

Thus left regular bands can be thought of as “skew semilattices,” and, for this reason, Schützenberger called left regular bands by the name *treillis gauches* [102].

The set $\Lambda(B)$ of principal left ideals of a left regular band is closed under intersection and hence is a semilattice called the *support semilattice* of B ; see [34, 35, 45]. More precisely we have the following proposition, which is a special case of a theorem of Clifford on semigroups all of whose elements belong to a subgroup [45].

Proposition 2.2. *Let B be a left regular band and $a, b \in B$. Then $Aa \cap Ab = Aab = Aba$. Thus $\Lambda(B) = \{Bb \mid b \in B\}$ is a semilattice.*

Proof. First note that $aba = ab$ and $bab = ba$ implies that $Aab = Aba$. Since $Aba \subseteq Aa$ and $Aab \subseteq Ab$, we conclude that $Aab \subseteq Aa \cap Ab$. Conversely, if $x \in Aa \cap Ab$, then $xa = a$ and $xb = b$. Thus $xab = xb = x$ and so $x \in Aab$. We conclude that $Aa \cap Ab = Aab$. $\square$

In the case that B is a monoid, $\Lambda(B)$ will be a lattice with B as the top. Equipping $\Lambda(B)$ with the binary operation of intersection, the mapping

$$\sigma: B \longrightarrow \Lambda(B)$$

given by $\sigma(a) = Ba$ becomes a surjective homomorphism (by Proposition 2.2) called the *support map*. Observe that $\sigma(a) = \sigma(b)$ if and only if a, b are $\mathcal{L}$ -equivalent, if and only if both $ab = a$ and $ba = b$. The following proposition will be used many times throughout the text without explicit reference.

Proposition 2.3. *Let B be a left regular band and let $B' \subseteq B$ be a subsemigroup. Then $\Lambda(B')$ is isomorphic to the image of B' under the support map $\sigma: B \longrightarrow \Lambda(B)$.*

Proof. It suffices to show that if $a, b \in B'$, then $Ba = Bb$ if and only if $B'a = B'b$. But both these equalities hold if and only if $ab = a$ and $ba = b$. $\square$

If P is a poset and $p \in P$, put

$$P_{\geq p} = \{q \in P \mid q \geq p\}.$$

The subposets $P_{\leq p}$, $P_{> p}$ and $P_{< p}$ are defined analogously.

If $X \in \Lambda(B)$, then the *contraction* of B to X is the subsemigroup

$$B_{\geq X} = \sigma^{-1}(\Lambda(B)_{\geq X}) = \{a \in B \mid \sigma(a) \geq X\}.$$

Later we shall see that this corresponds to the operation of contraction in oriented matroid theory. Observe that $\Lambda(B_{\geq X}) \cong \Lambda(B)_{\geq X}$ by Proposition 2.3.

The support semilattice of B is the abelianization of B in the following categorical sense.

Proposition 2.4. *Let B be a left regular band and let Λ be a meet semilattice. Then any homomorphism $\tau: B \longrightarrow \Lambda$ factors uniquely through $\Lambda(B)$,*

that is, there is a unique homomorphism $\tau': \Lambda(B) \longrightarrow \Lambda$ such that the diagram

$$\begin{array}{ccc} B & \xrightarrow{\sigma} & \Lambda(B) \\ & \searrow \tau & \swarrow \tau' \\ & \Lambda & \end{array}$$

commutes.

Proof. If $\sigma(a) = \sigma(b)$, that is, $Ba = Bb$, then $ab = a$ and $ba = b$. Therefore, as Λ is commutative, we have that $\tau(a) = \tau(ab) = \tau(a)\tau(b) = \tau(b)\tau(a) = \tau(ba) = \tau(b)$. It follows that τ factors uniquely through σ . $\square$

If B is a left regular band and $a \in B$, we want to think of aB as the face poset of a closed cell of B and hence we put $\partial aB = aB \setminus \{a\}$, which we think of as the boundary of aB . In particular, if B is a left regular band monoid, then $\partial B = B \setminus \{1\}$. Note that $aB, \partial aB$ are right ideals of B . The subsemigroup aB is a left regular band monoid with identity a as $aB = aBa$. Readers familiar with oriented matroid theory should think of aB as being the result of applying a deletion to B . Observe that $\Lambda(aB) \cong \Lambda(B)_{\leq Ba}$ by Proposition 2.3.

Lemma 2.5. *If $Ba = Bb$, then aB is isomorphic to bB as a left regular band monoid via $x \mapsto bx$ for $x \in aB$ (with inverse $y \mapsto ay$ for $y \in bB$).*

Proof. Indeed, $b(xy) = bxb y$ by the left regular band axiom and if $x \in aB$ and $y \in bB$, then $abx = ax = x$ and $bay = by = y$ because $ab = a$ and $ba = b$. $\square$

It is also crucial that the action of B on itself by left multiplication is order preserving.

Lemma 2.6. *The action of B on itself by left multiplication is order preserving.*

Proof. Indeed, if $e \leq f$, then $fe = e$ and so if $b \in B$, then $(bf)(be) = bfe = be$. $\square$

It will be convenient to place into a single theorem several characterizations of left regular bands that we shall use without comment throughout the text.

Theorem 2.7. *Let S be a semigroup. Then the following are equivalent.*

- (1) S is a left regular band.
- (2) S is an $\mathcal{R}$ -trivial band, that is, a band in which $aS = bS$ implies $a = b$.
- (3) S is a band in which each left ideal is two-sided.
- (4) S is a semigroup with a homomorphism $\rho: S \longrightarrow \Lambda$ to a semilattice Λ such that each fiber (which is necessarily a semigroup) satisfies the identity $xy = x$.

	0	+	-
0	0	+	-
+	+	+	+
-	-	-	-

TABLE 1. The multiplication table for L

Proof. Suppose that S is a left regular band and $aS = bS$. Then $b = ab = aba = aa = a$ as $b \in aS$ and $a \in bS$ imply $ab = b$ and $ba = a$. Thus S is an $\mathcal{R}$ -trivial band. Conversely, if S is an $\mathcal{R}$ -trivial band and $a, b \in S$, then $abaS = abS$ because $abab = ab$. Therefore, $aba = ab$ by $\mathcal{R}$ -triviality and hence S is a left regular band. This proves the equivalence of (1) and (2).

Suppose that S is a left regular band and L is a left ideal of S . Let $a \in S$ and $b \in L$. Then from $ba = bab$ we conclude that $ba \in L$ and hence L is a two-sided ideal. Thus (1) implies (3). Assume that (3) holds and let $a, b \in S$. Then Ba is a two-sided ideal and so $ab \in Ba$. Therefore, $aba = ab$ and hence S is a left regular band. This establishes the equivalence of (1) and (3).

If S is a left regular band, then the support map $\sigma: S \rightarrow \Lambda(S)$ is a homomorphism to a semilattice with the property that $\sigma(a) = \sigma(b)$ implies $ab = a$ and $ba = b$ (as $Ba = Bb$). Therefore, (1) implies (4). If (4) holds and $a \in S$, then $aa = a$ because the fiber over $\rho(a)$ satisfies $xy = x$. Also, if $a, b \in S$, then $\rho(aba) = \rho(a)\rho(b)\rho(a) = \rho(a)^2\rho(b) = \rho(a)\rho(b) = \rho(ab)$ and so ab, aba belong to the same fiber of ρ . Thus $aba = (ab)^2a = (ab)(aba) = ab$ and so S is a left regular band. This completes the proof of the theorem. $\square$

We define here some further operations on left regular bands, inspired by topology, that will come into play later in the theory.

If B, B' are (disjoint) left regular bands, we define their *join* $B * B'$ to be $B \cup B'$ where B and B' are subsemigroups and $b'b = b = bb'$ for all $b \in B$ and $b' \in B'$. Notice that $B * B'$ is a monoid if and only if B' is a monoid. Also observe that $\Lambda(B * B') = \Lambda(B) * \Lambda(B')$ (where the join is as left regular bands).

The variety (in the sense of universal algebra [37]) of left regular bands is well known to be generated by the three-element left regular band $L = \{0, +, -\}$ with the multiplication table in Table 1. See [92, Proposition 7.3.2].

One can view L as the monoid of self-mappings of $[-1, 1]$ (acting on the left) generated by the identity map r_0 , and the constant maps $r_{\pm}$ with image ± 1 . When we come to the relationship between hyperplane face semigroups and zonotopes, described later on, this way of thinking about L will make sense because L is the face monoid of the hyperplane arrangement in $\mathbb{R}$ with the origin as the hyperplane and $[-1, 1]$ is the corresponding zonotope. In oriented matroid theory, elements of L^n are called *covectors*. Note that $\partial L = \{-, +\}$.

Define the *suspension* of a left regular band B to be $S(B) = \partial L * B$; for instance, $L = S(\{0\})$. Note that $S(B)$ is a monoid if and only if B is a monoid. Also $\Lambda(S(B)) = \Lambda(B) \cup \{-\infty\}$ where $-\infty$ is an adjoined minimum element.

We turn now to some particularly salient examples of left regular bands.

2.1. Free left regular bands and matroids. In this section we consider the free left regular band monoid $F(A)$ on a set A and a generalization to matroids.

2.1.1. Free left regular bands. The free monoid on the set A , denoted A^* , is the set of all words over the alphabet A (including the empty word) with the binary operation of concatenation. By standard universal algebra (or the adjoint functor theorem), there is a *free left regular band monoid* $F(A)$ on the set A . Let $\rho: A^* \rightarrow F(A)$ be the canonical surjective homomorphism. We claim that the repetition-free words form a cross-section to ρ , that is, they constitute a set of normal forms. Indeed, $\rho(u) = \rho(\bar{u})$ where $\bar{u}$ is the word obtained from u by removing repetitions as you scan u from left-to-right (for example, $\overline{ababcbac} = abc$) because of the identities $x^2 = x$ and $xyx = xy$ satisfied by $F(A)$.

The power set $P(A)$ is a commutative left regular band with respect to the operation of union and the mapping $c: A^* \rightarrow P(A)$ sending a word $w \in A^*$ to the set $c(w)$ of letters appearing in w is a homomorphism. Thus if u, v are words with the same image under ρ , then $c(u) = c(v)$. Thus to show that the repetition-free words are a set of normal forms, it suffices to show that if u, v are distinct repetition-free words with $c(u) = c(v)$, then there is a homomorphism $\psi: A^* \rightarrow L$ with $\psi(u) \neq \psi(v)$. (This also shows that L generates the variety of left regular bands.) Let w be the longest common prefix of u and v . Then $u = wax$ and $v = wby$ with $a \neq b$ in A and $x, y \in A^*$. Since u, v are repetition-free, the letters a and b do not appear in w . Hence if we define a homomorphism $\psi: A^* \rightarrow L$ on A by $\psi(a) = +$, $\psi(b) = -$ and $\psi(c) = 0$ for $c \in A \setminus \{a, b\}$, then we obtain that $\psi(u) = +$ and $\psi(v) = -$.

It follows that we can identify $F(A)$ with the set of repetition-free words over the alphabet A with the binary operation $u \cdot v = \overline{uv}$ where $\overline{uv}$ is the result of removing repetitions as you scan uv from left to right. The free left regular band on A is obtained from $F(A)$ by removing the empty word. If A has at least two elements, the free left regular band (semigroup) does not have a unital algebra over any base ring (as is easily verified). For this reason we are only interested in free left regular band monoids.

It is not difficult to see that $u \leq v$ in $F(A)$ if and only if v is a prefix of u and that $F(A)u \subseteq F(A)v$ if and only if $c(v) \subseteq c(u)$. Hence $\Lambda(F(A))$ can be identified with $P(A)$ ordered by reverse inclusion. With this identification if $X \subseteq A$, then $F(A)_{\geq X} = F(X)$. If $w \in F(A)$ with $c(w) = X$, then $wF(A) \cong F(A \setminus X)$. Thus every contraction and “deletion” of a free left regular band monoid is again a free left regular band monoid.

2.1.2. *Matroids.* Brown [34] generalized the free left regular band monoid to matroids. Matroids were introduced by Whitney in 1932 as an abstraction of linear independence in vector spaces, algebraic independence over fields and independence in graphs. Since then, matroid theory has expanded in many directions, with applications in geometry, topology, algebraic combinatorics and optimization theory. We give a brief introduction to matroids and the related notion of geometric lattice suited to the needs of the current work. See [117] for further details.

A *matroid* is a pair $\mathcal{M} = (E, \mathcal{F})$ where E is the *ground set* and $\mathcal{F}$ is a *hereditary* collection of subsets of E (that is, $X \in \mathcal{F}$ and $Y \subseteq X$ implies $Y \in \mathcal{F}$) satisfying the following Exchange Axiom.

(EA) If I_1 and I_2 are elements of $\mathcal{F}$ with $|I_1| < |I_2|$, then there is an $e \in I_2 - I_1$ such that $I_1 \cup \{e\} \in \mathcal{F}$.

Elements of $\mathcal{F}$ are called *independent subsets*.

A *loop* in $\mathcal{M}$ is an element $e \in E$ that belongs to no independent subset or, equivalently, $\{e\} \notin \mathcal{F}$. We will often restrict our attention to matroids that do not contain loops. Two elements e, f of E are *parallel* if they are not loops and $\{e, f\}$ is not an independent subset. This is a clear generalization of the notion of parallel vectors in a vector space. A matroid is *simple* if it contains no loops and no distinct pair of parallel elements.

The motivating examples are the matroid of all linearly independent subsets of a vector space, the matroid of all forests in a graph (thought of as sets of edges of the graph) and the matroid of all algebraically independent subsets of a field. It follows easily from the exchange axiom that all maximal independent subsets (called *bases*) in a matroid have the same size. In the three motivating examples, bases in a vector space matroid are the usual bases of linear algebra, bases in a graphical matroid are spanning forests, and bases in the field example are transcendence bases.

Let $\mathcal{M} = (E, \mathcal{F})$ be a matroid and let A be a subset of E . The *rank* of A is defined by

$$r(A) = \max\{|I| \mid I \in \mathcal{F}, I \subseteq A\}.$$

In the vector space matroid, the rank of a set of vectors is the dimension of the space that they span. The *rank* of a matroid is the cardinality of a basis; this is also $r(E)$.

The rank function, $r: 2^E \rightarrow \mathbb{N}$ is a *semimodular* rank function, meaning that it satisfies the following axioms.

- (R1) $r(A) \leq |A|$
- (R2) $A \subseteq B \implies r(A) \leq r(B)$
- (R3) *Upper Semimodularity:* $r(A \cup B) + r(A \cap B) \leq r(A) + r(B)$

One can prove that the independent sets $\mathcal{F}$ are precisely the subsets $I \subseteq E$ such that $r(I) = |I|$. Thus, the rank function gives another way to axiomatize matroids.

Flats are the matroid theoretic analogue of subspaces of a vector space. They are the maximal subsets of E of a given rank. That is, a subset A is a

flat if and only if $r(A) < r(A \cup \{e\})$ for all $e \in E \setminus A$. It can be shown that the intersection of a collection of flats is a flat.

By the usual considerations, we can define a closure operator on 2^E by sending a subset A of E to the subset $\overline{A}$ which is the intersection of all the flats containing A . More concretely,

$$\overline{A} = A \cup \{e \in E \mid r(A) = r(A \cup \{e\})\}.$$

The assignment of A to $\overline{A}$ is a closure operator on 2^E , in that it is an order preserving, idempotent and non-decreasing function. In addition, it satisfies the following version of the exchange axiom, familiar in the case of vector spaces and linear independence.

(EX) If $e, f \in E$ and $A \subseteq E$, then if $f \in \overline{A \cup \{e\}} \setminus \overline{A}$, then $e \in \overline{A \cup \{f\}}$.

It can be shown that, conversely, a closure operator $c: 2^E \rightarrow 2^E$ satisfying the Exchange Axiom (EX) defines a unique matroid where the flats are precisely those subsets A of E such that $c(A) = A$.

The collection $\Lambda(\mathcal{M})$ of all flats of a matroid $\mathcal{M}$ is a lattice where the meet operation is intersection and the join is defined by $F \vee G = \overline{F \cup G}$ for flats F and G . In fact, $\Lambda(\mathcal{M})$ is a geometric lattice. In order to define this notion, recall that an element a in a lattice Λ is an *atom* if it covers the bottom element of Λ . The lattice Λ is *ranked* or *graded* if all maximal chains between x and y in Λ , with $x < y$, have the same length. The *rank function* of a graded lattice Λ is the function $r: \Lambda \rightarrow \mathbb{N}$ where $r(x)$ is the length of a maximal chain from the bottom element of Λ to x .

Definition 2.8 (Geometric lattice). A lattice Λ is then said to be a *geometric lattice* if it satisfies the following axioms.

- (G1) Λ is an atomic lattice, that is, every element in Λ is a join of the atoms.
- (G2) Λ is ranked.
- (G3) Λ is *upper semimodular*: $r(A \vee B) + r(A \wedge B) \leq r(A) + r(B)$.

A theorem of Birkhoff [117] shows that there is a one-to-one correspondence between geometric lattices and simple matroids. The correspondence associates with a geometric lattice Λ the simple matroid $\mathcal{M}(\Lambda)$ defined as follows. The ground set of this matroid is the set E of atoms of Λ . A collection I of atoms is independent if the rank of the join of I is the cardinality of I . Then the lattice of flats of $\mathcal{M}(\Lambda)$ is isomorphic to Λ .

If $\mathcal{M}$ is a matroid with ground set E , then Brown considers the left regular band monoid whose elements are all linearly ordered independent subsets of E . The product is given by concatenation and then removing elements that are dependent on the previous elements where we say $e \in E$ is *dependent* on a subset $A \subseteq E$ if $\overline{A \cup \{e\}} = \overline{A}$. The free left regular band is obtained from the matroid in which all subsets of elements are independent. The Hasse diagram of the left regular band associated to a matroid is a rooted tree. Björner extended this construction to interval greedoids, which form a natural extension of the notion of matroids. See [23] for details.

2.2. Free partially commutative left regular bands. If $\Gamma = (V, E)$ is a (finite simple) graph, then the *free partially commutative left regular band* $B(\Gamma)$ is the left regular band *monoid* with presentation

$$\langle V \mid xy = yx, \text{ if } \{x, y\} \in E \rangle$$

(where this presentation is as a left regular band monoid). They are left regular band analogues of free partially commutative monoids [41] (or trace monoids [50]) and right-angled Artin groups (cf. [16, 118]).

The elements of $B(\Gamma)$ are in bijection with acyclic orientations of induced subgraphs of the complementary graph $\bar{\Gamma}$ [75]. Namely, if w is a word in V and W is the set of letters appearing in w , then we associate to w the induced subgraph of $\bar{\Gamma}$ on the vertex set W , acyclically oriented by orienting an edge $\{x, y\}$ from x to y if the first occurrence of x in w is *before* the first occurrence of y in w . Conversely, given an acyclically oriented induced subgraph of $\bar{\Gamma}$ we can find a repetition-free word giving rise to that subgraph by performing a topological sorting.

For example, if $E = \emptyset$, i.e., the graph Γ is edgeless, then $B(\Gamma)$ is the free left regular band monoid on the vertices of Γ and elements correspond to acyclic orientations of induced subgraphs of the complete graph on the vertex set V . But these correspond exactly to repetition-free words over the alphabet V . At the other extreme, if Γ is complete, then $\bar{\Gamma}$ is edgeless. Hence each induced subgraph has only one acyclic orientation. Thus the class of each word is determined by its support and so $B(\Gamma) \cong (P(V), \cup)$.

Like the case of free left regular band monoids, free partially commutative left regular bands behave well under contraction and “deletion.” First observe that the natural homomorphism $F(V) \rightarrow P(V)$ factors through $B(\Gamma)$ and hence $\Lambda(B(\Gamma)) \cong (P(V), \cup)$ as a semilattice via the map taking an acyclically oriented induced subgraph of the complement of Γ to its set of vertices. If $X \subseteq V$, then one easily checks that $B(\Gamma)_{\geq X} \cong B(\Gamma[X])$ where $\Gamma[X]$ denotes the induced subgraph of Γ with vertex set X . On the other hand, if $b \in B(\Gamma)$ is the image of a word w in $F(V)$ with $c(w) = X$, then $bB(\Gamma) \cong B(\Gamma[V \setminus X])$. Details can be found in [75] or verified directly.

If Γ_1, Γ_2 are graphs, then it is not difficult to check that $B(\Gamma_1) \times B(\Gamma_2) \cong B(\Gamma_1 * \Gamma_2)$, where $\Gamma_1 * \Gamma_2$ is the join of the two graphs. In other words, $\Gamma_1 * \Gamma_2$ is the graph obtained by connecting each vertex of Γ_1 to each vertex of Γ_2 by an edge.

Free partially commutative left regular bands turn out to give a simple model of a family of random walks considered by Athanasiadis and Diaconis [7] and their cohomology turns out to be connected to classical invariants of the clique (or flag) complex of Γ . More details on free partially commutative left regular bands can be found in [75].

2.3. Hyperplane arrangements and oriented matroids. We follow here Brown [34, 35], Brown and Diaconis [36] and Björner *et al.* [26]. See

also [1]. A *hyperplane arrangement* in a real vector space V is a finite collection $\mathcal{A} = \{H_1, \dots, H_n\}$ of (affine) hyperplanes in V . The arrangement is called *central* $\bigcap_{i=1}^n H_i \neq \emptyset$ and otherwise is called *affine*. Without loss of generality, in the case of a central arrangement, we may (and do) assume that the hyperplanes are linear (that is, pass through the origin). A hyperplane arrangement in V is said to be *essential* if the unit normals to the hyperplanes span V . A central arrangement is essential if and only if the intersection of all the hyperplanes is the origin. We can always assume a central arrangement is essential by considering $V / \bigcap_{i=1}^n H_i$. Similarly, we may restrict our attention to essential affine arrangements without loss of generality. See [111] for details. The connected components of $V \setminus \bigcup_{i=1}^n H_i$ are called *chambers*.

2.3.1. Central arrangements and oriented matroids. For the moment we consider the case that $\mathcal{A}$ is central and we assume that it is given by linear forms $f_i = 0$, for $i = 1, \dots, n$, with $f_1, \dots, f_n \in V^*$. If $x \in \mathbb{R}$, put

$$\operatorname{sgn}(x) = \begin{cases} +, & \text{if } x > 0 \\ -, & \text{if } x < 0 \\ 0, & \text{if } x = 0. \end{cases}$$

Then we can define a mapping $\theta: V \longrightarrow L^n$ by

$$\theta(x) = (\operatorname{sgn}(f_1(x)), \dots, \operatorname{sgn}(f_n(x))). \quad (2.1)$$

It turns out that the set of covectors $\theta(V)$ is a submonoid of L^n , introduced independently by Tits [113, 114] and Bland. Trivially, we have $\theta(x)\theta(x) = \theta(x)$. If $x \neq y$, then $\theta(x)\theta(y) = \theta(z)$ where z is an element on the open line segment (x, y) such that $f_i(z) = f_i(x)$ whenever $f_i(x) \neq 0$. (Such a z exists because $f_1, \dots, f_n$ are continuous so we can find $\varepsilon > 0$ such that for each i with $f_i(x) \neq 0$ we have that $\operatorname{sgn}(f_i(B_\varepsilon(x))) = \operatorname{sgn}(f_i(x))$.) Note that $\theta(0) = 0$ (the identity). The fibers of θ are relative interiors of polyhedral cones and the fibers over elements of $\theta(V) \cap (\partial L)^n$ are the chambers. Define $\mathcal{F}(\mathcal{A}) = \theta(V)$ to be the *face monoid* of $\mathcal{A}$. Note that up to isomorphism it does not depend on the choice of the f_i . Being a submonoid of L^n , the monoid $\mathcal{F}(\mathcal{A})$ is a left regular band. Figure 1 gives an example of the covectors associated to a hyperplane arrangement in $\mathbb{R}^2$.

Associated to the arrangement $\mathcal{A}$ is the Minkowski sum

$$\begin{aligned} Z(\mathcal{A}) &= [-f_1, f_1] + \dots + [-f_n, f_n] \\ &= \left\{ \sum_{i=1}^n t_i f_i \mid -1 \leq t_i \leq 1, \text{ for } i = 1, \dots, n \right\}, \end{aligned}$$

which is a convex polytope known as the *zonotope polar to $\mathcal{A}$* . We refer the reader to [122] for a good source on polytopes and, in particular, zonotopes. Formally, a *zonotope* is an image of a hypercube under an affine map or,




FIGURE 1. The covectors of the faces of the hyperplane arrangement in $\mathbb{R}^2$ consisting of three distinct lines.

equivalently, a Minkowski sum of line segments. If $\sigma = (\sigma_1, \dots, \sigma_n) \in L^n$ is any covector, then we can associate to it the zonotope

$$Z_\sigma = \sum_{\sigma_i=0} [-f_i, f_i] + \sum_{\sigma_i=-} \{-f_i\} + \sum_{\sigma_i=+} \{f_i\}. \quad (2.2)$$

Clearly, $Z_\sigma \subseteq Z(\mathcal{A}) = Z_0$. It turns out that the faces of $Z(\mathcal{A})$ are exactly the zonotopes Z_σ with $\sigma \in \mathcal{F}(\mathcal{A})$ and that $\sigma \leq \tau$ if and only if $Z_\sigma \leq Z_\tau$. Consequently, $\mathcal{F}(\mathcal{A})$ is isomorphic, as a poset, to the face poset of $Z(\mathcal{A})$. Indeed, if $x \in (V^*)^* = V$, then the face of $Z(\mathcal{A})$ on which x (viewed as a functional) is maximized is $Z_{\theta(x)}$. See [26, Proposition 2.2.2] for details. Notice that the vertices of the zonotope are in bijection with the chambers of the arrangement.

The *intersection lattice* $\mathcal{L}(\mathcal{A})$ is the poset of subspaces which are intersections of hyperplanes from $\mathcal{A}$, ordered by reverse inclusion. It is a geometric lattice [111]. One can verify directly that $\Lambda(\mathcal{F}(\mathcal{A})) \cong \mathcal{L}(\mathcal{A})$ and the support map takes a covector $x \in \mathcal{F}(\mathcal{A})$ to the linear span of the fiber $\theta^{-1}(x)$. See [34, 35] for details.

Some important examples are the following. The *boolean arrangement* is the arrangement $\mathcal{B}_n$ of coordinate hyperplanes $x_i = 0$ in $\mathbb{R}^n$. The zonotope $Z(\mathcal{A})$ is the n -cube $[-1, 1]^n$. One has $\mathcal{F}(\mathcal{B}_n) = L^n$. Figure 2 depicts the zonotope polar to the hyperplane arrangement in Figure 1.

If Γ is a simple graph (say with vertex set $[n]$), then the *graphical arrangement* $\mathcal{A}(\Gamma)$ associated to Γ is defined by the hyperplanes

$$H_{ij} = \{x \in \mathbb{R}^n \mid x_i - x_j = 0\}$$

with $\{i, j\}$ an edge of Γ and $j > i$ (note that this arrangement is not essential). The special case $\mathcal{A}(K_n)$ of the complete graph is known as the *braid arrangement*. The zonotope corresponding to the braid arrangement is the




FIGURE 2. The zonotope polar to the arrangement in Figure 1.

permutohedron, which is the convex hull of the vectors

$$\{(\sigma(1), \dots, \sigma(n)) \mid \sigma \in S_n\}.$$

The braid arrangement is the Coxeter arrangement associated to S_n , viewed as a Coxeter group. More generally, if W is any finite Coxeter group, there is a corresponding reflection arrangement $\mathcal{A}_W$. One can obtain the associated zonotope, called the W -*permutohedron*, $Z(\mathcal{A}_W)$ by choosing a point x in some chamber and then taking the convex hull of the orbit Wx of x . The 1-skeleton of $Z(\mathcal{A}_W)$ is the Cayley graph of W with respect to its Coxeter generators.

The sets of covectors of a central hyperplane arrangement are the primary examples of oriented matroids. The reader is referred to the treatise [26] for a definitive account of the theory of oriented matroids, including all facts that we state below. We need some notation to define this notion. There is a natural automorphism of L that fixes 0 and switches $+$, $-$, which we write $x \mapsto -x$. Thus L can be viewed as a unary monoid and hence L^n can be viewed as a unary monoid via

$$(x_1, \dots, x_n) \mapsto -(x_1, \dots, x_n) = (-x_1, \dots, -x_n).$$

For covectors $x, y \in L^n$, define their *separation set* to be

$$S(x, y) = \{i \in [n] \mid x_i = -y_i \neq 0\}$$

that is, it is the set of indices where x, y have opposite signs. The following properties are enjoyed by the set $\mathcal{F}(\mathcal{A})$ of covectors of a central hyperplane arrangement and form the set of axioms for an oriented matroid.

Definition 2.9 (Oriented matroid). A set of $\mathcal{L}$ of covectors in $\{+, -, 0\}^E$ is said to form an *oriented matroid* (with (finite) *ground set* E) if it satisfies: (OM0) $0 \in \mathcal{L}$;

- (OM1) $x \in \mathcal{L}$ implies $-x \in \mathcal{L}$;
- (OM2) $x, y \in \mathcal{L}$ implies $xy \in \mathcal{L}$;
- (OM3) *Strong Elimination*: $x, y \in \mathcal{L}$ and $e \in S(x, y)$ implies there exists $z \in \mathcal{L}$ such that $z_e = 0$ and $z_f = (xy)_f = (yx)_f$ for all $f \notin S(x, y)$.

Sometimes we say that the oriented matroid is the pair $(E, \mathcal{L})$ to emphasize that the ground set is really part of the data.

Note that (OM0)–(OM2) say that $\mathcal{L}$ is a unary submonoid of L^E . The face monoid $\mathcal{F}(\mathcal{A})$ of a central hyperplane arrangement is a unary submonoid because $\theta(-x) = -\theta(x)$. To see that $\mathcal{F}(\mathcal{A})$ satisfies (OM3), if $e \in S(x, y)$ and $x = \theta(x')$, $y = \theta(y')$, then the line segment $[x', y']$ will intersect the hyperplane corresponding to e at some point z' . Putting $z = \theta(z')$, we have $z_e = 0$, and if $f \notin S(x, y)$, then $z_f = (xy)_f = (yx)_f$. An oriented matroid is said to be *realizable* if it comes from a hyperplane arrangement. Not all oriented matroids are realizable (in fact, in some sense most are not).

Let us introduce some oriented matroid terminology and compare it with semigroup terminology. Let $\mathcal{L} \subseteq L^E$ be an oriented matroid. Let us warn the reader that the oriented matroid literature customarily uses the reverse of the order that we have been considering on left regular bands. However, we shall stick to using the order $x \leq y$ if $yx = x$. With our convention, maximal non-identity elements of $\mathcal{L}$ are called *cocircuits* and the elements of the minimal ideal (that is, the minimal elements) are called *topes* (in hyperplane theory, topes are called *chambers*). The set of all topes is denoted by $\mathcal{T}$. It is known that the cocircuits generate $\mathcal{L}$ as a monoid and that

$$\mathcal{L} = \{x \in L^E \mid x\mathcal{T} \subseteq \mathcal{T}\};$$

see the dual of [26, Proposition 3.7.2] and [26, Theorem 4.2.13]. In particular, the set of topes determines the oriented matroid.

If $x \in \mathcal{L}$, the *zero set* of x is

$$Z(x) = \{e \in E \mid x_e = 0\}$$

The mapping $Z: \mathcal{L} \rightarrow (P(E), \cap)$ given by $x \mapsto Z(x)$ is a monoid homomorphism whose corestriction $Z: \mathcal{L} \rightarrow Z(\mathcal{L})$ can be identified with the support map $\sigma: \mathcal{L} \rightarrow \Lambda(\mathcal{L})$. One has that $\Lambda(\mathcal{L})$ is a geometric lattice and hence the lattice of flats of a unique simple matroid. The *rank* of $\mathcal{L}$ is defined to be the rank of the corresponding matroid. When $\mathcal{L}$ is the set of covectors associated to a central hyperplane arrangement $\mathcal{A}$, one has that the support lattice $\Lambda(\mathcal{F}(\mathcal{A}))$ is isomorphic to the intersection lattice of $\mathcal{A}$ [35, 111] and the rank is the codimension of the intersection of the hyperplanes in $\mathcal{A}$. We remark that there is a notion of isomorphism of oriented matroids and an unpublished result of the third author and Hugh Thomas shows that it coincides with isomorphism of the corresponding unary left regular band monoids [112].

If $A \subseteq E$, then the *contraction* of $(E, \mathcal{L})$ at A is the oriented matroid with ground set $E \setminus A$ and whose set $\mathcal{L}/A$ of covectors is $\{x|_{E \setminus A} \mid A \subseteq Z(x)\}$.

This monoid is isomorphic to the contraction $\mathcal{L}_{\geq \overline{A}}$ where

$$\overline{A} = \bigcap \{B \in Z(\mathcal{L}) \mid B \supseteq A\}$$

(under the identification of $\Lambda(\mathcal{L})$ with the zero sets of elements of $\mathcal{L}$). This is why we use the terminology contraction in this context for left regular bands in general. In the case of an oriented matroid $(E, \mathcal{L})$ coming from a hyperplane arrangement $\mathcal{A}$, one has that $\mathcal{L}/A$ is the face monoid of the hyperplane arrangement $\mathcal{A}/A$ whose underlying vector space W is the intersection of the hyperplanes coming from A and whose hyperplanes are the proper intersections of hyperplanes from $E \setminus A$ with W .

If $(E, \mathcal{L})$ is an oriented matroid and $A \subseteq E$, then the *deletion* of A from $(E, \mathcal{L})$ is the oriented matroid with ground set $E \setminus A$ and whose set $\mathcal{L} \setminus A$ of covectors is $\{x|_{E \setminus A} \mid x \in \mathcal{L}\}$. If $x \in \mathcal{L}$ with $E \setminus Z(x) = A$, then one easily checks that $x\mathcal{L} \cong \mathcal{L} \setminus A$, cf. the discussion after [26, Lemma 4.1.8]. For hyperplane arrangements, the operation of deletion corresponds to removing a subset of the hyperplanes.

An element $e \in E$ is called a *loop* of $(E, \mathcal{L})$ if $x_e = 0$ for all $x \in \mathcal{L}$ or, equivalently, if $T_e = 0$ for each tope T . If $e \in E$ is not a loop, then there is an associated partition of $\mathcal{T}$ into *half-spaces*. Namely, we have

$$\mathcal{T}_e^+ = \{T \in \mathcal{T} \mid T_e = +\} \quad \text{and} \quad \mathcal{T}_e^- = \{T \in \mathcal{T} \mid T_e = -\}.$$

For hyperplane arrangements these correspond to the sets of chambers in the positive and negative half-spaces determined by the hyperplane corresponding to e .

The reader should consult [26] for more details and motivation for studying oriented matroids. In this paper, we mostly need results and notions from [26, Chapter 4]. However, in the next subsection we will explain how zonotopal tilings of a zonotope correspond to certain oriented matroids. In particular, we will need the following notion. Let $(E, \mathcal{L})$ be an oriented matroid. Then an oriented matroid $(E \cup \{g\}, \mathcal{L}')$ is a *single-element lifting* of $(E, \mathcal{L})$ if g is not a loop of $\mathcal{L}'$ and $\mathcal{L}'/\{g\} = \mathcal{L}$.

2.3.2. Affine arrangements, affine oriented matroids, T -convex sets of topes and zonotopal tilings. Next we turn to affine hyperplane arrangements and affine oriented matroids. Suppose that $\mathcal{A} = \{H_1, \dots, H_n\}$ is an affine hyperplane arrangement in $V = \mathbb{R}^d$ given by equations $f_i(x) = c_i$ with $f_i \in V^*$ and $c_i \in \mathbb{R}$ for $i = 1, \dots, n$. See Figure 3. Then one can define a mapping $\theta: V \rightarrow L^n$ by

$$\theta(x) = (\text{sgn}(f_1(x) - c_1), \dots, \text{sgn}(f_n(x) - c_n))$$

and obtain a subsemigroup of L^n . It will not be a monoid for a non-central arrangement. It turns out, however, to be more convenient to describe affine hyperplane face semigroups via central arrangements.




FIGURE 3. An affine arrangement

To do this, we embed V in $\mathbb{R}^{d+1}$ as the affine hyperplane $x_{d+1} = 1$. Now let H'_i be the hyperplane in $\mathbb{R}^{d+1}$ defined by

$$f'_i(x_1, \dots, x_{d+1}) = f_i(x_1, \dots, x_d) - x_{d+1}c_i$$

and observe that $H'_i \cap \{x \in \mathbb{R}^{d+1} \mid x_{d+1} = 1\} = \{(x, 1) \mid x \in H_i\}$. Let $H = \{x \in \mathbb{R}^{d+1} \mid x_{d+1} = 0\}$ and put $\mathcal{A}' = \{H'_1, \dots, H'_n, H\}$. We now define the *hyperplane face semigroup* of $\mathcal{A}$ to be $\mathcal{F}(\mathcal{A}) = \{x \in \mathcal{F}(\mathcal{A}') \mid x_{n+1} = +\}$. The reader should verify that this gives an isomorphic semigroup to the one constructed in the previous paragraph. The support semilattice of $\mathcal{F}(\mathcal{A})$ is isomorphic to the intersection semilattice of $\mathcal{A}$ [111] (that is the poset, of all non-empty intersections of hyperplanes of $\mathcal{A}$ ordered by reverse inclusion). Notice that $\mathcal{F}(\mathcal{A})$ is a right ideal in $\mathcal{F}(\mathcal{A}')$.

An *affine oriented matroid* consists of a triple $(E, \mathcal{L}, g)$ where $\mathcal{L}$ is an oriented matroid on E and $g \in E$ is not a loop. The corresponding left regular band is

$$\mathcal{L}^+(g) = \{x \in \mathcal{L} \mid x_g = +\}.$$

For example, each affine hyperplane arrangement $\mathcal{A}$ determines an affine oriented matroid $([n+1], \mathcal{F}(\mathcal{A}'), n+1)$ in the manner described above. Affine oriented matroids play a role in linear programming over oriented matroids [26, Chapter 10]. See [26, Section 4.5] for more on affine oriented matroids.

Affine oriented matroids are closely related to zonotopal tilings via the Bohne-Dress theorem. We follow here Richter-Gebert and Ziegler [93]. Let $\mathcal{A}$ be a hyperplane arrangement with associated oriented matroid $\mathcal{F}(\mathcal{A})$ and zonotope $Z(\mathcal{A})$. A (simple) *zonotopal tiling* of $Z(\mathcal{A})$ is a collection $\mathcal{Z} = \{Z_1, \dots, Z_m\}$ of zonotopes such that:

- (ZT1) $Z(\mathcal{A}) = \bigcup_{i=1}^m Z_i$;
- (ZT2) each face of an element of $\mathcal{Z}$ belongs to $\mathcal{Z}$;
- (ZT3) if $Z_i, Z_j \in \mathcal{Z}$, then $Z_i \cap Z_j$ is a common face of Z_i, Z_j ;
- (ZT4) every edge of $Z(\mathcal{A})$ belongs to $\mathcal{Z}$.

Notice that (ZT1)–(ZT3) just assert that $\mathcal{Z}$ is the structure of a polyhedral complex on $Z(\mathcal{A})$. The Bohné-Dress theorem asserts that zonotopal tilings of $Z(\mathcal{A})$ are in bijection with single-element liftings of $\mathcal{F}(\mathcal{A})$. If $(E \cup \{g\}, \mathcal{L})$ is a single element lifting of $\mathcal{F}(\mathcal{A})$, then the corresponding zonotopal tiling is $\mathcal{Z} = \{Z_\sigma \mid (\sigma, +) \in \mathcal{L}^+(g)\}$ where Z_σ is defined as in (2.2). Hence the faces of the polyhedral complex $\mathcal{Z}$ can be identified with the elements of the left regular band $\mathcal{L}^+(g)$ associated to the affine oriented matroid $(E \cup \{g\}, \mathcal{L}, g)$. The Bohné-Dress theorem is a good motivation for studying oriented matroids even when one is only interested in zonotopes. (It is known that not all zonotopal tilings come from realizable single-element extensions.)

The construction of affine oriented matroids admits the following generalization. A set of topes of an oriented matroid $(E, \mathcal{L})$ is called *T-convex* if it is an intersection of half-spaces (in the sense defined earlier). For example, the topes belonging to $\mathcal{L}^+(g)$ are precisely those in $\mathcal{T}_g^+$, and hence form a *T-convex* set. In [34], *T-convex* sets of topes for a central hyperplane arrangement are called convex sets of chambers. Up to reorientation of the oriented matroid, we may assume without loss of generality that we are considering an intersection of positive half-spaces. So let $A \subseteq E$ be a subset, none of whose elements are loops. Put $\mathcal{L}^+(A) = \bigcap_{e \in A} \mathcal{L}^+(e)$. Note that $\mathcal{L}^+(\emptyset) = \mathcal{L}$. If non-empty, then we call $\mathcal{L}^+(A)$ the *face semigroup* of the *T-convex* set of topes $\bigcap_{e \in A} \mathcal{T}_e^+$. Random walks on these semigroups, in the case of hyperplane arrangements, were considered by Brown [34]. Details on *T-convex* sets can be found in [26, Section 4.2].

2.4. Strong elimination systems, lopsided systems and COMs. As we were preparing this paper for submission, the paper [13] appeared on ArXiv that includes a large source of new examples of left regular bands and, in particular, includes all the examples of the previous subsection.

The weakest notion considered in [13] is that of a strong elimination system.

Definition 2.10 (Strong elimination system). A *strong elimination system* $(E, \mathcal{L})$ with *ground set* E is a collection $\mathcal{L} \subseteq L^E$ of covectors satisfying (OM2) and (OM3) from Definition 2.9.

The notion of a strong elimination system abstracts the following setup. A *realizable strong elimination system* consists of a central hyperplane arrangement $\mathcal{A}$ in $\mathbb{R}^d$ and a convex subset $C \subseteq \mathbb{R}^d$. It is convenient to assume that the strong elimination system is *essential* meaning that C spans $\mathbb{R}^d$ and that every hyperplane in $\mathcal{A}$ passes through C . This assumption does not change the semigroups one can obtain. Again, we choose forms defining the hyperplane arrangement to obtain a mapping $\theta: \mathbb{R}^d \rightarrow \mathbb{L}^E$, as per (2.1), where E is the set of hyperplanes. Then $\theta(C)$ is a subsemigroup of the face monoid $\mathcal{F}(\mathcal{A})$ because if $x, y \in C$, then the line segment $[x, y]$ from x to y is contained in C and hence the element $z \in \mathbb{R}^n$ with $\theta(x)\theta(y) = \theta(z)$ belongs

to C . Also, if $x, y \in C$ and H is a hyperplane with x, y on opposite sides of H , then the intersection z of $[x, y]$ with H belongs to C and so $\theta(C)$ satisfies (OM3). We remark that one could replace central arrangements with affine ones, but one obtains nothing new since our embedding of face semigroups of affine arrangements as right ideals in face monoids of central arrangements shows that the strong elimination system obtained from intersecting a convex set with the faces of an affine arrangement can also be obtained from intersecting a convex set with a central arrangement in a Euclidean space of one larger dimension.

Lawrence introduced the notion of a lopsided system in [68], which has applications in statistics, combinatorics, learning theory and computational geometry; see [11, 12, 79]. A *lopsided system* $(E, \mathcal{L})$ with ground set E is a right ideal in $\mathcal{L} \subseteq L^E$ satisfying (OM3). It turns out that CAT(0) cube complexes are examples of lopsided systems, which we essentially proved before we became aware of the notion of a lopsided system, and which was proved explicitly and independently in [13]. Lopsided systems abstract the strong elimination systems obtained by considering the regions obtained by intersecting orthants (i.e., faces of the Boolean arrangement) with an open convex set in Euclidean space.

More generally, you can obtain a right ideal of a hyperplane face monoid satisfying (OM3) by considering those faces of a hyperplane arrangement $\mathcal{A}$ in $\mathbb{R}^d$ intersecting a convex open subset C in $\mathbb{R}^d$. More precisely, following [13], a *realizable COM*¹ is a set of covectors of a realizable strong elimination system corresponding to a central hyperplane arrangement $\mathcal{A}$ in $\mathbb{R}^d$ and a convex *open* subset C of $\mathbb{R}^d$ (and again we may assume it is *essential*, meaning that each hyperplane of $\mathcal{A}$ meets C). The right ideal property follows because if $x \in C$ and $y \in \mathbb{R}^d$, then any point z on $[x, y]$ sufficiently closed to x belongs to C and hence we may choose $z \in C$ with $\theta(x)\theta(y) = \theta(z)$ (by choosing z in an ε -ball around x small enough so that $z \in C$ and is in the same open half-space containing x associated to each hyperplane of $\mathcal{A}$ not containing x). Note that any face semigroup of an affine hyperplane arrangement is a realizable COM. Indeed, our embedding of the face semigroup of an affine arrangement in $\mathbb{R}^d$ into the face monoid of a central hyperplane arrangement in $\mathbb{R}^{d+1}$ consists of taking the covectors of elements in $\mathbb{R}^{d+1}$ belonging to the open convex set which is the positive half-space with $x_{d+1} > 0$ in the homogenization of the affine arrangement. As was the case for strong elimination systems, there is no extra generality obtained by considering the faces of an affine arrangement intersecting a convex open subset.

Let us now give the formal definition of a COM from [13].

¹COM is used in [13] as an acronym for both “complex of oriented matroids” and “conditional oriented matroid.”

Definition 2.11 (COM). A COM with ground set E is defined in [13] to be a pair $(E, \mathcal{L})$ with $\mathcal{L} \subseteq L^E$ such that $\mathcal{L}$ satisfies (OM3) of Definition 2.9 and the *face symmetry axiom* (FS):

(FS) $x, y \in \mathcal{L}$ implies $x(-y) \in \mathcal{L}$.

It is obvious that a right ideal $\mathcal{L}$ in an oriented matroid satisfies (FS). In particular, every lopsided system is a COM. It is apparently an open problem whether every COM is a right ideal in an oriented matroid. It is observed in [13] that every COM is a strong elimination system and hence a left regular band.

Proposition 2.12. *If $(E, \mathcal{L})$ is a COM, then $\mathcal{L}$ is a subsemigroup of L^E .*

Proof. If $x, y \in \mathcal{L}$, then $x(-y) \in \mathcal{L}$ and hence $x(-[x(-y)]) = x((-x)y) = [x(-x)]y = xy$ belongs to $\mathcal{L}$. $\square$

Affine oriented matroids and T -convex sets of topes are examples of COMs.

Proposition 2.13. *Every affine oriented matroid or T -convex set of topes is a COM.*

Proof. As any right ideal in an oriented matroid satisfies (FS), the only issue is to prove that (OM3) holds. The key point is that if $x_g = + = y_g$ and z is an element with $z_g = (xy)_g = (yx)_g$, then $z_g = +$. Thus the set of covectors of an affine oriented matroid or T -convex set of topes will inherit (OM3) from the ambient oriented matroid. $\square$

Elements of the minimal ideal of a COM are called *topes*. The following beautiful example of a COM associated to a poset is from [13].

Example 2.14 (Ranking COM of a poset). Let $(P, \preceq)$ be a poset. Without loss of generality we may assume that the underlying set of P is $\{1, \dots, n\}$. Consider then the braid arrangement $\mathcal{A}$ in $\mathbb{R}^n$ defined by the hyperplanes

$$H_{ij} = \{x \in \mathbb{R}^n \mid x_i = x_j\}$$

with $1 \leq i < j \leq n$. Let $C \subseteq \mathbb{R}^n$ be the convex open set defined by $x_i < x_j$ if $i \prec j$. The corresponding realizable COM $\mathcal{R}(P)$ is called the *ranking COM* associated to P [13]. The topes of $\mathcal{R}(P)$ are the linear extensions of P [13]. In fact, $\mathcal{R}(P)$ provides a regular CW decomposition of the order polytope [109] of P ; see [13] for details and some worked out examples.

If we view $\mathcal{F}(\mathcal{A})$ as consisting of all ordered partitions $(P_1, \dots, P_r)$ of $\{1, \dots, n\}$ (as per [17] or Example 8.10 below), then $\mathcal{R}(P)$ is the right ideal of $\mathcal{F}(\mathcal{A})$ consisting of those ordered partitions satisfying $i \prec j$ implies the block of i comes before the block of j .

An important fact is that every contraction of a COM is again a COM and every deletion is an oriented matroid [13, Section 3].

Proposition 2.15. *Let $(E, \mathcal{L})$ be a COM. Then each contraction $\mathcal{L}_{\geq X}$ with $X \in \Lambda(\mathcal{L})$ is a COM. Moreover, if $x \in \mathcal{L}$, then $x\mathcal{L}$ is isomorphic to the monoid of covectors of an oriented matroid.*

2.5. Complex hyperplane arrangements. In [27], Björner and Ziegler associated covectors to complex hyperplane arrangements. Björner later observed that these sets of covectors are left regular band monoids [23]. Ziegler [121] generalized the notion of oriented matroid to complex oriented matroids (which again are left regular band monoids) and our results should also apply in this context, but we did not work out the details.

Let $\tilde{L} = \{0, +, -, i, j\}$ with the multiplication table in Table 2. The Hasse diagram of $\tilde{L}$ is drawn in Figure 4. Note that $\tilde{L} = S(L)$, the suspension of L as defined above.

	0	+	-	i	j
0	0	+	-	i	j
+	+	+	+	i	j
-	-	-	-	i	j
i	i	i	i	i	i
j	j	j	j	j	j

TABLE 2. The multiplication table of $\tilde{L}$.




FIGURE 4. The Hasse diagram of $\tilde{L}$

Elements of $\tilde{L}^n$ will again be called *covectors*. To define the covectors of a complex hyperplane arrangement, we must first associate elements of $\tilde{L}$ to complex numbers. Define a function $\text{sgn}: \mathbb{C} \rightarrow \tilde{L}$ by

$$\text{sgn}(x + iy) = \begin{cases} i, & \text{if } y > 0, \\ j, & \text{if } y < 0, \\ +, & \text{if } y = 0, x > 0, \\ -, & \text{if } y = 0, x < 0, \\ 0, & \text{if } x = 0 = y \end{cases}$$




FIGURE 5. Complex “signs”

as depicted in Figure 5.

A (central) *complex hyperplane arrangement* in $\mathbb{C}^d$ is a collection $\mathcal{A} = \{H_1, \dots, H_n\}$ of (complex) codimension one subspaces, which we take to be the zero sets of complex linear forms $f_1, \dots, f_n$ on $\mathbb{C}^d$. We may always assume that the arrangement is *essential*, that is, $H_1 \cap \dots \cap H_n = \{0\}$. Analogously to the case of real hyperplane arrangements, we can define a map $\tau: \mathbb{C}^d \rightarrow \tilde{L}^n$ given by

$$\tau(z_1, \dots, z_d) = (\operatorname{sgn}(f_1(z_1)), \dots, \operatorname{sgn}(f_d(z_d))).$$

The image $\mathcal{F}(\mathcal{A}) = \tau(\mathbb{C}^d)$ is a submonoid of $\tilde{L}^n$ called the *face monoid* of $\mathcal{A}$. The minimal ideal of $\mathcal{F}(\mathcal{A})$ consists of the elements of $\mathcal{F}(\mathcal{A}) \cap \{i, j\}^n$, cf. [23, Proposition 3.1]. It is shown in [27, Theorem 3.5] that the right ideal $\mathcal{F}(\mathcal{A}) \cap (\partial \tilde{L})^n$ of $\mathcal{F}(\mathcal{A})$ is the face poset of a regular CW complex that is homotopy equivalent to the complement $\mathbb{C}^d \setminus (H_1 \cup \dots \cup H_n)$.

The *augmented intersection lattice* $\mathcal{L}(\mathcal{A})$ of $\mathcal{A}$ is the collection of all intersections of elements of

$$\mathcal{A}_{\text{aug}} = \{H_1, \dots, H_n, H_1^{\mathbb{R}}, \dots, H_n^{\mathbb{R}}\}$$

ordered by reverse inclusion. Here

$$H_i^{\mathbb{R}} = \{z \in \mathbb{C}^d \mid \operatorname{sgn}(f_i(z)) \in \{0, +, -\}\},$$

which is a real hyperplane defined by $\Im(f_i(z)) = 0$. One has that $\mathcal{L}(\mathcal{A})$ is the support lattice of $\mathcal{F}(\mathcal{A})$ and that the support map takes $F \in \mathcal{F}(\mathcal{A})$ to the intersection of all elements of $\mathcal{A}_{\text{aug}}$ containing $\tau^{-1}(F)$ [23, Proposition 3.3]. The lattice $\mathcal{L}(\mathcal{A})$ is a semimodular lattice of length $2d$ [23, Proposition 3.2]. More generally, if $X < Y$ in $\mathcal{L}(\mathcal{A})$, then the length of the longest chain from X to Y in $\mathcal{L}$ is $\dim_{\mathbb{R}} X - \dim_{\mathbb{R}} Y$.

It is not true in general that contractions of the left regular band $\mathcal{F}(\mathcal{A})$ along elements of $\mathcal{L}(\mathcal{A})$ are again complex hyperplane arrangements, but

their underlying posets are always face posets of regular (even PL) cell decompositions of a ball. See [75, Section 4.9] for details.

3. REGULAR CW COMPLEXES AND CW POSETS

The reader is referred to [26, Section 4.7] and [1, Appendix A.2] for basics about regular CW complexes and CW posets. See also [21] and [77, Chapter IX]. In this text all posets and CW complexes are finite unless otherwise stated.

3.1. Simplicial complexes. A *simplicial complex* K is a pair $(V, \mathcal{F})$ where V is a (finite) set of *vertices* and $\mathcal{F}$ is a collection of non-empty subsets of V such that:

- $\emptyset \neq \sigma \subseteq \tau \in \mathcal{F} \implies \sigma \in \mathcal{F}$;
- $V = \bigcup \mathcal{F}$.

Elements of $\mathcal{F}$ are called *simplices* or *faces*. More precisely, if $|\sigma| = q + 1$, then σ is called a q -*simplex* and we write $\dim \sigma = q$. Maximal faces are called *facets*. A simplicial complex K is said to be *pure* if all its facets have the same dimension. The dimension $\dim K$ of K is the maximum dimension of a facet of K .

If σ is a simplex of $K = (V, \mathcal{F})$, then the *link* of σ is the simplicial complex $\text{lk}_K(\sigma)$ with vertex set $W = \{v \in V \setminus \sigma \mid \{v\} \cup \sigma \in \mathcal{F}\}$ and with simplices all subsets $\tau \subseteq W$ such that $\sigma \cup \tau \in \mathcal{F}$. For example, if v is a vertex, then $\text{lk}_K(v)$ has vertex set the neighbors of v . A subset τ of neighbors of v is a simplex in $\text{lk}_K(v)$ if and only if $\tau \cup \{v\}$ is a simplex of K .

Let $\mathbb{k}$ be a commutative ring with unit. A simplicial complex K is called *$\mathbb{k}$ -Cohen-Macaulay* if, for all simplices σ of K , the reduced homology of $\text{lk}_K(\sigma)$ satisfies

$$\tilde{H}_q(\text{lk}_K(\sigma); \mathbb{k}) = 0$$

for all $q < \dim \text{lk}_K(\sigma)$. When $\mathbb{k}$ is a field, this is equivalent to the Stanley-Reisner ring of K over $\mathbb{k}$ being Cohen-Macaulay, whence the name. See [110] for details. If K is $\mathbb{Z}$ -Cohen-Macaulay, we shall just say that K is *Cohen-Macaulay*. By the universal coefficient theorem, K is Cohen-Macaulay if and only if K is $\mathbb{k}$ -Cohen-Macaulay for all fields $\mathbb{k}$.

The *geometric realization* of a simplicial complex $K = (V, \mathcal{F})$ is denoted $\|K\|$. By definition $\|K\|$ is the subspace of $\mathbb{R}^V$ consisting of all non-negative mappings $f: V \rightarrow \mathbb{R}$ such that $\sum_{v \in V} f(v) = 1$ and whose support is a simplex of K . Notice that the vertex of $\|K\|$ corresponding to a vertex $v \in V$ is the mapping $\delta_v: V \rightarrow \mathbb{R}$ with

$$\delta_v(x) = \begin{cases} 1, & \text{if } x = v \\ 0, & \text{else.} \end{cases}$$

A *subcomplex* of $K = (V, \mathcal{F})$ is a simplicial complex $(V', \mathcal{F}')$ with $V' \subseteq V$ and $\mathcal{F}' \subseteq \mathcal{F}$. If $W \subseteq V$, then the subcomplex of K *induced* by W is $K[W] = (W, \mathcal{F} \cap P(W))$, that is, the vertex set of $K[W]$ is W and a subset

of W is a simplex of $K[W]$ if and only if it is a simplex of K . It is the largest subcomplex of K whose vertex set is W . The q -skeleton of K is the subcomplex K^q consisting of all simplices of K of dimension at most q .

A *flag complex* is a simplicial complex $K = (V, \mathcal{F})$ such that whenever the 1-skeleton of a simplex belongs to K , then so does the simplex. Formally, this means that the minimal non-empty subsets of V that do not belong to $\mathcal{F}$ have size two.

Let us consider some important examples of simplicial complexes.

The *order complex* $\Delta(P)$ of a poset P is the simplicial complex whose vertex set is P and whose simplices are the chains in P .

If $\Gamma = (V, E)$ is a (simple) graph, the *clique complex* of Γ is the simplicial complex $\text{Clique}(\Gamma)$ whose vertex set is V and whose simplices are the cliques (subsets of vertices inducing complete subgraphs) of Γ . It is the unique flag complex whose 1-skeleton is Γ .

A *simplicial map* $f: K \rightarrow L$ of simplicial complexes is a map between their vertex sets such that the image of a simplex of K under f is a simplex of L . Simplicial complexes and simplicial maps form a category.

If $K = (V, \mathcal{F})$ is a simplicial complex and $v \notin V$, then the *cone* on K with *cone point* v is the simplicial complex

$$K * v = (V \cup \{v\}, \{\sigma \subseteq V \cup \{v\} \mid \sigma \cap V \in \mathcal{F}\}).$$

It is easy to see that $\|K * v\|$ is contractible via a straightline homotopy to the vertex δ_v . In fact, $\|K * v\|$ is homeomorphic to the topological cone $(\|K\| \times I) / (\|K\| \times \{1\})$ on $\|K\|$. If P is a poset with a maximum $\hat{1}$, then $\Delta(P) = \Delta(P \setminus \{\hat{1}\}) * \hat{1}$ and a similar statement holds for a minimum.

More generally, if $K = (V, \mathcal{F})$, $K' = (V', \mathcal{F}')$ with $V \cap V' = \emptyset$, then their *join* is the simplicial complex

$$K * K' = (V \cup V', \mathcal{F} \cup \mathcal{F}' \cup \{\sigma \cup \tau \mid \sigma \in \mathcal{F}, \tau \in \mathcal{F}'\}).$$

One has that $\|K * K'\| \cong \|K\| * \|K'\|$ where the *join* $A * B$ of topological spaces A, B is $(A \times B \times I) / R$ where R is the equivalence relation identifying all elements of $A \times B \times \{0\}$ which project to the same element of A and identifying all elements of $A \times B \times \{1\}$ that project to the same element of B . It is well known that the join is associative up to homeomorphism for locally compact spaces and that $S^n * S^m$ is homeomorphic to S^{n+m+1} . The join $S(X) = S^0 * X$ is called the *suspension* of X . Notice that $S(S^m)$ is homeomorphic to S^{m+1} . Similarly, we can define the suspension of a simplicial complex by taking the join with the simplicial complex consisting of two vertices and no simplices of dimension greater than zero.

If P, Q are disjoint posets, their *join* $P * Q$ is $P \cup Q$ where we define $p \leq q$ for all $p \in P$ and $q \in Q$ and where P and Q retain their respective orders. It is easy to see that $\Delta(P * Q) = \Delta(P) * \Delta(Q)$. The *suspension* $S(P)$ of a poset P is the join of the two element antichain with P . The operations of taking joins or suspensions for left regular bands correspond to performing these operations on their underlying posets.

Let $\mathbb{k}$ be a commutative ring with unit. Define the $\mathbb{k}$ -Leray number $L_{\mathbb{k}}(K)$ of a simplicial complex $K = (V, \mathcal{F})$ as follows:

$$L_{\mathbb{k}}(K) = \min\{d \mid \tilde{H}^i(K[W]; \mathbb{k}) = 0, \forall i \geq d, \forall W \subseteq V\}$$

where $\tilde{H}^i(L; \mathbb{k})$ denotes the reduced cohomology of L with coefficients in $\mathbb{k}$. The Leray number is classically defined when $\mathbb{k}$ is a field, and usually the definition is formulated in terms of homology. Of course, over a field homology and cohomology are dual vector spaces. It is known that $L_{\mathbb{k}}(K) = 0$ if and only if K is a simplex and $L_{\mathbb{k}}(K) \leq 1$ if and only if $K = \text{Clique}(\Gamma)$ for a chordal graph Γ (recall that Γ is *chordal* if it contains no induced cycle on four or more vertices). See [63, 64, 116, 120] for details, as well as for connections with Castelnuovo-Mumford regularity of Stanley-Reisner rings and with representability of simplicial complexes as nerves of families of compact convex subspaces of Euclidean space.

An important theorem in poset topology is Rota's cross-cut theorem. A proof can be found in [22].

Theorem 3.1 (Rota). *Let P be a finite poset such that any subset with a common lower bound has a meet. Let $\mathcal{M}(P)$ be the set of maximal elements of P and let K be the simplicial complex with vertex set $\mathcal{M}(P)$ and whose simplices are those subsets of $\mathcal{M}(P)$ with a common lower bound. Then $\|\Delta(P)\|$ is homotopy equivalent to $\|K\|$.*

There is, of course, a dual version of the cross-cut theorem concerning upper bounds, joins and minimal elements.

3.2. Regular CW complexes and CW posets. We use the books of Fritsch and Piccinini [53] and Lundell and Weingram [72] as our main references on CW complexes. A *regular CW complex* is a CW complex all of whose attaching maps are homeomorphisms to their images. Simplicial complexes, convex polytopes and polyhedral complexes are examples. In this text, we consider only finite CW complexes. Let us give the formal definition.

Let E^n be the closed unit ball in $\mathbb{R}^n$ and S^{n-1} be the unit sphere. Let X be a Hausdorff space and let Φ be a finite collection of continuous injective maps $\varphi: E^n \rightarrow X$ (ranging over different n). We call $\varphi(E^n)$ a *closed n -cell* of (X, Φ) and $\varphi(E^n \setminus S^{n-1})$ an *open n -cell*. We usually denote an open cell by e , the corresponding closed cell by $\bar{e}$ (which will be the topological closure of e) and we put $\partial e = \bar{e} - e$. The pair (X, Φ) is called a *regular CW complex* if:

- (1) the open cells partition X ;
- (2) for each open n -cell e with $n \geq 1$, we have that ∂e is contained in the union of the open k -cells with $k < n$.

A subcomplex of (X, Φ) is a CW complex (Y, Ψ) with Y a subspace of X and $\Psi \subseteq \Phi$. Equivalently, Y is a subspace with the property that it contains the closure of every open cell that it intersects. Note that each

closed cell is a subcomplex in a regular CW complex; see [72, Theorem III.2.1] or [53, Theorem 1.4.10]. A 0-cell of a CW complex is called a *vertex*.

If $q \geq 0$, then the q -skeleton of X is the subcomplex X^q consisting of all closed k -cells with $k \leq q$. A continuous mapping $f: X \rightarrow Y$ of CW complexes is called *cellular* if $f(X^q) \subseteq Y^q$ for all $q \geq 0$. A cellular mapping $f: X \rightarrow Y$ is called *regular* if it maps each open cell of X onto an open cell of Y [53, 72]. One can then show that the image under a regular cellular mapping f of each closed cell is a closed cell and that the image of each subcomplex is a subcomplex (cf. [53, Proposition 2.1.1]); in fact, if e is an open cell, then $f(\bar{e}) = \overline{f(e)}$, as is shown in the proof of [53, Proposition 2.1.1]. A bijective regular cellular map is called an *isomorphism*. The inverse of a bijective regular cellular map is automatically a regular cellular map [72, Proposition I.4.7].

The face poset $\mathcal{P}(X)$ of a regular CW complex X is the set of open cells of X ordered by $\sigma \leq \tau$ if $\sigma \subseteq \bar{\tau}$. We do not admit an empty face.

A (finite) poset P is said to be *graded* (or *ranked*) if $\Delta(P_{\leq p})$ is a pure simplicial complex for each $p \in P$. Define the *dimension* $\dim \sigma$ of $\sigma \in P$ by $\dim \sigma = \dim \Delta(P_{\leq \sigma})$. The *rank* of a closed interval $[\sigma, \tau]$ in a graded poset is given by

$$\text{rk}[\sigma, \tau] = \dim \tau - \dim \sigma.$$

It is the length of all maximal chains starting at σ and ending at τ . The face poset of a regular CW complex is graded (cf. [26, Corollary 4.7.12]).

A graded poset P is called a *CW poset* if $\|\Delta(P_{<\sigma})\|$ is homeomorphic to a $(\dim \sigma - 1)$ -sphere for each $\sigma \in P$ (where for this purpose we consider $S^{-1} = \emptyset$). In this case $\|\Delta(P_{\leq \sigma})\|$ is a cone on $\|\Delta(P_{<\sigma})\|$ and hence a topological $\dim \sigma$ -ball. Any (non-empty) lower set of a CW poset is a CW poset in its own right. An element σ of a CW poset with $\dim \sigma = q$ will be called a q -cell.

If X is a regular CW complex, then $\|\Delta(\mathcal{P}(X))\|$ is homeomorphic to X and $\Delta(\mathcal{P}(X))$ is the barycentric subdivision of X [26, 53, 72, 77]. In particular, if e is an open n -cell of X , then ∂e is a subcomplex and $\|\Delta(\mathcal{P}(X)_{<e})\| = \|\Delta(\mathcal{P}(\partial e))\| \cong S^{n-1}$. Thus if X is a regular CW complex, then $\mathcal{P}(X)$ is a CW poset and lower sets correspond to subcomplexes of X . If $f: X \rightarrow Y$ is a regular cellular map, then it induces an order-preserving map

$$\mathcal{P}(f): \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$$

via $e \mapsto f(e)$. An additional property of $\mathcal{P}(f)$ will be exploited later.

Conversely, there is a functor Σ from CW posets and appropriate morphisms to regular CW complexes and regular cellular maps which can be described on objects as follows. The geometric realization $\|\Delta(P)\|$ of the order complex of a CW poset P admits a regular CW complex structure $\Sigma(P)$ whose closed cells are the subcomplexes of $\|\Delta(P)\|$ of the form $\bar{e}_a = \|\Delta(P_{\leq a})\|$ with $a \in P$. Note that $\bar{e}_a$ is indeed a $\dim a$ -ball as described above. The open cell e_a corresponding to $a \in P$ is the union of the geometric simplices of $\|\Delta(P)\|$ spanned by chains with maximal element

a. Hence each point of $\|\Delta(P)\|$ belongs to a unique open cell. Indeed, a mapping $f: V \rightarrow \mathbb{R}$ from $\|\Delta(P)\|$ belongs to e_a if and only if a is the largest element of its support. Also, $\partial e_a = \|\Delta(P_{<a})\|$ and is contained in the union of the open cells e_b with $b < a$. Thus $\|\Delta(P)\|$ with the closed cells $\{\bar{e}_a\}_{a \in P}$ is a regular CW complex $\Sigma(P)$. Notice that $P \cong \mathcal{P}(\Sigma(P))$ via the map $a \mapsto e_a$. On the other hand, if X is a regular CW complex, the homeomorphism $\|\Delta(\mathcal{P}(X))\| \rightarrow X$ of [26, Proposition 4.7.9] is induced by an invertible regular cellular map and so $X \cong \Sigma(\mathcal{P}(X))$.

Let us say that an order-preserving map $f: P \rightarrow Q$ of posets is *cellular* if $\tau \leq f(\sigma)$ implies there exists $\sigma' \leq \sigma$ with $f(\sigma') = \tau$. Equivalently, f is cellular if $f(P_{\leq \sigma}) = Q_{\leq f(\sigma)}$ for all $\sigma \in P$. Alternatively, f is cellular if $f(L)$ is a lower set for each lower set L of P .

Proposition 3.2. *Let $f: X \rightarrow Y$ be a regular cellular map between regular cell complexes. Then the induced map $\mathcal{P}(f): \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$ is cellular.*

Proof. Clearly, $\mathcal{P}(f)$ is order-preserving. Let $e \in \mathcal{P}(X)$. Suppose that $e' \leq f(e)$, that is, $e' \subseteq \overline{f(e)}$. Let $y \in e'$. Then as $f(\bar{e}) = \overline{f(e)}$ (by the proof of [53, Proposition 2.1.1]), we can choose $x \in \bar{e}$ such that $f(x) = y$. There is a unique open cell e'' contained in $\bar{e}$ such that x belongs to e'' . Then $e'' \leq e$ and $f(e'')$ is an open cell containing y . Since each element of a CW complex belongs to a unique open cell, we conclude that $f(e'') = e'$. This concludes the proof that $\mathcal{P}(f)$ is cellular. $\square$

It follows that $\mathcal{P}$ is a functor from the category of regular CW complexes and regular cellular maps to the category of CW posets and cellular maps. We now want to show that Σ extends to a functor going the opposite way. The functoriality of Σ seems to be novel to this text.

Lemma 3.3. *Let $f: P \rightarrow Q$ be a cellular map of posets. Then given any chain $\tau_0 < \dots < \tau_q = \tau$ in Q and any $\sigma \in P$ with $f(\sigma) = \tau$, there is a chain $\sigma_0 < \dots < \sigma_q = \sigma$ in P with $f(\sigma_i) = \tau_i$ for $0 \leq i \leq q$.*

Proof. Set $\sigma_q = \sigma$. Let $0 \leq i < q$ and assume that σ_j has been defined for $i < j \leq q$ with $\sigma_{i+1} < \dots < \sigma_q = \sigma$ and $f(\sigma_j) = \tau_j$. By definition of a cellular map there exists $\sigma_i < \sigma_{i+1}$ such that $f(\sigma_i) = \tau_i$. This completes the proof. $\square$

Lemma 3.4. *If $f: P \rightarrow Q$ is a cellular map of CW posets, then the induced simplicial map $\Delta(f): \Delta(P) \rightarrow \Delta(Q)$ induces a regular cellular map $\Sigma(f): \Sigma(P) \rightarrow \Sigma(Q)$.*

Proof. Lemma 3.3 easily implies that if $p \in P$, then $\Delta(f)$ takes the closed cell $\Delta(P_{\leq p})$ of $\Sigma(P)$ onto the closed cell $\Delta(Q_{\leq f(p)})$ of $\Sigma(Q)$ and $\dim \Delta(Q_{\leq f(p)}) \leq \dim \Delta(P_{\leq p})$. Thus $\Sigma(f)$ is cellular. The open cell of $\Sigma(P)$ corresponding to $p \in P$ is the union of those simplices in $\|\Delta(P)\|$ spanned by chains with maximum element p . Hence $\Delta(f)$ takes the open cell corresponding to p into the open cell corresponding to $f(p)$, and in fact onto it by Lemma 3.3. Thus $\Delta(f)$ is a regular cellular map. $\square$

We conclude that Σ is a functor from the category of CW posets and cellular maps to the category of regular CW complexes and regular cellular maps with the property that $\Sigma(\mathcal{P}(X)) \cong X$ for any regular CW complex X .

Remark 3.5. It is easy to see that $\mathcal{P} \circ \Sigma$ is isomorphic to the identity functor. But it is possible that $\mathcal{P}(f) = \mathcal{P}(g)$ for two distinct regular cellular maps $f, g: X \rightarrow Y$. Nonetheless, it is not difficult to show that $\Sigma(\mathcal{P}(f))$ is always homotopy equivalent to f for any regular cellular map (by an aspherical carrier type argument). In particular, as far as homology is concerned there is not much of a difference between working with CW posets and cellular maps and with regular CW complexes and regular cellular maps.

Notice that if K, L are simplicial complexes and $f: K \rightarrow L$ is a simplicial map, then the induced map $\mathcal{P}(K) \rightarrow \mathcal{P}(L)$ is cellular. Moreover, $\Sigma(f): K = \Sigma(\mathcal{P}(K)) \rightarrow \Sigma(\mathcal{P}(L)) = L$ is the map of geometric realizations induced by the simplicial map f . So simplicial complexes with simplicial maps can be viewed either as a subcategory of CW posets and cellular maps, or a subcategory of regular CW complexes with regular cellular maps.

We now state some further properties of cellular maps that will be used in our study of left regular bands. The first shows that the support homomorphism is cellular.

Proposition 3.6. *If B is a left regular band and $\sigma: B \rightarrow \Lambda(B)$ is the support map, then σ is cellular. Moreover, $a < b$ implies $\sigma(a) < \sigma(b)$.*

Proof. The second statement follows because $a \leq b$ and $\sigma(a) = \sigma(b)$ implies $ba = a$ and $ba = b$, whence $a = b$. To see that σ is cellular, let $X \leq \sigma(b)$. Choose a with $Ba = X$. Then $\sigma(ba) = \sigma(b) \wedge X = X$ and $ba \leq b$. $\square$

Corollary 3.7. *Let $f: P \rightarrow Q$ be a surjective cellular map of posets and suppose, furthermore, that $\sigma < \sigma'$ implies that $f(\sigma) < f(\sigma')$. Then P is graded if and only if Q is graded and, moreover, if this is the case then f preserves ranks of intervals.*

Proof. Let $p \in P$ and $q = f(p)$. Because f preserves strict inequalities, it is immediate that the simplicial map $f: \Delta(P_{\leq p}) \rightarrow \Delta(Q_{\leq q})$ preserves dimensions of simplices. Also Lemma 3.3 implies that each simplex of $\Delta(Q_{\leq q})$ is the image of a simplex of $\Delta(P_{\leq p})$. It follows that the facets of $\Delta(Q_{\leq q})$ are precisely the images under f of the facets of $\Delta(P_{\leq p})$. We conclude $\Delta(P_{\leq p})$ is pure if and only if $\Delta(Q_{\leq q})$ is pure. The result now follows. $\square$

Recall that a topological space is *acyclic* if its reduced homology vanishes over $\mathbb{Z}$. Contractible spaces are of course acyclic. If X is acyclic, then the reduced homology of X vanishes over any coefficient ring by the universal coefficient theorem. If P is a poset, we will say that P has a topological property (e.g., contractibility, acyclicity, connectivity) if $\|\Delta(P)\|$ has that property.

Proposition 3.8. *Let $f: P \rightarrow P$ be an idempotent cellular mapping on a poset. Then $f(P)$ is a lower set and hence if $f(p) = p$, then $f(q) = q$ for all $q \leq p$. If P is a (connected/contractible/acyclic) CW poset then so is $f(P)$.*

Proof. The image $f(P)$ of the lower set P under a cellular map is a lower set. Hence, if $f(p) = p$ and $q \leq p$, then $q \in f(P)$ and thus $f(q) = q$ because f is idempotent. If P is a CW poset, then the subcomplex $\Sigma(f(P)) = \Sigma(f)(\Sigma(P))$ is a retract of $\Sigma(P)$ and the result follows. $\square$

The following trivial observation will be useful later.

Proposition 3.9. *Let P be a CW poset with a maximum. Then $\Sigma(P)$ is a regular CW decomposition of a $\dim \Delta(P)$ -ball. Conversely, if Q is the face poset of a regular CW decomposition of a $\dim \Delta(Q)$ -sphere, then $P = Q \cup \{\hat{1}\}$ (where $\hat{1}$ is an adjoined maximum) is the face poset of a regular CW decomposition of a $\dim \Delta(P)$ -ball.*

Proof. Suppose first that P is a CW poset with maximum $\hat{1}$. Then by definition $\|\Delta(P_{<\hat{1}})\|$ is homeomorphic to a $(\dim \Delta(P) - 1)$ -sphere and so $\|\Delta(P)\| = \|\Delta(P_{<\hat{1}}) * \hat{1}\|$ is homeomorphic to a $\dim \Delta(P)$ -ball, being a cone on a sphere. Suppose now that Q is the face poset of a regular CW decomposition of a $\dim \Delta(Q)$ -sphere and $P = Q \cup \{\hat{1}\}$. Then for $q \in Q$, we have $P_{<q} = Q_{<q}$ and $P_{<\hat{1}} = Q$ and so P is a CW poset. By the above P is the face poset of a $\dim \Delta(P)$ -ball. $\square$

The following propositions will also be helpful.

Proposition 3.10. *Suppose that P, Q are CW posets with*

$$\|\Delta(P)\| \cong S^{\dim(P)}.$$

Then $P * Q$ is a CW poset. In particular, the suspension $S(Q)$ of Q is a CW poset.

Proof. Clearly if $p \in P$, then $(P * Q)_{<p} = P_{<p}$ and hence has order complex a sphere of the appropriate dimension. If $q \in Q$, then $(P * Q)_{\leq q} = P * Q_{\leq q}$ has dimension $\dim \Delta(P) + \dim \Delta(Q_{\leq q}) + 1$. On the other hand, $(P * Q)_{<q} = P * Q_{<q}$ and so $\|(P * Q)_{<q}\|$ is a sphere of the desired dimension from the formula $S^m * S^n = S^{m+n+1}$. The final statement is the special case when $\|\Delta(P)\| \cong S^0$. $\square$

Recall that if P, Q are posets, then their direct product is a poset by putting $(p, q) \leq (p', q')$ if $p \leq p'$ and $q \leq q'$.

Proposition 3.11. *If P, P' are CW posets, then so is $P \times P'$.*

Proof. Suppose that $(p, p') \in P \times P'$, $\dim \Delta(P_{\leq p}) = r$ and $\dim \Delta(P'_{\leq p'}) = s$. Clearly $\dim \Delta((P \times P')_{\leq (p, p')}) = r + s$. A general result in poset topology [22, Equation (9.7)] says that

$$\|\Delta((P \times P')_{<(p, p')})\| \cong \|\Delta(P_{<p}) * \Delta(P'_{<p'})\| \cong S^{r-1} * S^{s-1} \cong S^{r+s-1}.$$

Thus $P \times P'$ is a CW poset. $\square$

3.3. Oriented interval greedoids. In 2008, Billera, Hsiao, and Provan introduced a construction that associates with every antimatroid a polytopal subdivision of a sphere [20]. This result is structurally similar to the sphericity theorem for oriented matroids, which associates a regular cell decomposition of the sphere with every oriented matroid (as already discussed in Section 2.3). The second author and Thomas developed a theory of “oriented interval greedoids,” akin to the theory of “oriented matroids,” in order to unify these results [100]. Their starting point was the observation that both complexes can be endowed with the structure of a left regular band monoid. We review the results here.

3.3.1. Motivation from oriented matroids. In order to motivate the definition of an oriented interval greedoid, we begin by recasting the definition of a covector for an oriented matroid.

Let $\mathcal{M} = (E, \mathcal{F})$ be a matroid, where we retain the notation of Section 2.1. A note on terminology is in order. In the matroid literature, the notions of “closed set” and “flat” coincide, but these admit different generalizations for greedoids. Each notion gives rise to a poset (with the former a lattice), which turn out to be isomorphic to each other in the case of interval greedoids [25, Theorem 3.6]. In particular, the greedoid notion of flat does not conform with the matroid notion of flat (cf. Section 2.1). To distinguish between these notions, we will use the term “greedoid flat” for flats in the sense of greedoid theory.

Introduce an equivalence relation on $\mathcal{F}$ by declaring $X \sim Y$ if and only if $\overline{X} = \overline{Y}$. The equivalence classes for this relation, denoted by $[X]$, are called the *greedoid flats* of the matroid. The set Φ of all greedoid flats of the matroid admits a partial order induced by inclusion: $[Y] \leq [X]$ if and only if there exists $Z \in E \setminus Y$ with $Y \cup Z \in \mathcal{F}$ and $Y \cup Z \sim X$. In particular, we have $[Y] \leq [X]$ whenever $Y \subseteq X$.

For $X \in \mathcal{F}$, let $\xi(X)$ be the union of all subsets in $[X]$:

$$\xi(X) = \bigcup_{Y \sim X} Y.$$

For matroids without loops (recall that a loop is an element that does not belong to any set in $\mathcal{F}$), the sets $\xi(X)$ are the *flats* of the matroid (but will be called closed sets in the greedoid context). In this case, the elements of $\xi(X)$ are those elements that can be added to X without increasing its rank. Define $\Gamma(X)$ to be the set of $e \in E$ such that $X \cup \{e\} \in \mathcal{F}$.

As briefly pointed out in Section 2.3, there is a matroid underlying every oriented matroid. For convenience, assume that the oriented matroid contains no loops. It is readily obtained by passing from the set of covectors of the oriented matroid to their zero sets:

$$x \mapsto Z(x) = \{e \in E \mid x_e = 0\}.$$

These are precisely the flats of a unique matroid (called the *underlying matroid* of the oriented matroid). With this terminology and notation, it

is straightforward to see that for every covector x of an oriented matroid, there exists a greedoid flat $[X]$ of the underlying matroid satisfying:

- $x_e = 0$ for each $e \in \xi(X)$ (i.e., $x_e = 0$ if and only if $e \in \overline{X}$);
- $x_e \in \{+, -\}$ for all $e \in \Gamma(X)$ (i.e., $x_e = \pm$ if and only if $X \cup \{e\}$ is independent).

The notion of a covector for an interval greedoid is based on these properties.

3.3.2. Interval greedoids. Let E denote a finite set and $\mathcal{F}$ a set of subsets of E . An *interval greedoid* is a pair $(E, \mathcal{F})$ satisfying the following properties for all $X, Y, Z \in \mathcal{F}$:

- (IG1) If $X \neq \emptyset$, then there exists an $x \in X$ such that $X \setminus \{x\} \in \mathcal{F}$.
- (IG2) If $|X| > |Y|$, then there exists an $x \in X \setminus Y$ such that $Y \cup x \in \mathcal{F}$.
- (IG3) If $X \subseteq Y \subseteq Z$ and $e \in E \setminus Z$ with $X \cup e \in \mathcal{F}$ and $Z \cup e \in \mathcal{F}$, then $Y \cup e \in \mathcal{F}$.

A *loop* in an interval greedoid $(E, \mathcal{F})$ is an element $e \in E$ that belongs to no subset in $\mathcal{F}$. We will restrict our attention to interval greedoids that do not contain loops.

Every matroid is an interval greedoid. Further examples of interval greedoids will be given below (in particular, antimatroids/convex geometries).

Let $(E, \mathcal{F})$ be an interval greedoid. For $X \in \mathcal{F}$, let $\mathcal{F}/X$ denote the collection of subsets that can be added to X with the result still in $\mathcal{F}$:

$$\mathcal{F}/X = \{Y \subseteq E \setminus X \mid Y \cup X \in \mathcal{F}\}.$$

Similarly, the set of *continuations* $\Gamma(X)$ of $X \in \mathcal{F}$ is the set of elements in E that can be added to X with the result still in $\mathcal{F}$:

$$\Gamma(X) = \{x \in E \setminus X \mid X \cup \{x\} \in \mathcal{F}\}.$$

Define an equivalence relation on $\mathcal{F}$ by declaring $X \sim Y$ if and only if $\mathcal{F}/X = \mathcal{F}/Y$. For interval greedoids, $\mathcal{F}/X = \mathcal{F}/Y$ if and only if $\Gamma(X) = \Gamma(Y)$. The equivalence classes of this relation are called *greedoid flats* and we denote the equivalence class of X by $[X]$. The set of greedoid flats Φ is a poset with partial order induced by inclusion, analogously to the matroid case. In fact, Φ is a *semimodular lattice* [23, Theorem 5.3], that is, a lattice satisfying axioms (G2) and (G3) of Definition 2.8. In particular, the minimal element of Φ is $[\emptyset]$, and $[Y] \leq [X]$ whenever $Y \subseteq X$. For a greedoid flat $[X]$, define

$$\xi(X) = \bigcup_{\substack{Y \in \mathcal{F} \\ \Gamma(Y) = \Gamma(X)}} Y.$$

The sets of the form $\xi(X)$ are the *closed sets* of the interval greedoid.

3.3.3. Oriented interval greedoids. A greedoid flat $[X]$ partitions the ground set E into three subsets (possibly empty):

- $\Gamma(X)$, the set of continuations of X ;
- $\xi(X)$, the set of elements that lie on a greedoid flat; and

- the set of elements that belong to neither $\Gamma(X)$ nor $\xi(X)$.

A *covector with underlying greedoid flat* $[X]$ is any vector $x \in \{0, +, -, 1\}^E$ satisfying:

- $x_e \in \{+, -\}$ if $e \in \Gamma(X)$;
- $x_e = 0$ if $x \in \xi(X)$;
- $x_e = 1$ if $x \notin \Gamma(X) \cup \xi(X)$.

Let $x \mapsto \text{flat}(x)$ denote the map that sends a covector to its underlying greedoid flat. For two covectors x and y of $(E, \mathcal{F})$, define their *separation set* by

$$S(x, y) = \{e \in E \mid x_e = -y_e \in \{+, -\}\}.$$

To define the *composition of covectors*, we first need an auxiliary operation. For $x, y \in \{0, +, -, 1\}^E$, define

$$(x \star y)_e = \begin{cases} y_e, & \text{if } y_e < x_e, \\ x_e, & \text{otherwise,} \end{cases}$$

where the symbols $0, +, -, 1$ are partially ordered so that 1 and 0 are the (unique) minimal and maximal elements, respectively, and $+$ and $-$ are incomparable. See Figure 6 for the Hasse diagram of $\{0, +, -, 1\}$. Note that we use the opposite ordering convention of [100]. Then the composition




FIGURE 6. The Hasse diagram of $\{0, +, -, 1\}$

$x \circ y$ of two covectors x and y with underlying greedoid flats $[X]$ and $[Y]$, respectively, is defined as

$$(x \circ y)_e = \begin{cases} (x \star y)_e, & \text{if } e \in \Gamma(W) \cup \xi(W), \\ 1, & \text{otherwise.} \end{cases}$$

where W is maximal among the sets in $\mathcal{F}$ that are contained in $\xi(X) \cap \xi(Y)$. (Note that, unlike for oriented matroids, this operation is not necessarily computed componentwise. See Figure 11 and the surrounding discussion.)

Since we are restricting our attention to interval greedoids without loops, the all zero covector is the identity element for this operation. (If the interval greedoid admits loops, then the identity element is the covector w with $w_e = 1$ if e is a loop and $w_e = 0$, otherwise.)

An *oriented interval greedoid* is a triple $(E, \mathcal{F}, \mathcal{L})$, where $(E, \mathcal{F})$ is an interval greedoid and $\mathcal{L}$ is a set of covectors of $(E, \mathcal{F})$ satisfying the following axioms.

- (OG1) The map $\text{flat}: \mathcal{L} \rightarrow \Phi$ is surjective, where Φ is the lattice of greedoid flats of $(E, \mathcal{F})$.
- (OG2) If $x \in \mathcal{L}$, then $-x \in \mathcal{L}$ (where $-1 = 1$).
- (OG3) If $x, y \in \mathcal{L}$, then $x \circ y \in \mathcal{L}$.
- (OG4) If $x, y \in \mathcal{L}$, $e \in S(x, y)$ and $(x \circ y)_e \neq 1$, then there exists $z \in \mathcal{L}$ such that $z_e = 0$ and for all $f \notin S(x, y)$, if $(x \circ y)_f \neq 1$, then $z_f = (x \circ y)_f = (y \circ x)_f$.

These axioms are modeled on the covector axioms for oriented matroids; compare with Definition 2.9. Moreover, if $(E, \mathcal{F})$ is a matroid without loops, then this notion coincides with that of an oriented matroid. More precisely, $\mathcal{L}$ is the collection of covectors of an oriented matroid with underlying matroid $(E, \mathcal{F})$ if and only if $(E, \mathcal{F}, \mathcal{L})$ is an oriented interval greedoid [100, Theorem 3.12]. Similarly, if the underlying interval greedoid is an antimatroid (see below), then this recovers the Billera–Hsiao–Provan complex [100, Theorem 3.14]. Other examples of oriented interval greedoids include complexified hyperplane arrangements [100, Theorem 3.20].

The covectors of an oriented interval greedoid form a left regular band monoid.

Proposition 3.12. *Let $\mathcal{L}$ be the set of covectors for an oriented interval greedoid on an interval greedoid without loops $(E, \mathcal{F})$.*

- (1) *$\mathcal{L}$ equipped with the composition of covectors forms a left regular band monoid with identity element equal to the all zero covector.*
- (2) *The support lattice $\Lambda(\mathcal{L})$ is isomorphic to the lattice of greedoid flats Φ of $(E, \mathcal{F})$ and the support map corresponds to $\text{flat}: \mathcal{L} \rightarrow \Phi$.*

Since $\mathcal{L}$ is a left regular band monoid, it is a poset with respect to the $\mathcal{R}$ -order, as we have been considering previously. The order is characterized by $x \geq y$ if and only if $x \circ y = y$, or equivalently $x \geq y$ if and only if $x_e \geq y_e$ for all $e \in E$ [100, Lemma 3.7]. The poset $\mathcal{L}^{\text{op}} \setminus \{0\}$ obtained from $\mathcal{L}$ by considering the *opposite* of the $\mathcal{R}$ -order and removing the identity is the face poset of a regular CW complex. The following result is [100, Theorem 6.1], where the reader is referred to [26, Chapter 4] for the notions of shellable and piecewise linear regular CW decompositions of a sphere.

Theorem 3.13. *Let $\mathcal{L}$ denote the set of covectors of an oriented interval greedoid. Then $\mathcal{L}^{\text{op}} \setminus \{0\}$ is isomorphic to the face poset of a shellable (hence piecewise linear) regular CW decomposition of a sphere.*

Here is an example extracted from [100, Example 6.2], which we illustrate in Figures 7, 8 and 9. Consider the interval greedoid on the ground set $E = \{a, b, c, e\}$ defined by

$$\mathcal{F} = \{\emptyset\} \cup F \cup \{X \cup \{e\} \mid X \in F\}, \quad (3.1)$$

where $F = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}\}$. The lattice of greedoid flats Φ of this interval greedoid is depicted in Figure 7. Next, consider a two-




FIGURE 7. The lattice of greedoid flats Φ of the interval greedoid defined on the ground set $E = \{a, b, c, e\}$ by (3.1).

dimensional sphere: draw three great circles on it, each passing through the north and south poles, and mark the points where each great circle crosses the equator (see Figure 8). The result is a regular CW decomposition of a sphere with 6 two-cells, 12 one-cells, and 8 zero-cells, whose cells correspond to a collection of covectors for an oriented interval greedoid structure for $(E, \mathcal{F})$ (see Figure 9).

3.3.4. Contraction and deletion of oriented interval greedoids. Let $(E, \mathcal{F})$ be an interval greedoid with lattice of greedoid flats Φ . The *contraction* of $(E, \mathcal{F})$ along $X \in \mathcal{F}$ is the interval greedoid $(\bigcup_{Y \in \mathcal{F}/X} Y, \mathcal{F}/X)$, where

$$\mathcal{F}/X = \{Y \subseteq E \setminus X \mid Y \cup X \in \mathcal{F}\}.$$

Let Φ/X , Γ/X and ξ/X denote the corresponding notions in the contraction.

Fix $X \in \mathcal{F}$. Let y be a covector of $(E, \mathcal{F})$ such that $\text{flat}(y) \geq [X]$ in Φ . Then there exists $Y \in \mathcal{F}/X$ such that $\text{flat}(y) = [X \cup Y]$. Define the *contraction* of y along X as

$$\text{con}_X(y)_e = \begin{cases} y_e, & \text{if } e \in (\Gamma/X)(Y), \\ 0, & \text{if } e \in (\xi/X)(Y), \\ 1, & \text{otherwise.} \end{cases} \quad (3.2)$$

If $\mathcal{L}$ is the set of covectors of an oriented interval greedoid on $(E, \mathcal{F})$, then

$$\mathcal{L}/X = \{\text{con}_X(y) \mid y \in \mathcal{L} \text{ and } \text{flat}(y) \geq [X]\}$$




FIGURE 8. A regular CW decomposition of a sphere corresponding to an oriented interval greedoid on the interval greedoid defined in (3.1). The cells are labelled by the covectors of the oriented interval greedoid. See Figure 9 for its (opposite) face poset.




FIGURE 9. The underlying poset of the left regular band of covectors from Figure 8. It is isomorphic to the opposite poset of the face poset (including an empty face) of the regular CW decomposition of the sphere in Figure 8.

defines an oriented interval greedoid over the contraction of $(E, \mathcal{F})$ by X [100, Proposition 4.3]. By [100, Proposition 4.4], $\mathcal{L}/X$ is isomorphic to the contraction of $\mathcal{L}$ to $[X]$ (in the sense of left regular bands) via the monoid homomorphism

$$\text{con}_X: \mathcal{L}_{\geq[X]} \longrightarrow \mathcal{L}/X.$$

We now describe the deletion operation on oriented interval greedoids. Let W be an arbitrary subset of E . The *deletion* of $(E, \mathcal{F})$ to W is the

interval greedoid $(W, \mathcal{F}|_W)$ defined by

$$\mathcal{F}|_W = \{X \in \mathcal{F} \mid X \subseteq W\}.$$

Let $\Phi|_W$, $\Gamma|_W$ and $\xi|_W$ denote the corresponding notions for the deletions.

Fix $W \subseteq E$. Let y be a covector of $(E, \mathcal{F})$ with $\text{flat}(y) = [Y]$. Define the *deletion* of y to W as

$$\text{res}_W(y)_w = \begin{cases} y_w, & \text{if } w \in \Gamma|_W(Y), \\ 0, & \text{if } w \in \xi|_W(Y), \\ 1, & \text{otherwise.} \end{cases}$$

If $\mathcal{L}$ is the set of covectors of an oriented interval greedoid on $(E, \mathcal{F})$ and $W \subseteq E$, then define the *deletion* of $\mathcal{L}$ to W to be

$$\mathcal{L}|_W = \{\text{res}_W(y) \mid y \in \mathcal{L}\}.$$

It is not true in general that deletions of oriented interval greedoids to arbitrary subsets $W \subseteq E$ are again oriented interval greedoids. However, it is true for deletion to $\Gamma(\emptyset)$, in which case the deletion is an oriented matroid [100, Proposition 4.12]; and for deletion to $\xi(X)$ for any $X \in \mathcal{F}$ [100, Theorem 4.15], in which case

$$\text{res}_{\xi(X)}: x\mathcal{L} \longrightarrow \mathcal{L}|_{\xi(X)}$$

is a monoid isomorphism for any covector x with $\text{flat}(x) = [X]$, with inverse $\text{res}_{\xi(X)}(y) \mapsto x \circ y$ for all $y \in \mathcal{L}$ [100, Proposition 4.16].

3.3.5. Oriented interval greedoids from convex geometries. Convex geometries give rise to a special class of interval greedoids called antimatroids. In this setting the covectors and the composition of covectors admit a very nice geometric description. We begin by defining convex geometries.

Just as matroids can be viewed as an abstraction of linear independence of vectors in $\mathbb{R}^d$, convex geometries can be viewed as an abstraction of convexity of vectors in $\mathbb{R}^d$. In the following, think of E as a finite subset of $\mathbb{R}^d$; of τ as the convex hull operator $\tau(A) = \text{conv}(A) \cap E$ for $A \subseteq E$; and of $\text{ext}(A)$ as the set of extreme points of the convex hull of A .

A *convex geometry* is a pair (E, τ) with E a finite set and $\tau: 2^E \longrightarrow 2^E$ a non-decreasing, order preserving and idempotent function, satisfying the following *anti-exchange* property:

(AE) If $x, y \notin \tau(X)$, $x \neq y$, and $y \in \tau(X \cup x)$, then $x \notin \tau(X \cup y)$.

The subsets $A \subseteq E$ satisfying $\tau(A) = A$ are called *closed sets*. The *extreme points* $\text{ext}(A)$ of $A \subseteq E$ are the points $x \in A$ satisfying $x \notin \tau(A \setminus x)$.

If (E, τ) is a convex geometry, then $(E, \mathcal{F})$ is an interval greedoid, where $\mathcal{F}$ consists of the complements of the closed sets. These interval greedoids are called *antimatroids* or *upper interval greedoids*. If $X \in \mathcal{F}$, then $E \setminus X$ is a closed set of the convex geometry and we define $\Gamma(X) = \text{ext}(E \setminus X)$.

A covector for an antimatroid admits a nice geometric description in terms of the corresponding convex geometry. Let $X \in \mathcal{F}$. Then X corresponds to

the closed set $C = E \setminus X$ in the convex geometry. A covector with underlying greedoid flat $[X]$ assigns $+$ or $-$ to the extreme points of C , assigns 0 to the points of the exterior of C , and assigns 1 to the non-extreme points contained in C . See Figure 10.




FIGURE 10. Covectors of an antimatroid: extreme points of a closed set are labelled by $+$ or $-$; non-extreme interior points are labelled 1; and exterior points are labelled 0.

The composition $x \circ y$ of two covectors x and y is computed as follows. (Figure 11 depicts the product of the two covectors in Figure 10.) Consider the convex hull of the union of the corresponding closed sets. The exterior points are labelled 0. The non-extreme interior points are labelled 1. If e is an extreme point, then it is an extreme point for x or for y (possibly both). If it is an extreme point for x , then e takes the sign x_e ; otherwise, it takes the sign y_e .

This example exemplifies that the product on covectors is not always computed componentwise. It is possible that $x_e = y_e = 0$ yet $(x \circ y)_e \neq 0$. This happens if e is exterior to both closed sets, yet it is interior to the convex hull of the union of the closed sets. Similarly, extreme points of the closed sets can become interior to the convex hull of their union.




FIGURE 11. The products $x \circ y$ (left) and $y \circ x$ (right) of the covectors x and y depicted in Figure 10.

The set of all covectors of an antimatroid forms an oriented interval greedoid [100, Theorem 3.14]. In fact, this is the only oriented interval greedoid structure on an antimatroid [100, Proposition 3.15]. This collection of

covectors is the central object of study in the work of Billera, Hsiao and Provan [20]. To state their results, we recall some definitions from [20].

Define a poset Q_L as follows. Let L denote the lattice of closed sets of the convex geometry ordered by inclusion. The elements of Q_L are equivalence classes of pairs (A, ε) with $A \in L$ and $\varepsilon: E \rightarrow \{+, -\}$, where $(A, \alpha) \sim (B, \beta)$ if $A = B$ and $\alpha|_{\text{ext}(A)} = \beta|_{\text{ext}(B)}$. The partial order on Q_L is defined as $(A, \alpha) \leq (B, \beta)$ if and only if $A \subseteq B$ and α and β agree on $\text{ext}(A) \cap \text{ext}(B)$. Note that Q_L admits a minimal element corresponding to the pair $(\emptyset, \varepsilon)$, where ε is the empty map. Q_L is also has a maximal element $\hat{1}$ adjoined.

Elements of Q_L are in bijection with covectors of the oriented interval greedoid on the antimatroid. To see this, note that elements of Q_L determine covectors, and vice-versa. Indeed, the equivalence class of (A, α) is determined by A and $\alpha|_{\text{ext}(A)}$, so one obtains a covector $\hat{\alpha}$ by declaring $\hat{\alpha}_e = \alpha(e)$ if $e \in \text{ext}(A)$, $\hat{\alpha}_e = 1$ if $e \in A \setminus \text{ext}(A)$ and $\hat{\alpha}_e = 0$ if $e \notin A$. Conversely, starting with a covector x , consider the closed set $A = \{e \in E \mid x_e \neq 0\}$ and any map $\alpha: E \rightarrow \{+, -\}$ satisfying $\alpha(e) = x_e$ on $\text{ext}(A)$.

The partial order on Q_L is opposite to the partial order on $\mathcal{L}$. Indeed, if $A \subseteq B$ and α and β agree on $\text{ext}(A) \cap \text{ext}(B)$, then the corresponding covectors $\hat{\alpha}$ and $\hat{\beta}$ satisfy $\hat{\alpha}_e \geq \hat{\beta}_e$ for all $e \in E$:

- if $e \notin A$, then $\hat{\alpha}_e = 0 \geq \hat{\beta}_e$;
- if $e \in \text{ext}(A) \cap \text{ext}(B)$, then $\hat{\alpha}_e = \hat{\beta}_e$ by assumption;
- if $e \in \text{ext}(A) \setminus \text{ext}(B)$, then $\hat{\alpha}_e \in \{+, -\}$ and $\hat{\beta}_e = 1$, so $\hat{\alpha}_e \geq \hat{\beta}_e$;
- if $e \in A \setminus \text{ext}(A)$, then $e \notin B \setminus \text{ext}(B)$ and $\hat{\alpha}_e = 1 = \hat{\beta}_e$.

Billera, Hsiao and Provan prove that $Q_L \setminus \{\hat{0}, \hat{1}\}$ is the face poset of a regular CW decomposition of a sphere. It follows that $\mathcal{L} \setminus \{0\}$ is the opposite poset of a face poset of a regular CW decomposition of a sphere.

3.4. The topology of left regular bands. In this section, we describe the topology of the families of left regular bands discussed in the previous section (where by the topology of a left regular band, we mean its topology as a poset). Note that if B is a left regular band monoid and also a CW poset, then it is, in fact, the face poset of a regular CW decomposition of a ball by Proposition 3.9 (since the identity is a maximum). This need not be the case for left regular bands without an identity element.

Proposition 3.14. *The following hold.*

- (1) *If $\mathcal{L}$ is an oriented matroid, then $\mathcal{L}_{\geq X}$ is the face poset of a regular cell decomposition of a ball for each $X \in \Lambda(\mathcal{L})$.*
- (2) *If $(E, \mathcal{L}, g)$ is an affine oriented matroid and $X \in \Lambda(\mathcal{L}^+(g))$, then $\mathcal{L}^+(g)_{\geq X}$ is the face poset of a regular CW decomposition of a ball.*
- (3) *Let $(E, \mathcal{L})$ be a COM and $X \in \Lambda(\mathcal{L})$. Then $\mathcal{L}_{\geq X}$ is the face poset of a contractible regular CW complex.*
- (4) *If $\mathcal{A}$ is a complex hyperplane arrangement, then $\mathcal{F}(\mathcal{A})_{\geq X}$ is the face poset of a regular CW decomposition of a ball for each $X \in \Lambda(\mathcal{F}(\mathcal{A}))$.*

- (5) If $(E, \mathcal{F}, \mathcal{L})$ is an oriented interval greedoid and $X \in \Lambda(\mathcal{L})$, then $\mathcal{L}_{\geq X}$ is the face poset of a regular CW decomposition of a ball.
- (6) If $\Gamma = (V, E)$ is a graph, then $\Lambda(B(\Gamma)) \cong (P(V), \cup)$. If $Y \subseteq X \subseteq V$ and $e_Y \in B(\Gamma)$ has support Y , then $\|\Delta(\partial e_Y B(\Gamma)_{\geq X})\|$ is homotopy equivalent to $\|\text{Cliq}(\Gamma[X \setminus Y])\|$.

Proof. For (1), we observe that $\mathcal{L}_{\geq X}$ is the set of covectors of a contraction of $\mathcal{L}$ and so it suffices to prove that $\mathcal{L}$ is a CW poset for any oriented matroid. From [26, Corollary 4.3.4], we have that $\mathcal{L} \setminus \{0\}$ is the face poset of a regular CW decomposition of a sphere. Thus $\mathcal{L}$ is the face poset of a regular CW decomposition of a ball by Proposition 3.9.

Similarly to (1), each contraction of an affine oriented matroid is again an affine oriented matroid and so to prove (2), we just need that $\mathcal{L}^+(g)$ is the face poset of a regular CW decomposition of a ball. But this is the content of [26, Corollary 4.5.8].

To prove (3), again we can use contractions to reduce to proving that $\mathcal{L}$ is a contractible CW poset by Proposition 2.15. But this is proved in [13, Section 11]. To see that $\mathcal{L}$ is the face poset of a regular cell complex, one just uses that each $x\mathcal{L}$ with $x \in \mathcal{L}$ is isomorphic to the monoid of covectors of an oriented matroid by Proposition 2.15.

Item (4) is the most complicated case because contractions of complex hyperplane face monoids need not again be complex hyperplane face monoids. It is shown in the proof of [75, Proposition 4.21] that if $X \in \Lambda(\mathcal{F}(\mathcal{A}))$, then $\mathcal{F}(\mathcal{A})_{\geq X}^{\text{op}} \setminus \{0\}$ is the face poset of a PL (piecewise linear) regular CW decomposition of a sphere (note that in [75] we followed the convention that regular CW complexes have empty faces). Thus $\mathcal{F}(\mathcal{A})_{\geq X} \setminus \{0\}$ is the face poset of a regular CW decomposition of a sphere by [26, Proposition 4.7.26] and so $\mathcal{F}(\mathcal{A})_{\geq X}$ is the face poset of a regular CW decomposition of a ball by Proposition 3.9.

To prove (5), we use closure of oriented interval greedoids under contraction to reduce to proving that $\mathcal{L}$ is the face poset of a regular CW decomposition of a ball. By Theorem 3.13, $\mathcal{L}^{\text{op}} \setminus \{0\}$ is the face poset of a PL regular CW decomposition of a sphere and hence $\mathcal{L} \setminus \{0\}$ is face poset of a regular CW decomposition of a sphere by [26, Proposition 4.7.26]. Therefore, $\mathcal{L}$ is the face poset of a regular CW decomposition of a ball by Proposition 3.9.

The final item is proved in the course of the proof of [75, Theorem 4.16]. As $e_Y B(\Gamma_{\geq X}) \cong B(\Gamma[X \setminus Y])$, the key point is to show that $\Delta(\partial B(\Gamma))$ is homotopy equivalent to $\text{Cliq}(\Gamma)$. We sketch the argument from [75, Theorem 4.16], where the reader can find the details. Since the stabilizer of each element in $B(\Gamma)$ is commutative, any set of elements of $B(\Gamma)$ with a common lower bound has a meet. The maximal elements of $\partial B(\Gamma)$ are the elements of V . A collection of elements of V have a common lower bound if and only if they mutually commute, which occurs if and only if they form a clique in Γ . Rota's cross-cut theorem (Theorem 3.1) then provides the desired homotopy equivalence. $\square$

In light of Proposition 3.14 it is natural to formulate the following definition. Let $\mathcal{P}$ be a class of posets closed under isomorphism, e.g., CW posets or contractible posets. We say that a left regular band B is a $\mathcal{P}$ -left regular band if $B_{\geq X} \in \mathcal{P}$ for all $X \in \Lambda(B)$. For example, the set of covectors of an (affine) oriented matroid is a contractible CW left regular band. A *spherical left regular band* will then mean a left regular band B such that $B_{\geq X}$ is the face poset of a regular CW decomposition of a $\dim \Delta(B_{\geq X})$ -sphere for all $X \in \Lambda(B)$. For example, if $\mathcal{A}$ is a real or complex hyperplane arrangement, then $\mathcal{F}(\mathcal{A}) \setminus \{0\}$ is a spherical left regular band, as is $\mathcal{L} \setminus \{0\}$ for any oriented matroid $(E, \mathcal{L})$.

The next two propositions give us easy ways to create new CW left regular bands from old ones.

Proposition 3.15. *If B, B' are (connected/contractible) CW left regular bands, then so is $B \times B'$.*

Proof. First note that $\Lambda(B \times B') \cong \Lambda(B) \times \Lambda(B')$ and that if $(X, Y) \in \Lambda(B \times B')$, then $(B \times B')_{\geq (X, Y)} = B_{\geq X} \times B_{\geq Y}$. Thus it suffices to show that the poset $B \times B'$ is a (connected/contractible) CW poset. It is a CW poset by Proposition 3.11. As products of connected/contractible spaces are connected/contractible and $\|\Delta(B \times B')\| \cong \|\Delta(B)\| \times \|\Delta(B')\|$ (cf. [22, Equation (9.6)]), the result follows. $\square$

Proposition 3.16. *If B' is a spherical left regular band and B is a connected CW left regular band (monoid), then $B' * B$ is a connected CW left regular band (monoid) with support lattice $\Lambda(B') * \Lambda(B)$. In particular, $S(B)$ is a connected CW left regular band (monoid) with support lattice $\Lambda(B) \cup \{-\infty\}$ where $-\infty$ is an external minimum.*

Proof. It is clear that $\Lambda(B' * B) \cong \Lambda(B') * \Lambda(B)$ and we identify these two semilattices. If $X \in \Lambda(B)$, then $(B' * B)_{\geq X} = B_{\geq X}$ and hence is a connected CW poset. If $X \in \Lambda(B')$, then $(B' * B)_{\geq X} = B'_{\geq X} * B$ and hence is a CW poset by Proposition 3.10. Moreover, the join of a sphere with a connected space is always connected. The monoid statements are clear. $\square$

In particular, if $\mathcal{A}, \mathcal{A}'$ are real or complex hyperplane arrangements, then $(\mathcal{F}(\mathcal{A}) \setminus \{0\}) * \mathcal{F}(\mathcal{A}')$ is a connected CW left regular band. We now give another example of a family of connected CW left regular band monoids, which is essentially from [27].

Example 3.17 (Ladders). Let $L_n = S^n(\{0\})$ be the n -fold suspension of the trivial left regular band. So $L_1 = L$ and $L_2 = \tilde{L}$. It is easy to see that $L_n \cong \{0, \pm 1, \dots, \pm n\}$ with the product given by

$$xy = \begin{cases} y, & \text{if } |x| < |y| \\ x, & \text{if } |x| \geq |y|. \end{cases}$$

Observe that $\Lambda(L_n) \cong \{0, \dots, n\}$ ordered by $\geq$ and the support map is given by $\sigma(x) = |x|$. We call the L_n *ladders* because the Hasse diagram of $L_n \setminus \{0\}$

is a ladder with n rungs. The partial order on L_n is given by $x < y$ if and only if $|x| > |y|$. It follows from Proposition 3.16 that L_n is a CW poset of dimension n and hence $\|\Delta(L_n)\|$ is homeomorphic to a closed n -ball. See Figure 12 for the Hasse diagram and the corresponding cell decomposition of the ball for $n = 3$.




FIGURE 12. The Hasse diagram of L_3 and corresponding cell decomposition of the 3-ball

The following lemma, providing an alternative characterization of connectedness for left regular bands, will be useful at a later juncture.

Lemma 3.18. *Let B be a left regular band and $X \in \Lambda(B)$. Define a graph $\Gamma(X)$ as follows. The vertex set of $\Gamma(X)$ is $L_X = \{b \in B \mid Bb = X\}$. Two vertices $x, y \in L_X$ are adjacent if they have a common upper bound, that is, $bx = x$ and $by = y$ for some $b \in B$ (necessarily belonging to $B_{\geq X}$). Then $\Delta(B_{\geq X})$ is connected if and only if $\Gamma(X)$ is connected.*

Proof. Suppose first that $\Gamma(X)$ is connected. Fix $e_X \in L_X$. First note that if $a \in B_{\geq X} \setminus L_X$, then $\{a, ae_X\}$ is an edge from a to ae_X in $\Delta(B_{\geq X})$. It thus suffices to show that any two elements of L_X are connected in $\Delta(B_{\geq X})$. If $x, y \in L_X$ are adjacent and $bx = x$, $by = y$ with $b \in B$, then $\{x, b\}$, $\{b, y\}$ is an edge path from x to y in $\Delta(B_{\geq X})$. We now deduce the connectivity of $\Delta(B_{\geq X})$ from the connectivity of $\Gamma(X)$.

Assume that $\Delta(B_{\geq X})$ is connected and that $x, y \in L_X$. Then there is an edge path from x to y in $\Delta(B_{\geq X})$. Consider a shortest edge path. If the path contains consecutive edges $\{a, b\}$ and $\{b, c\}$ where $a < b < c$ or $a > b > c$, then these two edges can be replaced by the single edge $\{a, c\}$, resulting in a shorter path. As x, y are minimal elements of $B_{\geq X}$, it follows that any minimal length edge path from x to y in $\Delta(B_{\geq X})$ is of the form

$$\{x_0, x_1\}\{x_1, x_2\} \cdots \{x_{n-2}, x_{n-1}\}\{x_{n-1}, x_n\}$$

with $x_0 = x$, $x_n = y$ and

$$x_0 < x_1 > x_2 < x_3 > \cdots > x_{n-2} < x_{n-1} > x_n.$$

As $a \geq ay \in L_X$ for all $a \in B_{\geq X}$, we deduce that

$$x_0 < x_1 > x_2y < x_3 > \cdots > x_{n-2}y < x_{n-1} > x_n$$

and so $x \sim x_2y \sim x_4y \sim \cdots \sim x_{n-2}y \sim y$ where $\sim$ denotes the adjacency relation of $\Gamma(X)$. This establishes that $\Gamma(X)$ is connected. $\square$

The graph constructed in Lemma 3.18 is implicit in [96] and is closely related to tope graphs of oriented matroids [26] and COMs [13]. We use the lemma to prove that strong elimination systems give rise to connected left regular bands.

Proposition 3.19. *Let $(E, \mathcal{L})$ be a strong elimination system. Then $\mathcal{L}$ is a connected left regular band.*

Proof. Let $X \in \Lambda(\mathcal{L})$. We prove that the graph $\Gamma(X)$ is connected. Let $x, y \in L_X$. We prove that x, y are in the same component of $\Gamma(X)$ by induction on $|S(x, y)|$. As $Z(x) = Z(y)$, if $S(x, y) = \emptyset$, then $x = y$ and there is nothing to prove. So assume that if $v, w \in L_X$ with $|S(v, w)| < |S(x, y)|$, then v, w are in the same component of $\Gamma(X)$. Let $e \in S(x, y)$. By (OM3), we can find $z \in \mathcal{L}$ such that $z_e = 0$ and $z_f = (xy)_f = (yx)_f$ for $f \in E \setminus S(x, y)$. Notice that since $Z(x) = Z(y)$, this implies that $z \in \mathcal{L}_{\geq X}$. Then we have that $zx, zy \in L_X$, $(zx)_f = x_f = y_f = (zy)_f$ for $f \in E \setminus S(x, y)$ and $(zx)_e = x_e$, $(zy)_e = y_e$. Therefore, $|S(x, zx)| < |S(x, y)|$ and $|S(y, zy)| < |S(x, y)|$, whence we may assume by induction that x, zx are in the same component of $\Gamma(X)$ and that y, zy are in the same component of $\Gamma(X)$. But z is a common upper bound of zx, zy and so zx and zy are adjacent in $\Gamma(X)$. We conclude that x, y are in the same component of $\Gamma(X)$, completing the proof. $\square$

3.5. CAT(0) cube complexes. Our next goal is to show that the face poset of every finite CAT(0) cube complex has the structure of a connected CW left regular band. We refer the reader to [118] for a survey of important developments in the theory of groups acting on CAT(0) cube complexes. In particular, they have played an important role in Agol's solution to Thurston's virtual fibering and virtual Haken conjectures [3], using heavily earlier work of Wise [118]. We follow here Abramenko and Brown [1, Appendix A.2] for cube complexes and the book of Bridson and Haefliger [29] for CAT(0) metric spaces. The unpublished monograph of Roller [94] is a good reference for combinatorially minded people on CAT(0) cube complexes. See also [10] for a more graph theoretic point of view. Independently, it is shown in [13] that CAT(0) cube complexes are lopsided systems (and hence left regular bands) and that certain, more general, CAT(0) zonotopal complexes are COMs. We have decided to leave here our original presentation, which was written and spoken about at various conferences in 2014–2015, and will discuss these further developments in the next subsection.

A (finite) *cube complex* is a regular CW complex K such that:

- (1) for each open n -cell e of K , the subcomplex carried by $\bar{e}$ is isomorphic to the standard n -cube (with its usual cell structure with face poset L^n);
- (2) the intersection of two closed cells, if non-empty, is again a closed cell.

Notice the second condition implies that if C is a cube all of whose vertices belong to a cube C' , then $C \subseteq C'$.

For example, if K is a simplicial complex, put $\widehat{\mathcal{P}(K)} = \mathcal{P}(K) \cup \{\emptyset\}$. Let

$$\mathcal{C}(K) = \|\Delta(\widehat{\mathcal{P}(K)})\| \cong \|K * v\|$$

where $v \notin K$. Each closed interval $[a, b]$ of $\widehat{\mathcal{P}(K)}$ is a boolean lattice and hence $\|\Delta([a, b])\|$ is homeomorphic to a cube. We take as our set of cubes for $\mathcal{C}(K)$ the cubes $\|\Delta([a, b])\|$ to give $\mathcal{C}(K)$ the structure of a cube complex.

If K is a cube complex and v is a vertex of K , then the *link* $\text{lk}_K(v)$ of v is the simplicial complex with vertices all edges of K incident on v and where a set of edges incident on v forms a simplex if the edges in question belong to a common cube. A *CAT(0) cube complex* is a simply connected cube complex such that the link of each vertex is a flag complex. By a theorem of Gromov [29, Theorem II.5.20] this is equivalent to K carrying a CAT(0) metric in which each n -cube C has a characteristic map which is simultaneously an isomorphism of CW complexes and an isometry from the n -cube $I^n = [0, 1]^n$ equipped with its Euclidean metric to C (and so in particular each edge length has 1). Let us briefly recall the definition and basic properties of a CAT(0) metric space. Details can be found [29, Section II.1].

If x, y are points of a metric space (X, d) , then a *geodesic* from x to y is a continuous map $\gamma: [0, \ell] \rightarrow X$ with $\gamma(0) = x$, $\gamma(\ell) = y$ and

$$d(\gamma(t_1), \gamma(t_2)) = |t_1 - t_2|$$

for all $t_1, t_2 \in [0, \ell]$ (and so, in particular, $\ell = d(x, y)$). A *geodesic metric space* is a metric space (X, d) in which any two points are connected by a geodesic.

A *geodesic triangle* Δ in (X, d) consists of three points $x, y, z \in X$ and geodesics γ_{xy} , γ_{xz} and γ_{yz} , where γ_{ab} is a geodesic from a to b . A *comparison triangle* for x, y, z is a triangle Δ' in the Euclidean plane $(\mathbb{R}^2, \|\cdot\|_2)$ with vertices x', y', z' with $d(a', b') = d(a, b)$ for $a, b \in \{x, y, z\}$ (such a triangle always exists and is unique up to isometry of labeled triangles). One says that a geodesic metric space X is a *CAT(0) metric space* if Δ is thinner than Δ' for all geodesic triangles Δ in X . Formally speaking, this means that if $a, b, c \in \{x, y, z\}$, if $p = \gamma_{ab}(t)$, $q = \gamma_{ac}(t')$ and if p', q' are the points of the edges $[a', b']$ and $[a', c']$, respectively, with $d(a, p) = \|a' - p'\|_2$ and $d(a, q) = \|a' - q'\|_2$, then $d(p, q) \leq \|p' - q'\|_2$; see Figure 13. CAT(0) metric spaces were popularized by Gromov and have played an important role in geometric group theory over the past 20 years.

The following theorem encompasses part of [29, Proposition II.1.4] and [29, Corollary II.1.5].

Theorem 3.20. *Let (X, d) be a CAT(0) metric space.*

- (1) *X is contractible.*
- (2) *For any points $x, y \in X$, there is a unique geodesic from x to y .*

A non-empty subspace C of a geodesic metric space (X, d) is *convex* if it contains all geodesics between points of C . In particular, if (X, d) is CAT(0),




FIGURE 13. Comparison triangle for a CAT(0) space

then C is convex if and only if it contains the unique geodesic between any two of its points. A convex subspace of a CAT(0) metric space is CAT(0) in the induced metric, cf. [29, Example II.1.15]. Intersections of convex subspaces, if non-empty, are again convex.

Example 3.21 (CAT(0) cone). Let K be a simplicial complex. The reader should check that the cube complex $\mathcal{C}(K)$ constructed above is CAT(0) if and only if K is a flag complex. In this case, $\mathcal{C}(K)$ is called the *CAT(0) cone* on K .

A graph is a CAT(0) cube complex (consisting of 0-cubes and 1-cubes) if and only if it is a tree. A simply connected square complex (pure 2-dimensional cube complex) is CAT(0) if and only if each internal vertex is surrounded by at least 4 squares because a graph is a flag complex if and only if it is triangle-free and the link of a vertex in a square complex is a graph. See Figure 14 for a CAT(0) square complex whose unique internal vertex has a link which is a 5-cycle.




FIGURE 14. A CAT(0) square complex with a 5-cycle link

Example 3.22 (Phylogenetic Tree Space). In [19], Billera, Holmes and Vogtman introduced a CAT(0) cube complex structure on the space of phylogenetic trees called *phylogenetic tree space*. Speyer and Sturmfels [106] later showed that phylogenetic tree space is the tropical Grassmannian. This

seems to be the most commonly studied CAT(0) cube complex outside of geometric group theory. Strictly speaking, phylogenetic tree space is not a finite cube complex, and so we shall work with the truncation of this complex in which the edge lengths of interior edges of phylogenetic trees are at most 1.

Following the formalism of [19], the *phylogenetic tree space* $\mathcal{T}_n$ will be defined as follows. We consider labelled n -trees with $n \geq 2$. These are trees with $n + 1$ leaves labelled by $\{0, \dots, n\}$ and each non-leaf has valence at least 3. We think of the leaf labelled by 0 as the root. See Figure 15 for the case $n = 3$.




FIGURE 15. The four labelled 3-trees

By an *interior edge* of a labelled n -tree, we mean an edge not incident on a leaf. A *metric* labelled n -tree is one whose interior edges are assigned positive lengths in the interval $(0, 1]$. (Note that in [19] arbitrary positive lengths are allowed.) See Figure 16. There is an obvious notion of isomorphism of




FIGURE 16. A metric tree

n -labelled metric trees as graph isomorphisms preserving the labellings and the edge lengths. We shall identify isomorphic metric trees.

It will be convenient to allow n -labelled trees to have interior edges of length 0 and to identify such a tree as being isomorphic to the metric tree obtained by contracting to a point each edge of length 0. In this way, each metric tree can be represented by a trivalent tree (all inner nodes have degree 3) with $n - 2$ interior edges with lengths in the interval $[0, 1]$. See Figure 17.





FIGURE 17. A degenerate metric tree

Let us say an n -labelled tree T' is a *degeneration* of an n -labelled tree T if it is obtained from T by contracting some number of interior edges to a point. For example, the n -labelled tree with no interior edges is a degeneration of every n -labelled tree.

Fix a trivalent n -labelled tree T . Then we can identify the set C_T of isomorphism classes of metric trees obtained from assigning lengths from $[0, 1]$ to the $n - 2$ interior edges with the cube $[0, 1]^{n-2}$. The proper faces of C_T come from allowing some interior edges to have length 0 and correspond bijectively to n -labelled trees which are degenerations of T . If T, T' are two trivalent trees with a common degeneration, then we isometrically glue together the faces of C_T and $C_{T'}$ corresponding to their common degenerations. The resulting cube complex $\mathcal{T}_n$ is CAT(0). The metric n -labelled tree with no interior edges is a cone point for $\mathcal{T}_n$. See Figure 18 for $\mathcal{T}_3$.

FIGURE 18. Phylogenetic tree space $\mathcal{T}_3$

In fact, $\mathcal{T}_n$ is the CAT(0) cone on the following flag complex. Call a partition of $\{0, \dots, n\}$ into two subsets *thick* if both parts have at least two elements. If T is an n -labelled tree and e is an interior edge of T , then $T - \{e\}$ has two connected components which induce a thick partition on $\{0, \dots, n\}$. Define a graph with vertices the thick partitions. If $\{A, B\}$ and $\{A', B'\}$ are thick partitions, then we connect them by an edge if

$$A \subseteq A' \text{ or } A \subseteq B' \text{ or } B \subseteq A' \text{ or } B \subseteq B'.$$

Let L_n be the clique complex of this graph. Then the simplices of L_n are in bijection with n -labelled trees. Given an n -labelled tree T , one obtains a simplex by considering all thick partitions obtained by removing some interior edge from T . The facets of L_n are in bijection with the trivalent trees. It is then easy to check that $\mathcal{T}_n$ is the CAT(0) cone on L_n and that L_n is the link of the origin. See [19, 106] for details.

Fundamental to the theory of CAT(0) cube complexes is the notion of a hyperplane.

A *midcube* of I^n is the intersection of I^n with one of the affine hyperplanes

$$H_i = \left\{ x \in \mathbb{R}^n \mid x_i = \frac{1}{2} \right\}.$$

If e is an edge of I^n , then the midcube dual to e is the intersection of I^n with the perpendicular bisector of e . See Figure 19.




FIGURE 19. A midcube

Midcubes of an n -cube of a CAT(0) cube complex can be defined via the characteristic map identifying it isometrically with I^n . Let X be a CAT(0) cube complex and let $\sim$ be the smallest equivalence relation on the edges of X such that opposite edges of any square of X are equivalent. If e is an edge of a CAT(0) cube complex, then the *hyperplane* H_e of X dual to e is the union of all midcubes dual to edges in the $\sim$ -equivalence class of e . Equivalently, a hyperplane is a maximal connected subspace which is a union of midcubes. See [94, 118] for details. Figure 20 depicts a hyperplane in a CAT(0) cube complex.




FIGURE 20. A hyperplane in a CAT(0) cube complex

We need the following fundamental facts about CAT(0) cube complexes. See [118, Theorem 2.13].

Theorem 3.23. *Let K be a $CAT(0)$ cube complex.*

- (1) *Each hyperplane of K is convex and hence a $CAT(0)$ cube complex (whose cubes are the faces of the midcubes that comprise it).*
- (2) *If C is an n -cube of K , then each of the n -midcubes of C lie on different hyperplanes of K .*
- (3) *If H is a hyperplane, then $K \setminus H$ has two connected components called half-spaces.*

Although we do not use it, we mention the following important theorem obtained, independently by Chepoi [42] and Roller [94]. We recall that a *median graph* is a connected graph Γ such that if x, y, z are vertices of Γ and γ_{ab} are geodesic edge paths from a to b for $a, b \in \{x, y, z\}$, then $|\gamma_{xy} \cap \gamma_{xz} \cap \gamma_{yz}| = 1$. For example, the hypercube graph is a median graph, as is the Hasse diagram of any distributive lattice.

Theorem 3.24. *A $CAT(0)$ cube complex K is uniquely determined by its 1-skeleton, which is a median graph. Namely, the cubes of K are obtained by filling in with an n -cube each induced subgraph of K^1 which is isomorphic to the 1-skeleton of an n -cube.*

Conversely, if Γ is a median graph, then the cube complex obtained by filling in 1-skeleta of induced hypercube subgraphs is a $CAT(0)$ cube complex.

Let K be a $CAT(0)$ cube complex with hyperplanes $H_1, \dots, H_n$. We define the *intersection semilattice* $\mathcal{L}(K)$ to consist of all $H_{i_1} \cap \dots \cap H_{i_k} \neq \emptyset$, where if $k = 0$, then we interpret the intersection to be K . We order it by reverse inclusion. Note that the semilattice operation takes two elements X, Y of $\mathcal{L}(K)$ and brings it to the intersection of all hyperplanes containing both X, Y (with the empty intersection being K). Hyperplanes in $CAT(0)$ cube complexes are convex in the $CAT(0)$ metric by Theorem 3.23 and hence each element $X \in \mathcal{L}(K)$ is convex and therefore a $CAT(0)$ cube complex in its own right. The cubes of X are exactly the non-empty intersections of cubes of K with the subspace X . One can recover the set $A = \{H_{i_1}, \dots, H_{i_k}\}$ of hyperplanes from the vertices of X , namely each vertex is the midpoint of a k -cube whose midcubes are precisely its intersections with the elements of A . In other words, each element X of $\mathcal{L}(K)$ can be uniquely expressed as an intersection of hyperplanes. In general, if C is an n -cube of K with $C \cap X \neq \emptyset$, then $C \cap X$ is an $(n - k)$ -cube of X .

The set of hyperplanes in a $CAT(0)$ cube complex satisfies the *Helly property*: any collection of hyperplanes with pairwise non-empty intersections has non-empty intersection; see [42, Page 143] for a proof. Recall that if $\mathcal{F} = \{X_1, \dots, X_n\}$ is a collection of subsets of a set X , then the *nerve* of $\mathcal{F}$ is the simplicial complex $\mathcal{N}(\mathcal{F})$ with vertex set $\mathcal{F}$ and where $\{X_{i_0}, \dots, X_{i_q}\}$ is a q -simplex if $X_{i_0} \cap \dots \cap X_{i_q} \neq \emptyset$. Clearly, $\mathcal{N}(\mathcal{F})$ is a flag complex if and only if $\mathcal{F}$ has the Helly property.

Proposition 3.25. *Let K be a finite $CAT(0)$ cube complex with set $\mathcal{H}$ of hyperplanes. Then the intersection semilattice $\mathcal{L}(K)$ can be identified with*

the face poset of the nerve $\mathcal{N}(\mathcal{H})$ of $\mathcal{H}$, including an empty face. Moreover, $\mathcal{N}(\mathcal{H})$ is a flag complex and every finite flag complex comes about this way.

Proof. We already saw above that each element X of $\mathcal{L}(K)$ can be uniquely expressed as an intersection of a subset A_X of hyperplanes. The mapping $X \mapsto A_X$ gives an isomorphism of $\mathcal{L}(K)$ with $\mathcal{P}(\mathcal{N}(\mathcal{H})) \cup \{\emptyset\}$ (sending K to the empty face). Since $\mathcal{H}$ has the Helly property, $\mathcal{N}(\mathcal{H})$ is a flag complex. Conversely, the reader should verify that if X is a flag complex and K is the CAT(0) cone on X , then $\mathcal{N}(\mathcal{H})$ is isomorphic to X (where $\mathcal{H}$ is the set of hyperplanes of K). $\square$

One can consider the vertex set K^0 of a CAT(0) cube complex as a metric space with the path metric d_p . The distance $d_p(v, w)$ between two vertices v, w is the length of a shortest edge path between them in the 1-skeleton K^1 . Let $S(v, w)$ be the set of hyperplanes H separating v and w , meaning that v and w are in different components of $K \setminus H$. Then it is known that

$$d_p(v, w) = |S(v, w)|.$$

In particular, two vertices are connected by an edge if and only if they are separated by exactly one hyperplane, which is necessarily the hyperplane dual to that edge. See [118, Remark 2.14] or [94].

We shall need the following lemma stating that a cube can be separated from each vertex outside of it by a hyperplane.

Lemma 3.26. *Let K be a CAT(0) cube complex, let v be a vertex of K and let C be a cube of K not containing v . Then there is a hyperplane H separating v from C , that is, v and C are contained in different connected components of $K \setminus H$.*

Proof. Let w be a vertex of C at a minimal distance from v with respect to the path metric d_p . Then $|S(v, w)| = d_p(v, w) \geq 1$. Let $H \in S(v, w)$. Suppose first that $H \cap C \neq \emptyset$. Then there is a unique edge e of C dual to H and incident on w . Let w' be the other endpoint of e . Then w' is in the same component of $K \setminus H$ as v and so $H \notin S(v, w')$. If $H' \in S(v, w')$, then because $S(w, w') = \{H\}$, we must have that w, w' are in the same component of $K \setminus H'$ and hence $H' \in S(v, w)$. We conclude that $S(v, w') \subseteq S(v, w) \setminus \{H\}$ and so $d_p(v, w') < d_p(v, w)$, a contradiction. Thus $H \cap C = \emptyset$. Since C is connected and w is in a different component of $K \setminus H$ than v , we conclude that H separates v and C . $\square$

Another important fact is that every finite CAT(0) cube complex embeds as a subcomplex of a hypercube; this was first proved by Bandelt in the language of median graphs [9]. We provide a proof here using covectors for both completeness, and to make the semigroup structure transparent. The final item of the next theorem owes a great deal to Daniel Wise (private communication).

Theorem 3.27. *Let K be a finite $CAT(0)$ cube complex with hyperplanes $\{H_1, \dots, H_d\}$. Fix, for each H_i , positive and negative half-spaces H_i^+ and H_i^- of $K \setminus H_i$. To each cube C of K assign the covector $\tau(C) \in L^d$ given by*

$$\tau(C)_i = \begin{cases} +, & \text{if } C \subseteq H_i^+ \\ -, & \text{if } C \subseteq H_i^- \\ 0, & \text{if } C \cap H_i \neq \emptyset. \end{cases}$$

Let $\mathcal{F}(K)$ be the set of all $\tau(C)$ such that C is a cube of K . Then the following hold.

- (1) $\tau: \mathcal{P}(K) \longrightarrow L^d$ is an order embedding.
- (2) $\mathcal{F}(K)$ is a right ideal in L^d .
- (3) K is isomorphic to a subcomplex of $[-1, 1]^d$.
- (4) $\Lambda(\mathcal{F}(K)) \cong \mathcal{L}(K)$.
- (5) If $X \in \mathcal{L}(K)$, then $\mathcal{F}(K)_{\geq X} \cong \mathcal{F}(X)$.

In particular, $\mathcal{F}(K)$ is a connected CW left regular band.

Proof. Clearly if C is a face of C' , then $\tau(C) \leq \tau(C')$. Suppose that $\tau(C) \leq \tau(C')$. Let v be a vertex of C . If $v \notin C'$, then there is a hyperplane H_i separating v from C' by Lemma 3.26. But then $\tau_i(C') = -\tau_i(v) \neq 0$ and $\tau_i(v) \leq \tau_i(C) \leq \tau_i(C')$, a contradiction. Thus each vertex of C is a vertex of C' , whence $C \subseteq C'$. Therefore, τ is an order embedding.

Suppose that C is an n -cube of K . Then exactly n hyperplanes intersect C and they intersect it in its midcubes by Theorem 3.23. Thus $\tau(C)$ has exactly n zeroes and therefore 3^n faces. But C also has 3^n faces and τ is an order embedding. Thus every face of $\tau(C)$ is in the image of τ and so $\mathcal{F}(K)$ is a lower set, that is, a right ideal of L^d .

Since $\mathcal{P}(K)$ is order isomorphic to a lower set of $L^d = \mathcal{P}([-1, 1]^d)$, it follows that K is isomorphic to a subcomplex of $[-1, 1]^d$ because regular cell complexes are determined up to isomorphism by their face posets.

Because $\mathcal{F}(K)$ is a right ideal in L^d , it follows that $\sigma: \mathcal{F}(K) \longrightarrow \Lambda(\mathcal{F}(K))$ can be identified with the zero set map $Z: \mathcal{F}(K) \longrightarrow Z(\mathcal{F}(K))$. If $A \subseteq \{1, \dots, n\}$, then $A = Z(\tau(C))$ if and only if $A = \{i \mid H_i \cap C \neq \emptyset\}$ and hence $\bigcap_{i \in A} H_i \in \mathcal{L}(K)$. Conversely, if $X = \bigcap_{i \in A} H_i \in \mathcal{L}(K) \neq \emptyset$ choose a vertex v of X . By definition of the cube complex structure on X , there is a $|A|$ -cube C of K with $C \cap X = \{v\}$. Then $Z(\tau(C)) = A$. Thus $\Lambda(\mathcal{F}(K))$ can be identified with $\mathcal{L}(K)$ as a meet semilattice.

If $A \in \Lambda(\mathcal{F}(K))$ and $X = \bigcap_{i \in A} H_i \in \mathcal{L}(K) \neq \emptyset$ are as above, then the cubes C with $A \subseteq Z(\tau(C))$ are exactly the cubes C of K meeting X . But the cubes of X are precisely the non-empty intersections of cubes of K with X . Thus the map $\tau(C) \mapsto C \cap X$ gives an order isomorphism between $\mathcal{F}(K)_{\geq A}$ and the face poset $\mathcal{P}(X)$ of X . In particular, $\mathcal{F}(K)_{\geq A}$ is a connected CW poset. But more is true. The hyperplanes of X are precisely the $H'_i = H_i \cap X$ with $i \notin A$ and so if we choose the positive (respectively, negative) half-space of H'_i to be $H_i^+ \cap X$ (respectively, $H_i^- \cap X$) and let $\tau': \mathcal{P}(X) \longrightarrow L^{[n] \setminus A}$

be the corresponding embedding, then the map $\tau(C) \mapsto \tau'(C \cap X)$ is a semigroup isomorphism between $\mathcal{F}(K)_{\geq A}$ and $\mathcal{F}(X)$. $\square$

We call $\mathcal{F}(K)$ the *face semigroup* of the CAT(0) cube complex K . We shall identify $\mathcal{F}(K)$ with the set of faces of K . The reader should verify that the semigroup structure does not depend on the choice of orientation for the hyperplanes. In fact, there is the following geometric interpretation of the product. If C, C' are cubes, then $C \cdot C'$ is the face of C on which the mapping $x \mapsto d_p(x, C')$ is minimized.

3.6. CAT(0) zonotopal complexes. By a *Coxeter zonotope* we mean a zonotope (called an *even polyhedron* in [61]) that is symmetric around the perpendicular bisector, or *mid-hyperplane*, of each edge. A zonotope is combinatorially isomorphic to a Coxeter zonotope if and only if it is a W -permutohedron for a finite Coxeter group W , that is, the zonotope dual to a Coxeter hyperplane arrangement $\mathcal{A}_W$. Of course, the n -cube is the Coxeter zonotope corresponding to the finite right-angled Coxeter group $(\mathbb{Z}/2\mathbb{Z})^n$. See [13, 61] for details.

By a *zonotopal complex*, we mean a polyhedral complex whose cells are zonotopes. So it is a collection of zonotopes closed under taking faces and such that the intersection of any two zonotopes in the collection is a common face of them both. A *CAT(0) zonotopal complex* is a zonotopal complex equipped with a CAT(0) metric. If K is a CAT(0) zonotopal complex, each of whose zonotopes is isometric to a Coxeter zonotope, then one can define a hyperplane in K similarly to the CAT(0) cube complex case. Namely, consider the equivalence relation on the set of edges generated by pairs of edges that are on opposite sides of a two-dimensional face (two-dimensional zonotopes always have an even number of sides). The union of all mid-hyperplanes from an equivalence class of edges gives a hyperplane in K and one has that hyperplanes do not self-intersect and have many of the same properties of hyperplanes in CAT(0) cube complexes such as having exactly two connected components in their complement and being convex. They are CAT(0) zonotopal complexes in their own right with Coxeter zonotopes as cells. See [13, Section 11] for details.

If K is a CAT(0) zonotopal complex with Coxeter zonotopes as cells, then one can define a collection of covectors $\mathcal{L} \subset L^E$, where E is the set of hyperplanes (which we assume have an orientation selected for their two half-spaces), via the same recipe we used for CAT(0) cube complexes. If Z is a face of K , we assign it the covector $\tau(Z)$ with

$$\tau(Z)_e = \begin{cases} 0, & \text{if } e \cap Z \neq \emptyset \\ +, & \text{if } Z \subseteq e^+ \\ -, & \text{if } Z \subseteq e^- \end{cases}$$

where e^+ and e^- denote the positive and negative half-spaces associated with the hyperplane e . The following beautiful theorem was proved in [13, Section 11].

Theorem 3.28. *If K is a $CAT(0)$ zonotopal complex whose cells are Coxeter zonotopes, then the collection of covectors defined above provides the face poset of K with the structure of a COM. Moreover, each contraction of K is isomorphic to the semigroup of covectors of an intersection of hyperplanes, which is again a $CAT(0)$ zonotopal complex whose cells are Coxeter zonotopes.*

We remark that the support semilattice for a $CAT(0)$ zonotopal complex with Coxeter zonotope cells is more complicated than in the cube complex case because the intersection of a zonotope with a subset of hyperplanes does not determine the subset. For example, the regular hexagon is the Coxeter zonotope associated to the symmetric group S_3 . There are 3 hyperplanes, but any two of them have the same intersection point: the center of the hexagon.

4. ALGEBRAS

Let R be a ring, not necessarily unital. We recall the definition of the Jacobson radical of R [95]. A left ideal L of R is said to be *modular* if there is an element $a \in R$ such that $r - ra \in L$ for all $r \in R$. Note that if R has a right identity, in particular, if R is unital, then every left ideal is modular (take a to be the right identity). By a *maximal modular left ideal* is meant a maximal proper modular left ideal. One can define (maximal) modular right ideals analogously. The (*Jacobson*) *radical* $\text{rad}(R)$ is the intersection of all maximal modular left ideals. It is a two-sided ideal and it is also the intersection of all maximal modular right ideals. If R has a right identity (e.g. if R is unital), then $\text{rad}(R)$ is the intersection of all maximal left ideals of R or, equivalently is the intersection of all annihilators of simple left R -modules. Every nilpotent ideal of R is contained in the radical. A ring R is called *semiprimitive* if $\text{rad}(R) = 0$. If R is semiprimitive and finite dimensional over a field, then R is automatically unital and semisimple [95]. A commutative ring with unit is semiprimitive if and only if it is a subdirect product of fields.

In this text, module unmodified shall always means left module. If we want to discuss right modules we shall explicitly say so. Sometimes, we shall write “left module” for emphasis. A module over a unital ring is called *unitary* if $1M = M$.

Let us remark that if R is a ring with a right identity e , then R is a projective left module, as is any direct summand in R . Indeed, $\text{Hom}_R(R, M) \cong eM$ for any R -module M and the functor $M \mapsto eM$ is obviously exact.

4.1. Finite dimensional algebras. The reader is referred to [6, 8, 15, 48] for background on the modern theory of finite dimensional algebras. We

summarize here what we need for this paper. For this subsection, we fix a field $\mathbb{k}$ and all $\mathbb{k}$ -algebras are assumed unital and finite dimensional (except path algebras of non-acyclic quivers). All modules will be assumed unitary. A $\mathbb{k}$ -algebra A is said to be *split* if its semisimple quotient $A/\text{rad}(A)$ is isomorphic to a direct product of matrix algebras over $\mathbb{k}$ or, equivalently, each simple A -module is absolutely irreducible. In particular, if $\mathbb{k}$ is algebraically closed, then all finite dimensional $\mathbb{k}$ -algebras are split.

A $\mathbb{k}$ -algebra is *basic* if $A/\text{rad}(A)$ is isomorphic to a direct product of division rings. Consequently, A is split basic if and only if $A/\text{rad}(A) \cong \mathbb{k}^n$ for some $n \geq 1$ or, in other words, A is split basic if and only if every simple A -module is one-dimensional. Every split finite dimensional algebra over $\mathbb{k}$ is Morita equivalent to a unique (up to isomorphism) split basic algebra. Therefore, the representation theory of finite dimensional algebras over an algebraically closed field can be reduced to the case of split basic algebras. Recall that two algebras are *Morita equivalent* if their module categories are equivalent.

An A -module M is *indecomposable* if it cannot be expressed as a direct sum of proper submodules. The Krull-Schmidt theorem guarantees that every finite dimensional A -module can be expressed as a direct sum of indecomposables and the number of indecomposable summands isomorphic to a given indecomposable module is independent of the decomposition. In particular, the regular A -module can be expressed as

$$A = P_1 \oplus \cdots \oplus P_r \quad (4.1)$$

where the P_i are projective indecomposable modules and, moreover, each projective indecomposable A -module is isomorphic to some P_i . Furthermore, we can find a complete set of orthogonal primitive idempotents $e_1, \dots, e_r$ such that $P_i = Ae_i$ for $i = 1, \dots, r$ and all complete sets of orthogonal primitive idempotents arise in this way. Moreover, any two complete sets of orthogonal primitive idempotents are conjugate by an element of the group of units of the algebra. Recall that a non-zero idempotent $e \in A$ is *primitive* if $e = e_1 + e_2$ with e_1, e_2 orthogonal idempotents implies $e_1 = 0$ or $e_2 = 0$.

The *radical* $\text{rad}(M)$ of a finite dimensional A -module M is the intersection of all maximal submodules of M . One has the equality

$$\text{rad}(M) = \text{rad}(A)M.$$

The quotient $M/\text{rad}(M)$ is semisimple and is called the *top* of M . Recall that a module is *semisimple* if it is a direct sum of simple modules. An important fact about finite dimensional algebras is that if P is a projective indecomposable module, then $P/\text{rad}(P)$ is simple. Moreover, if P and Q are projective indecomposable modules, then $P \cong Q$ if and only if $P/\text{rad}(P) \cong Q/\text{rad}(Q)$. In addition, if (4.1) is a decomposition of A into projective indecomposables, then

$$A/\text{rad}(A) \cong P_1/\text{rad}(P_1) \oplus \cdots \oplus P_r/\text{rad}(P_r)$$

and every simple A -module is isomorphic to one of the form $P_i/\text{rad}(P_i)$. Consequently, there is a bijection between isomorphism classes of projective indecomposable modules and simple modules.

We recall that a surjective homomorphism $\varphi: P \rightarrow M$ from a finite dimensional projective A -module P to a module M is a *projective cover* if $\ker \varphi \subseteq \text{rad}(P)$. An injective homomorphism $\varphi: M \rightarrow I$ where I is a finite dimensional injective module is called an *injective envelope* (or *injective hull*) if every non-zero submodule of I intersects $\varphi(M)$. Each projective indecomposable module is the projective cover of its simple top and each injective indecomposable module is the injective envelope of its simple socle. Recall that the *socle* of a module is its largest semisimple submodule.

Let $e_1, \dots, e_r$ be a complete set of orthogonal primitive idempotents. Then each module $I_j = \text{Hom}_{\mathbb{k}}(e_j A, \mathbb{k})$ is an injective indecomposable module with simple socle isomorphic to $(A/\text{rad } A)e_j$. Moreover, $I_j \cong I_k$ if and only if $(A/\text{rad } A)e_j \cong (A/\text{rad } A)e_k$ and every injective indecomposable A -module is isomorphic to one of the form I_j with $1 \leq j \leq r$. Thus isomorphism classes of injective indecomposable modules and simple modules are also in bijection.

A $\mathbb{k}$ -algebra A is said to be *hereditary* if each left ideal is a projective module or, equivalently, submodules of projective modules are projective. Alternatively, hereditary algebras are algebras of global dimension at most one. Recall that the *projective dimension* of a module M is the shortest length of a projection resolution

$$P_\bullet \rightarrow M = \cdots \rightarrow P_q \rightarrow P_{q-1} \rightarrow \cdots \rightarrow P_1 \rightarrow P_0 \rightarrow M.$$

The *global dimension* $\text{gl. dim } A$ of A is the supremum of the projective dimensions of all A -modules. When A is finite dimensional, $\text{gl. dim } A$ is the maximum projective dimension of a simple A -module [6, Theorem A.4.8]. Alternatively, $\text{gl. dim } A$ is the largest n such that $\text{Ext}_A^n(S, S') \neq 0$ for some simple A -modules S, S' .

A *quiver* Q is a directed graph (possibly with loops and multiple edges), always assumed finite here. The vertex set is denoted Q_0 and the edge set Q_1 . More generally, Q_n will denote the set of (directed) paths of length n with $n \geq 0$ (where the distinction between empty paths and vertices is blurred). Denote by Q^{op} the quiver obtained from Q by reversing the orientation of each arrow.

The *path algebra* $\mathbb{k}Q$ of Q has a basis consisting of all paths in Q (including an empty path ε_v at each vertex) with the product induced by concatenation (where undefined concatenations are set equal to zero). Here we compose paths from right to left like category theorists: so if $p: v \rightarrow w$ and $q: w \rightarrow z$ are paths, then their composition is denoted qp . Note that $\mathbb{k}Q = \bigoplus_{n \geq 0} \mathbb{k}Q_n$. The identity of $\mathbb{k}Q$ is the sum of all empty paths.

For example, if Q consists of a single vertex with n loops, then $\mathbb{k}Q$ is the free algebra on n generators. If Q is a directed path with n nodes, then $\mathbb{k}Q$ is isomorphic to the algebra of $n \times n$ upper triangular matrices over $\mathbb{k}$.

If A is a finite dimensional algebra, then the primitive idempotents of the center $Z(A)$ are called *central primitive idempotents*. The central primitive idempotents of $\mathbb{k}Q$ are precisely the sums of empty paths ranging over a connected component of Q . If f is a central idempotent, then $Af = fAf$ is a unital $\mathbb{k}$ -algebra with identity f . There is a unique complete set of orthogonal central primitive idempotents $f_1, \dots, f_n$ and, moreover,

$$A \cong Af_1 \times \cdots \times Af_n$$

as a $\mathbb{k}$ -algebra. The algebras Af_j are called the *blocks* of A .

An important theorem of Gabriel is that a split basic algebra is hereditary if and only if it is isomorphic to $\mathbb{k}Q$ for some acyclic quiver Q . More generally, Gabriel showed that every split basic algebra has a quiver presentation. A *bound quiver* (Q, I) consists of a quiver Q and an admissible ideal I of $\mathbb{k}Q$. To define an admissible ideal, let $J = \bigoplus_{n \geq 1} \mathbb{k}Q_n$ be the *arrow ideal* of $\mathbb{k}Q$, that is, the ideal generated by all paths of length one. Then an ideal I is *admissible* if $J^n \subseteq I \subseteq J^2$ for some $n \geq 2$. The algebra $A = \mathbb{k}Q/I$ will be a finite dimensional split basic $\mathbb{k}$ -algebra. Note that $\text{rad}(A) = J/I$ and the cosets $\varepsilon_v + I$ with $v \in Q_0$ form a complete set of orthogonal primitive idempotents. In particular, if Q is acyclic, then $\{0\}$ is admissible, $\text{rad}(\mathbb{k}Q) = J$ and the empty paths ε_v with $v \in Q_0$ form a complete set of orthogonal primitive idempotents of $\mathbb{k}Q$.

Conversely, every finite dimensional split basic algebra A is the algebra of a bound quiver for a unique quiver $Q(A)$ (the admissible ideal I is not unique). The vertices of $Q(A)$ are the (isoclasses of) simple A -modules. If S, S' are simple A -modules, then the number of directed edges $S \rightarrow S'$ in $Q(A)$ is $\dim_{\mathbb{k}} \text{Ext}_A^1(S, S')$.

To explain how the admissible ideal I is obtained, fix a complete set E of orthogonal primitive idempotents for A and let $e_S \in E$ be the idempotent with $S \cong (A/\text{rad}(A))e_S$. One has that

$$\text{Ext}_A^1(S, S') \cong e_{S'}[\text{rad}(A)/\text{rad}^2(A)]e_S$$

as $\mathbb{k}$ -vector spaces [6, 15]. One can define a homomorphism $\mathbb{k}Q(A) \rightarrow A$ by sending the empty path at S to e_S and by choosing a bijection of the set of arrows $S \rightarrow S'$ with a subset of $e_{S'}\text{rad}(A)e_S$ mapping to a basis of $e_{S'}[\text{rad}(A)/\text{rad}^2(A)]e_S$. The map is extended to paths of length greater than one in the obvious way. Such a homomorphism is automatically surjective and the kernel will be an admissible ideal. We remark that if there is no path of length greater than one from S to S' in $Q(A)$, then $e_{S'}\text{rad}^2(A)e_S = 0$ because $\text{rad}(A) = J/I$. Similarly, if there is no path at all from S to S' through S'' , then

$$e_{S'}\text{rad}(A)e_{S''}\text{rad}(A)e_S = e_{S'}(J/I)e_{S''}(J/I)e_S = 0.$$

Finally, we remark that if $S \neq S'$, then $e_{S'}Ae_S = e_{S'}\text{rad}(A)e_S$ [6, 15].

A *system of relations* for an admissible ideal I is a finite subset

$$R \subseteq \bigcup_{v,w \in Q_0} \varepsilon_w I \varepsilon_v$$

generating I as an ideal. Every admissible ideal admits a system of relations R and the pair (Q, R) is called a *quiver presentation* of A . A system of relations R is called *minimal* if no proper subset of R generates the ideal I . The following result, giving a homological interpretation of the number of elements of a minimal generating set, encapsulates [28, 1.1 and 1.2]. We include a proof for completeness both because it will play an essential role in this text and to make things accessible for the diverse audience we are targeting (the exposition in the original reference is aimed at specialists in finite dimensional algebras).

Theorem 4.1 (Bongartz). *Let (Q, I) be a bound quiver with Q acyclic and let $A = \mathbb{k}Q/I$. For $v \in Q_0$, let $\mathbb{k}_v$ be the simple A -module $(A/\text{rad}(A))\varepsilon_v$. Then, for any minimal system of relations R , one has that the cardinality of $R \cap \varepsilon_w I \varepsilon_v$ is $\dim_{\mathbb{k}} \text{Ext}_A^2(\mathbb{k}_v, \mathbb{k}_w)$.*

Proof. For $v, w \in Q_0$, let $R(v, w) = R \cap \varepsilon_w I \varepsilon_v$. Let J be the arrow ideal of $\mathbb{k}Q$. We claim that

$$|R(v, w)| = \dim_{\mathbb{k}} \varepsilon_w [I/(JI + IJ)] \varepsilon_v. \quad (4.2)$$

First we prove that the cosets of the elements of $R(v, w)$ are linearly independent in $\varepsilon_w [I/(JI + IJ)] \varepsilon_v$. Suppose that this is not the case. Then there exists $\alpha_0 \in R(v, w)$ such

$$\alpha_0 \in \sum_{\alpha \in R(v, w) \setminus \{\alpha_0\}} \mathbb{k}\alpha + JI + IJ. \quad (4.3)$$

Let K be the ideal generated by $R \setminus \{\alpha_0\}$ and let I_0 be the ideal generated by α_0 . Then we deduce from (4.3) that $\alpha_0 \in K + JI_0 + I_0J$ and so $I_0 \subseteq K + JI_0 + I_0J$. We claim that, for all $n \geq 1$,

$$I_0 \subseteq K + \sum_{i=0}^n J^{n-i} I_0 J^i$$

where J^0 is interpreted as $\mathbb{k}Q$. The base case $n = 1$ is already handled. Assume it is true for n . Then we have

$$\begin{aligned} I_0 &\subseteq K + JI_0 + I_0J \subseteq K + JK + J \sum_{i=0}^n J^{n-i} I_0 J^i + KJ + \sum_{i=0}^n J^{n-i} I_0 J^i J \\ &\subseteq K + \sum_{i=0}^{n+1} J^{n+1-i} I_0 J^i. \end{aligned}$$

As J is nilpotent and $J^{n-i} I_0 J^i \subseteq J^n$, we conclude that $I_0 \subseteq K$. This contradicts the minimality of R . Therefore, the cosets of the elements of

$R(v, w)$ are linearly independent in $\varepsilon_w [I/(JI + IJ)] \varepsilon_v$. Next we show that they span this $\mathbb{k}$ -vector space.

Let $r \in \varepsilon_w I \varepsilon_v$. Then

$$r = \sum_{\alpha \in R} \sum_{i=1}^{n_\alpha} c_{\alpha,i} u_{\alpha,i} \alpha v_{\alpha,i}$$

with the $c_{\alpha,i} \in \mathbb{k}$ and the $u_{\alpha,i}$ and $v_{\alpha,i}$ paths. If $u_{\alpha,i}$ or $v_{\alpha,i}$ is non-empty, then $u_{\alpha,i} \alpha v_{\alpha,i} \in \varepsilon_w (JI + IJ) \varepsilon_v$. It follows that

$$r + \varepsilon_w (JI + IJ) \varepsilon_v = \sum_{\alpha \in R(v,w)} c_\alpha \alpha + \varepsilon_w (JI + IJ) \varepsilon_v$$

for some $c_\alpha \in \mathbb{k}$. This completes the proof of (4.2).

It remains to prove that

$$\dim_{\mathbb{k}} \text{Ext}_A^2(\mathbb{k}_v, \mathbb{k}_w) = \dim_{\mathbb{k}} \varepsilon_w [I/(JI + IJ)] \varepsilon_v.$$

Consider the exact sequence of $\mathbb{k}Q/I$ -modules

$$0 \longrightarrow (I/IJ) \varepsilon_v \xrightarrow{h} (J/IJ) \varepsilon_v \xrightarrow{j} (\mathbb{k}Q/I) \varepsilon_v \xrightarrow{k} \mathbb{k}_v \longrightarrow 0$$

where $h(x + IJ) = x + IJ$, $j(x + IJ) = x + I$ and $k(x + I) = x + J/I$. We write $s(e)$ for the source of an edge $e \in Q_1$ and $t(e)$ for the target. Then there is an isomorphism of $\mathbb{k}Q$ -modules

$$\bigoplus_{\{e \in Q_1 | s(e)=v\}} \mathbb{k}Q \varepsilon_{t(e)} \longrightarrow J \varepsilon_v$$

that takes $p \in \mathbb{k}Q \varepsilon_{t(e)}$ to pe . Thus

$$(J/IJ) \varepsilon_v \cong \mathbb{k}Q/I \otimes_{\mathbb{k}Q} J \varepsilon_v \cong \bigoplus_{\{e \in Q_1 | s(e)=v\}} (\mathbb{k}Q/I) \varepsilon_{t(e)}$$

is a projective $\mathbb{k}Q/I$ -module.

Let $M \cong (J/I) \varepsilon_v$ be the cokernel of h . Then we have exact sequences

$$0 \longrightarrow (I/IJ) \varepsilon_v \xrightarrow{h} (J/IJ) \varepsilon_v \longrightarrow M \longrightarrow 0$$

$$0 \longrightarrow M \longrightarrow (\mathbb{k}Q/I) \varepsilon_v \longrightarrow \mathbb{k}_v \longrightarrow 0$$

of $\mathbb{k}Q/I$ -modules with the middle terms projective modules. The long exact sequence for the Ext-functor then yields that

$$\text{Ext}_{\mathbb{k}Q/I}^2(\mathbb{k}_v, \mathbb{k}_w) \cong \text{Ext}_{\mathbb{k}Q/I}^1(M, \mathbb{k}_w) \cong \text{coker } h^*$$

where $h^*: \text{Hom}_{\mathbb{k}Q/I}((J/IJ) \varepsilon_v, \mathbb{k}_w) \longrightarrow \text{Hom}_{\mathbb{k}Q/I}((I/IJ) \varepsilon_v, \mathbb{k}_w)$ is the homomorphism induced by h . But if $f: (J/IJ) \varepsilon_v \longrightarrow \mathbb{k}_w$ is a homomorphism, then because I is admissible, and hence $I \subseteq J^2$, we conclude that $fh((I/IJ) \varepsilon_v) \subseteq f((J^2/IJ) \varepsilon_v) \subseteq (J/I)f((J/IJ) \varepsilon_v) \subseteq (J/I) \mathbb{k}_w = 0$ as $J/I = \text{rad}(\mathbb{k}Q/I)$. We conclude that $h^* = 0$ and so

$$\text{coker } h^* \cong \text{Hom}_{\mathbb{k}Q/I}((I/IJ) \varepsilon_v, \mathbb{k}_w) \cong \text{Hom}_{\mathbb{k}Q/I}([I/(JI + IJ)] \varepsilon_v, \mathbb{k}_w).$$

As $\mathbb{k}Q/J$ is semisimple, it follows that $\text{Hom}_{\mathbb{k}Q/J}([I/(JI + IJ)]\varepsilon_v, \mathbb{k}_w) \cong \text{Hom}_{\mathbb{k}Q/J}(\mathbb{k}_w, [I/(JI + IJ)]\varepsilon_v) \cong \varepsilon_w [I/(JI + IJ)]\varepsilon_v$. We conclude that $\text{Ext}_A^2(\mathbb{k}_v, \mathbb{k}_w) \cong \varepsilon_w [I/(JI + IJ)]\varepsilon_v$, completing the proof. $\square$

Obtaining a quiver presentation of a finite dimensional algebra is a crucial step in order to be able to apply the apparatus developed by modern representation theorists.

In [75, Theorem 3.10], the authors showed that for any acyclic quiver Q , there is a left regular band monoid B_Q with $\mathbb{k}B_Q \cong \mathbb{k}Q$ for any field $\mathbb{k}$; in other words, the representation theory of acyclic quivers is a special case of the representation theory of left regular bands!

4.2. Gradings, quadratic algebras and Koszul algebras. In this subsection we recall the notions of quadratic and Koszul algebras. Motivation for these notions can be found in [14, 57, 76]. We continue to hold fixed a field $\mathbb{k}$ and assume that all $\mathbb{k}$ -algebras are unital. A $\mathbb{k}$ -algebra A is said to be *graded* if it has a vector space decomposition $A = \bigoplus_{n \geq 0} A_n$ such that $A_i A_j \subseteq A_{i+j}$. An ideal I of A is *homogeneous* if it is generated as an ideal by the elements of $I \cap A_i$, for $i \geq 0$. In this case, $A/I = \bigoplus_{n \geq 0} (A_n + I)/I$ is a grading. Note that $I = \bigoplus_{n \geq 1} A_i$ is a homogeneous ideal and $A/I \cong A_0$ with the grading trivial (i.e., concentrated in degree 0). In particular, we can view A_0 as an A -module. A *graded A -module* is an A -module $M = \bigoplus_{n \geq 0} M_n$ such that $A_i M_j \subseteq M_{i+j}$ for all i, j . We say that M is generated in degree i if $M_j = 0$ for $j < i$ and $M_j = A_{j-i} M_i$ for $j \geq i$. An A -module homomorphism $\varphi: M \rightarrow N$ between graded A -modules has *degree d* if $\varphi(M_i) \subseteq N_{i+d}$ for all $i \geq 0$.

The path algebra $\mathbb{k}Q = \bigoplus_{n \geq 0} \mathbb{k}Q_n$ of a quiver is naturally graded by path length. That is, the homogeneous component of degree $n \geq 0$ is the $\mathbb{k}$ -span of the set Q_n of paths of length n . The arrow ideal

$$J = \bigoplus_{n \geq 1} \mathbb{k}Q_n$$

is graded and $\mathbb{k}^{|Q|} \cong \mathbb{k}Q_0 \cong \mathbb{k}Q/J$. If I is a homogeneous admissible ideal, then $A = \mathbb{k}Q/I$ is a graded finite dimensional $\mathbb{k}$ -algebra.

A graded $\mathbb{k}$ -algebra A is *quadratic* if A_0 is semisimple and A is generated by A_1 over A_0 with relations of degree 2. Rather than define this formally at this level of generality, we stick to the case of path algebras factored by homogeneous ideals. If I is a homogeneous admissible ideal of $\mathbb{k}Q$ that is generated as an ideal by elements in degree 2 (i.e., by elements which are linear combinations of paths of length 2), then $A = \mathbb{k}Q/I$ is called a *quadratic algebra*. We now wish to define, restricted to this setting, the quadratic dual. Our presentation follows that of [76] and [57]. Define a non-degenerate bilinear form

$$\langle \cdot, \cdot \rangle: \mathbb{k}Q_2 \times \mathbb{k}Q_2^{op} \rightarrow \mathbb{k}$$

by putting, for paths p, q in Q_2 and Q_2^{op} , respectively,

$$\langle p, q \rangle = \begin{cases} 1, & \text{if } q = p^\circ \\ 0, & \text{else} \end{cases}$$

where p° is the path in Q^{op} obtained from p by reversing all the edges and their order. Let $I_2 = I \cap \mathbb{k}Q_2$ and let $I_2^\perp \subseteq \mathbb{k}Q_2^{op}$ be the orthogonal complement of I_2 . Then $A^\dagger = \mathbb{k}Q^{op}/\langle I_2^\perp \rangle$ is called the *quadratic dual* of A .

Suppose that A is a graded $\mathbb{k}$ -algebra. Then a graded A -module M has a *linear resolution* if it is generated in degree 0 and there is a projective resolution $P_\bullet \rightarrow M$ such that each P_i is a graded A -module generated in degree i for all $i \geq 0$ and the maps in the resolution all have degree 0. One says that A is a *Koszul algebra* if A_0 is a semisimple $\mathbb{k}$ -algebra and A_0 , considered as a graded A -module concentrated in degree 0, has a linear resolution. In the case $A = \mathbb{k}Q/I$ for a homogeneous admissible ideal, one has that $A_0 \cong \mathbb{k}^{|Q_0|}$ is a semisimple algebra. Also observe that in this case A_0 admits a linear resolution if and only if each simple A -module (viewed as a graded module concentrated in degree 0) has a linear resolution. A $\mathbb{k}$ -algebra A admits at most one grading (up to isomorphism of graded algebras) making it Koszul.

Koszul algebras are always quadratic [14] but the converse is false. The quadratic dual of a Koszul algebra is also called its *Koszul dual*. If A is a graded $\mathbb{k}$ -algebra, the *Ext-algebra* of A is the graded $\mathbb{k}$ -algebra

$$\text{Ext}(A) = \bigoplus_{n \geq 0} \text{Ext}_A^n(A_0, A_0)$$

with multiplication given by the Yoneda composition. If A is a Koszul algebra, then $A^\dagger \cong \text{Ext}(A)^{op}$ and $\text{Ext}(\text{Ext}(A)) \cong A$. See [14] for details.

4.3. The algebra of a left regular band. Fix a (finite) left regular band B and a commutative ring with unit $\mathbb{k}$ for this subsection. The semigroup algebra $\mathbb{k}B$ is the free $\mathbb{k}$ -module with basis B and multiplication extending that of B . Formally, the product is given by

$$\sum_{b \in B} c_b b \cdot \sum_{b \in B} d_b b = \sum_{b, b' \in B} c_b d_{b'} b b'.$$

Note that $\mathbb{k}B$ is not in general unital. Indeed, if X is a finite set with at least two elements and we make X into a left regular band via the *left zero multiplication*, that is, by setting $xy = x$ for all $x \in X$, then $\mathbb{k}X$ is not unital. Indeed, for each $x \in X$, one has that

$$\sum_{y \in X} c_y y \cdot x = \sum_{y \in X} c_y y$$

and hence each element of X is a right identity for $\mathbb{k}X$. It turns out, though, that $\mathbb{k}B$ does have a right identity (not necessarily unique) for each left regular band. However, we shall see that $\mathbb{k}B$ can be unital even when B

is not a monoid. In particular, we shall characterize when $\mathbb{k}B$ is unital in terms of the poset structure on B .

We first recall some basic results on the structure of the semigroup algebra $\mathbb{k}B$. These results can be found in [96] for the case when $\mathbb{k}$ is a field and in [75] for general rings under the assumption that B is a monoid. For completeness we reprove the results here for arbitrary finite left regular bands (not necessarily monoids).

4.3.1. Orthogonal idempotents. Let us recall the construction of a set of orthogonal idempotents from [96]. They form a complete set of orthogonal idempotents when B is a left regular band monoid; in general, they just sum up to a right identity. When $\mathbb{k}$ is a field and $\mathbb{k}B$ is unital, they are primitive idempotents. Fix for each $X \in \Lambda(B)$ an element e_X with $X = Be_X$. Denoting the minimum element of $\Lambda(B)$ by $\widehat{0}$, define η_X recursively by $\eta_{\widehat{0}} = e_{\widehat{0}}$ and, for $X > \widehat{0}$,

$$\eta_X = e_X - \sum_{Y < X} e_X \eta_Y. \quad (4.4)$$

Notice that, by induction, one can write

$$\eta_X = \sum_{b \in B} c_b b$$

with the c_b integers such that $e_X \geq b$ for all b with $c_b \neq 0$ and the coefficient of e_X in η_X is 1. Also note that by construction

$$e_X = \sum_{Y \leq X} e_X \eta_Y.$$

The following results are proved in Lemma 4.1 and Theorem 4.2 of [96] when $\mathbb{k}$ is a field and B is a monoid.

Theorem 4.2. *The elements $\{\eta_X\}_{X \in \Lambda(B)}$ enjoy the following properties.*

- (1) *If $a \in B$ and $X \in \Lambda(B)$ are such that $a \notin B_{\geq X}$, then $a\eta_X = 0$.*
- (2) *One has that $\{\eta_X \mid X \in \Lambda(B)\}$ is a set of orthogonal idempotents and that*

$$\eta = \sum_{X \in \Lambda(B)} \eta_X$$

is a right identity for $\mathbb{k}B$. In particular, if $\mathbb{k}B$ is unital (for instance, if B is a monoid), then $\eta = 1$ and $\{\eta_X\}_{X \in \Lambda(B)}$ is a complete set of orthogonal idempotents.

- (3) *If $\mathbb{k}$ is a field and $\mathbb{k}B$ is unital, then each η_X , with $X \in \Lambda(B)$, is a primitive idempotent.*

Proof. We prove the first item by induction on X . If $X = \widehat{0}$, then there is nothing to prove. So assume that it is true for $Y < X$ and we prove it for

X . Let $a \in B \setminus B_{\geq X}$ and put $Z = X \wedge Ba = Be_X \cap Ba = Bae_X$. Then we have that

$$\begin{aligned} a\eta_X &= ae_X - \sum_{Y < X} ae_X \eta_Y \\ &= ae_X - \sum_{Y \leq Z} ae_X \eta_Y - \sum_{Y < X, Y \not\leq Ba} ae_X \eta_Y \\ &= ae_X - ae_X \sum_{Y \leq Z} \eta_Y \end{aligned}$$

where the last equality uses the induction hypothesis. As $Bae_X = Z = Be_Z$, we conclude that $ae_X = ae_X e_Z$ and so

$$a\eta_X = ae_X - ae_X \sum_{Y \leq Z} \eta_Y = ae_X e_Z - ae_X e_Z \sum_{Y \leq Z} \eta_Y = ae_X e_Z - ae_X e_Z = 0$$

because

$$e_Z \sum_{Y \leq Z} \eta_Y = e_Z.$$

This completes the proof of the first item.

For the second item, we first verify that η_X is idempotent. Indeed,

$$\eta_X^2 = e_X \eta_X - \sum_{Y < X} e_X \eta_Y \eta_X = e_X \eta_X = \eta_X$$

by the first item because each element $a \in B$ with non-zero coefficient in $e_X \eta_Y$ for $Y < X$ satisfies $Ba \not\leq X$.

Next we check that $\eta_X \eta_Z = 0$ if $X \neq Z$. First suppose that $Z \not\leq X$. As every element of B with a non-zero coefficient in η_X belongs to $e_X B \subseteq Be_X$, we conclude that $\eta_X \eta_Z = 0$ by the first item. Thus we are reduced to the case that $Z < X$. We proceed by induction. Assume that $\eta_Y \eta_Z = 0$ for all Y with $Z < Y < X$. Then we have that

$$\eta_X \eta_Z = e_X \eta_Z - \sum_{Y < X} e_X \eta_Y \eta_Z = e_X \eta_Z - e_X \eta_Z = 0$$

because η_Z is idempotent, $\eta_Y \eta_Z = 0$ if $Z \not\leq Y$ by the first case and $\eta_Y \eta_Z = 0$ if $Z < Y < X$ by induction. This completes the proof that $\{\eta_X \mid X \in \Lambda(B)\}$ is an orthogonal set of idempotents. It remains to verify that

$$\eta = \sum_{X \in \Lambda(B)} \eta_X$$

is a right identity for $\mathbb{k}B$. Indeed, by the first item, if $b \in B$ and $X = Bb$, then $be_X = b$ and

$$b\eta = \sum_{Y \leq X} b\eta_Y = be_X \sum_{Y \leq X} \eta_Y = be_X = b$$

as required.

The final item will be proved in Theorem 4.6 below. $\square$

Corollary 4.3. *If B is a left regular band, then $\mathbb{k}B$ is a projective $\mathbb{k}B$ -module as are its direct summands.*

Proof. We have $B = \mathbb{k}B\eta$ and so $\text{Hom}_{\mathbb{k}B}(\mathbb{k}B, M) \cong \eta M$ for any left $\mathbb{k}B$ -module M . As the functor $M \mapsto \eta M$ is obviously exact, this completes the proof. $\square$

The following is [96, Corollary 4.4] for fields and is in [75] for the general case, but with B a monoid.

Corollary 4.4. *The set $\{b\eta_{\sigma(b)} \mid b \in B\}$ is a basis of idempotents for $\mathbb{k}B$.*

Proof. Clearly, $b\eta_{\sigma(b)}b\eta_{\sigma(b)} = b\eta_{\sigma(b)}^2 = b\eta_{\sigma(b)}$ by the left regular band axiom and idempotence of $\eta_{\sigma(b)}$. As $b\eta_{\sigma(b)} = b$, it follows that

$$b\eta_{\sigma(b)} = b + \sum_{a \in \partial b B} c_a a$$

and so the mapping $\mathbb{k}B \rightarrow \mathbb{k}B$ induced by $b \mapsto b\eta_{\sigma(b)}$ has matrix which is unipotent upper triangular with respect to any linear extension of the natural partial order on B . The result follows. $\square$

4.3.2. Schützenberger representations. We recall the classical (left) *Schützenberger representation* associated to an element $X \in \Lambda(B)$. We retain the notation of the previous subsection. Let $L_X = \sigma^{-1}(X)$. It is the $\mathcal{L}$ -class of e_X in the sense of semigroup theory [46], that is, it consists of all elements that generate the same principal left ideal of B as e_X . Define a $\mathbb{k}B$ -module structure on $\mathbb{k}L_X$ by putting, for $a \in B$ and $b \in L_X$,

$$a \cdot b = \begin{cases} ab & \text{if } \sigma(a) \geq X \\ 0 & \text{else.} \end{cases}$$

Let us compute the endomorphism ring of $\mathbb{k}L_X$.

Proposition 4.5. *Let $\mathbb{k}$ be a commutative ring with unit and B a left regular band. Let $X \in \Lambda(B)$. Then $\text{End}_{\mathbb{k}B}(\mathbb{k}L_X) \cong \mathbb{k}$. Consequently, if $\mathbb{k}$ is a field, then $\mathbb{k}L_X$ is indecomposable.*

Proof. Fix e_X with $Be_X = X$, as usual. Let $\lambda: \mathbb{k}L_X \rightarrow \mathbb{k}$ be the $\mathbb{k}$ -module homomorphism defined by $\lambda(a) = 1$ for $a \in L_X$ and define

$$\Psi: \text{End}_{\mathbb{k}B}(\mathbb{k}L_X) \rightarrow \mathbb{k}$$

by $\Psi(\varphi) = \lambda(\varphi(e_X))$. We claim that $\varphi(a) = \Psi(\varphi)a$ for all $a \in L_X$. The proposition will follow from this and the observation that $\Psi(c1_{\mathbb{k}L_X}) = c$ for $c \in \mathbb{k}$ because $\mathbb{k}L_X$ is a free $\mathbb{k}$ -module. Indeed, suppose that $\varphi(e_X) = \sum_{b \in L_X} c_b b$. If $a \in L_X$, then we have that $ab = a$ for all $b \in L_X$. Therefore, we have that

$$\varphi(a) = \varphi(ae_X) = a\varphi(e_X) = \sum_{b \in L_X} c_b ab = \left(\sum_{b \in L_X} c_b \right) a = \Psi(\varphi)a.$$

This establishes that $\text{End}_{\mathbb{k}B}(\mathbb{k}L_X) \cong \mathbb{k}$.

The second statement follows from the first because a field has no non-trivial idempotents. $\square$

The next theorem was proved in [96] for $\mathbb{k}$ a field and over a general base commutative ring in [75] under the assumption that B has an identity.

Theorem 4.6. *Let B be a left regular band and $\mathbb{k}$ a commutative ring with unit. Then the Schützenberger representation $\mathbb{k}L_X$ is isomorphic to $\mathbb{k}B\eta_X$, for $X \in \Lambda(B)$ via the mapping $b \mapsto b\eta_X$ and, consequently, is a projective module. Moreover,*

$$\mathbb{k}B \cong \bigoplus_{X \in \Lambda(B)} \mathbb{k}L_X. \quad (4.5)$$

If $\mathbb{k}$ is a field and $\mathbb{k}B$ is unital, then this is the decomposition of $\mathbb{k}B$ into projective indecomposables and hence each η_X is a primitive idempotent.

Proof. Fix $X \in \Lambda(B)$. Define a $\mathbb{k}$ -linear map $\varphi: \mathbb{k}L_X \rightarrow \mathbb{k}B\eta_X$ by $\varphi(b) = b\eta_X$ for $b \in L_X$. The mapping φ is injective as a consequence of Corollary 4.4. It is surjective because if $a \in B$, then $a\eta_X = 0$ unless $Ba \supseteq X$ by Theorem 4.2. But if $X = Be_X$ and $Ba \supseteq X$, then $a\eta_X = ae_X\eta_X$ and $ae_X \in L_X$. Thus $a\eta_X = \varphi(ae_X)$ and so φ is surjective. We check that φ is $\mathbb{k}B$ -linear. If $a \in L_X$ and $b \in B$ with $Bb \supseteq X$, then

$$\varphi(b \cdot a) = ba\eta_X = b(a\eta_X) = b\varphi(a).$$

If $Bb \not\supseteq X$, then $ba\eta_X = 0$ by Theorem 4.2 and so

$$\varphi(b \cdot a) = \varphi(0) = 0 = b(a\eta_X) = b\varphi(a).$$

This establishes that $\mathbb{k}B\eta_X \cong \mathbb{k}L_X$.

To prove (4.5), we recall that

$$\eta = \sum_{X \in \Lambda(B)} \eta_X$$

is a right identity and the η_X with $X \in \Lambda(B)$ form a set of orthogonal idempotents. Thus (4.5) follows from

$$\mathbb{k}B = \mathbb{k}B\eta = \bigoplus_{X \in \Lambda(B)} \mathbb{k}B\eta_X$$

and the first part of the theorem. The final statement is a consequence of Proposition 4.5 because $\mathbb{k}B\eta_X \cong \mathbb{k}L_X$ is indecomposable. $\square$

If $X \in \Lambda(B)$, then there is a $\mathbb{k}$ -algebra homomorphism $\rho_X: \mathbb{k}B \rightarrow \mathbb{k}B_{\geq X}$ given by

$$\rho_X(b) = \begin{cases} b & \text{if } \sigma(b) \geq X \\ 0 & \text{else} \end{cases}$$

for $b \in B$. This homomorphism allows us to consider any $\mathbb{k}B_{\geq X}$ -module M as a $\mathbb{k}B$ -module via the action $b \cdot m = \rho_X(b)m$ for all $b \in B$ and $m \in M$.

Denote by $\mathbb{k}_X$ the trivial $\mathbb{k}B_{\geq X}$ -module, viewed as a $\mathbb{k}B$ -module. It is known that if $\mathbb{k}$ is a field, then the $\mathbb{k}_X$ with $X \in \Lambda(B)$ form a complete set of non-isomorphic simple $\mathbb{k}B$ -modules and the augmentation map $\lambda_X: \mathbb{k}L_X \rightarrow \mathbb{k}_X$ is the projective cover [96]; in fact, this is an immediate consequence of Theorem 4.6 because the $\mathbb{k}L_X$ are a complete set of projective indecomposables modules and hence have simple tops. Let us state this as a corollary.

Corollary 4.7. *Let $\mathbb{k}$ be a field and B a left regular band such that $\mathbb{k}B$ is unital. Then the simple $\mathbb{k}B$ -modules $\mathbb{k}_X$ with $X \in \Lambda(B)$ form a complete set of representatives of the isomorphism classes of simple $\mathbb{k}B$ -modules and the mapping $\varphi_X: \mathbb{k}L_X \rightarrow \mathbb{k}_X$ given by $\varphi_X(b) = 1$ for $b \in L_X$ is the projective cover.*

If $Y \geq X$, then $\mathbb{k}L_Y$ is a $\mathbb{k}B_{\geq X}$ -module and is the projective indecomposable module for $\mathbb{k}B_{\geq X}$ corresponding to $Y \in \Lambda(B)_{\geq X} \cong \Lambda(B_{\geq X})$. We, therefore, obtain the following corollary of Theorem 4.6.

Corollary 4.8. *If $X \in \Lambda(B)$, then $\mathbb{k}B_{\geq X}$ is a projective $\mathbb{k}B$ -module and the decomposition*

$$\mathbb{k}B \cong \mathbb{k}B_{\geq X} \oplus \bigoplus_{Y \not\geq X} \mathbb{k}L_Y$$

holds. Consequently, any projective $\mathbb{k}B_{\geq X}$ -module is a projective $\mathbb{k}B$ -module (via ρ_X).

As a corollary, it follows that we can compute projective resolutions for $\mathbb{k}B_{\geq X}$ -modules, and compute Ext between them, over either $\mathbb{k}B$ or $\mathbb{k}B_{\geq X}$.

Corollary 4.9. *Let $X \in \Lambda(B)$ and let M, N be $\mathbb{k}B_{\geq X}$ -modules. Any projective resolution $P_\bullet \rightarrow M$ of M as a $\mathbb{k}B_{\geq X}$ -module is also a projective resolution as a $\mathbb{k}B$ -module. Consequently,*

$$\mathrm{Ext}_{\mathbb{k}B}^n(M, N) \cong \mathrm{Ext}_{\mathbb{k}B_{\geq X}}^n(M, N)$$

for all $n \geq 0$.

4.3.3. The Jacobson radical. In this section we compute the Jacobson radical of the algebra of a left regular band over a semiprimitive commutative ring with unit $\mathbb{k}$. Munn characterized more generally the radical of a band over any base ring [80].

First we need Solomon's theorem describing the algebra of a finite semilattice [104]. We include a proof for completeness. The *incidence algebra* $I(P; \mathbb{k})$ of a finite poset P is the $\mathbb{k}$ -algebra with $\mathbb{k}$ -basis consisting all pairs (σ, τ) with $\sigma \leq \tau$ and whose product on basis vectors is given by

$$(\sigma, \tau)(\alpha, \beta) = \begin{cases} (\alpha, \tau), & \text{if } \beta = \sigma \\ 0, & \text{else.} \end{cases}$$

If $\mathbb{k}$ is field, then $I(P; \mathbb{k}) \cong \mathbb{k}Q/\langle R \rangle$ where Q is the Hasse diagram of P and R is the set of all differences $p - q$ of parallel paths in Q [6].

We can identify elements of $I(P; \mathbb{k})$ with mappings $f: P \times P \rightarrow \mathbb{k}$ with $f(p, q) = 0$ whenever $p \not\leq q$. The mapping f corresponds to the element

$$\sum_{p \leq q} f(p, q) \cdot (p, q)$$

of $I(P; \mathbb{k})$. The identity element $\sum_{p \in P} (p, p)$ of $I(P; \mathbb{k})$ then corresponds to the Kronecker function δ . The product is given by

$$(f \cdot g)(p, q) = \sum_{p \leq x \leq q} f(x, q)g(p, x)$$

for $p \leq q$. An element $f \in I(P; \mathbb{k})$ is invertible if and only if $f(p, p)$ is a unit of $\mathbb{k}$ for all $p \in P$. The inverse of f is given recursively by

$$\begin{aligned} f^{-1}(p, p) &= f(p, p)^{-1} \\ f^{-1}(p, q) &= -f(p, p)^{-1} \sum_{p \leq r < q} f(r, q) f^{-1}(p, r), \quad \text{for } p < q. \end{aligned} \quad (4.6)$$

The *zeta function* of P is the element $\zeta \in I(P; \mathbb{k})$ given by

$$\zeta(p, q) = \begin{cases} 1, & \text{if } p \leq q \\ 0, & \text{else.} \end{cases}$$

It is invertible over any base ring $\mathbb{k}$ and has inverse the *Möbius function* μ of P . Normally, one considers the Möbius function to be the inverse of ζ in $I(P; \mathbb{Z})$ as the general case is then a specialization.

Theorem 4.10. *Let $\mathbb{k}$ be a commutative ring with unit and Λ a finite (meet) semilattice. Then $\mathbb{k}\Lambda \cong \mathbb{k}^\Lambda$ and, in particular, is unital.*

Moreover, if

$$\varepsilon_x = \sum_{y \leq x} \mu(y, x)y$$

for $x \in \Lambda$, then the ε_x form a complete set of orthogonal idempotents and $\mathbb{k}\Lambda \varepsilon_x = \mathbb{k} \varepsilon_x$. Furthermore, if $y \in \Lambda$ and $c \in \mathbb{k}$, then

$$y c \varepsilon_x = \begin{cases} c \varepsilon_x, & \text{if } y \geq x \\ 0, & \text{else} \end{cases}$$

holds. In particular, if $\mathbb{k}$ is a field, then the ε_x form a complete set of orthogonal primitive idempotents.

Proof. For $y \in \Lambda$, put

$$\delta_y(z) = \begin{cases} 1, & \text{if } z = y \\ 0, & \text{else.} \end{cases}$$

and note that the δ_y with $y \in \Lambda$ form a $\mathbb{k}$ -basis for $\mathbb{k}^\Lambda$.

Define $\varphi: \mathbb{k}\Lambda \longrightarrow \mathbb{k}^\Lambda$ and $\psi: \mathbb{k}^\Lambda \longrightarrow \mathbb{k}\Lambda$ by

$$\begin{aligned}\varphi(x) &= \sum_{z \leq x} \delta_z \\ \psi(f) &= \sum_{x \in \Lambda} \sum_{z \leq x} f(x) \mu(z, x) z\end{aligned}$$

for $x \in \Lambda$. In particular,

$$\psi(\delta_x) = \sum_{z \leq x} \mu(z, x) z.$$

First we verify that φ is a homomorphism. It suffices to check that $\varphi(x)\varphi(y) = \varphi(xy)$ for $x, y \in \Lambda$. Indeed, we have

$$\varphi(x)\varphi(y) = \sum_{z \leq x} \delta_z \cdot \sum_{u \leq y} \delta_u = \sum_{z \leq x, y} \delta_z = \sum_{z \leq xy} \delta_z = \varphi(xy).$$

Next we establish that φ and ψ are inverse mappings. For $x \in \Lambda$, we have

$$\begin{aligned}\psi(\varphi(x)) &= \sum_{y \leq x} \sum_{z \leq y} \mu(z, y) z = \sum_{z \leq x} \left(\sum_{z \leq y \leq x} \zeta(y, x) \mu(z, y) \right) z = x \\ \varphi(\psi(\delta_x)) &= \sum_{y \leq x} \sum_{z \leq y} \mu(y, x) \delta_z = \sum_{z \leq x} \left(\sum_{z \leq y \leq x} \mu(y, x) \zeta(z, y) \right) \delta_z = \delta_x\end{aligned}$$

as required. This completes the proof that φ is an isomorphism of $\mathbb{k}$ -algebras with inverse ψ .

As $\varepsilon_x = \psi(\delta_x)$, it remains to consider the action of $y \in \Lambda$ on $\mathbb{k}\Lambda\varepsilon_x$ or, equivalently, the action of $\varphi(y)$ on $\mathbb{k}^\Lambda\delta_x = \mathbb{k}\delta_x$. But we have

$$\varphi(y)\delta_x = \sum_{z \leq y} \delta_z \delta_x = \begin{cases} \delta_x, & \text{if } y \geq x \\ 0, & \text{else.} \end{cases}$$

This completes the proof. $\square$

Recall that if B is a left regular band, then $\Lambda(B) = \{Bb \mid b \in B\}$ is a meet semilattice with $Ba \cap Bb = Bab = Bba$ and we have the support homomorphism $\sigma: B \longrightarrow \Lambda(B)$ given by $\sigma(b) = Bb$. As before, if $X \in \Lambda(B)$, then $L_X = \sigma^{-1}(X)$ is an $\mathcal{L}$ -class of B . Abusing notation, we shall also write σ for the unique extension $\sigma: \mathbb{k}B \longrightarrow \mathbb{k}\Lambda(B)$. The following result is a special case of the results of [80]; see also [34].

Theorem 4.11. *Let B be a finite left regular band and $\mathbb{k}$ a commutative ring with unit. Let J be the kernel of the homomorphism $\sigma: \mathbb{k}B \longrightarrow \mathbb{k}\Lambda(B)$ induced by $b \mapsto Bb$. Let n be the largest integer such that there is a chain $X_1 > X_2 > \cdots > X_n$ in $\Lambda(B)$. Then $J^{n+1} = 0$ and hence J is nilpotent.*

Proof. If $r = \sum_{b \in B} c_b b$ with the $c_b \in \mathbb{k}$ and $X \in \Lambda(B)$, then put $r_X = \sum_{b \in L_X} c_b b$. Notice that $r = \sum_{X \in \Lambda(B)} r_X$ and that

$$\sigma(r) = \sum_{b \in B} c_b \sigma(b) = \sum_{X \in \Lambda(B)} \left(\sum_{b \in L_X} c_b \right) X.$$

Therefore, $r \in J$ if and only if, for each $X \in \Lambda(B)$, one has that $\sum_{b \in L_X} c_b = 0$. It follows from this that $r \in J$ if and only if $r_X \in J$ for all $X \in \Lambda(B)$.

Let us put $J_X = \{r \in J \mid r = r_X\}$. Then we have that

$$J = \bigoplus_{X \in \Lambda(B)} J_X$$

from the discussion in the previous paragraph. Thus to prove our theorem it suffices to prove that if $Y_1, \dots, Y_{n+1} \in \Lambda(B)$, then $J_{Y_1} \cdots J_{Y_{n+1}} = 0$. Put $X_i = Y_1 \wedge Y_2 \wedge \cdots \wedge Y_i$ for $1 \leq i \leq n+1$.

We claim that if $Y \leq Z$, then $J_Y J_Z = 0$. Indeed, if $r = \sum_{a \in L_Y} c_a a$ and $b \in B$ with $\sigma(b) = Z \geq Y$, then

$$rb = \sum_{a \in L_Y} c_a ab = \sum_{a \in L_Y} c_a a = r$$

because $ab = a$ if $Ba \subseteq Bb$. Therefore, if $r \in J_Y$ and $s = \sum_{b \in L_Z} d_b b \in J_Z$, then

$$rs = \sum_{b \in L_Z} d_b rb = \left(\sum_{b \in L_Z} d_b \right) r = 0$$

as $s \in J_Z$ implies $\sum_{b \in L_Z} d_b = 0$. We conclude that $J_Y J_Z = 0$ whenever $Y \leq Z$.

Next observe that $J_{Y_1} \cdots J_{Y_i} \subseteq J_{X_i}$ for $1 \leq i \leq n+1$ because if $\sigma(y_k) = Y_k$ for $1 \leq k \leq i$, then $\sigma(y_1 \cdots y_i) = \sigma(y_1) \cdots \sigma(y_i) = X_i$. As

$$X_1 \geq X_2 \geq \cdots \geq X_{n+1}$$

we deduce by choice of n that $X_i = X_{i+1}$ for some $1 \leq i < n+1$. But then $X_i = X_{i+1} = X_i \wedge Y_{i+1} \leq Y_{i+1}$. Therefore, we have

$$J_{Y_1} \cdots J_{Y_i} J_{Y_{i+1}} \subseteq J_{X_i} J_{Y_{i+1}} = 0$$

where the last equality follows from the claim because $X_i \leq Y_{i+1}$. We conclude that $J_{Y_1} \cdots J_{Y_{n+1}} = 0$, completing the proof. $\square$

As a consequence, we can deduce that J is the Jacobson radical of $\mathbb{k}B$ if the ground ring $\mathbb{k}$ is sufficiently nice.

Corollary 4.12. *Let B be a finite left regular band and $\mathbb{k}$ a semiprimitive commutative ring with unit (e.g. a field). Then the Jacobson radical of $\mathbb{k}B$ is the kernel of the homomorphism $\sigma: \mathbb{k}B \rightarrow \mathbb{k}\Lambda(B)$ induced by $b \mapsto Bb$ for $b \in B$.*

Proof. A nilpotent ideal is contained in the Jacobson radical of any ring. Thus $\ker \sigma \subseteq \text{rad}(\mathbb{k}B)$ by Theorem 4.11. The isomorphism $\mathbb{k}\Lambda(B) \cong \mathbb{k}^\Lambda$ of Theorem 4.10 implies that $\mathbb{k}\Lambda(B)$ is semiprimitive. We deduce that $\text{rad}(\mathbb{k}B) \subseteq \ker \sigma$ and the theorem follows. $\square$

It follows from Corollary 4.12 that $\mathbb{k}B$ is a split basic algebra over any field $\mathbb{k}$ whenever it is unital.

Note that as a $\mathbb{k}B$ -module we have that

$$\mathbb{k}\Lambda(B) \cong \bigoplus_{X \in \Lambda(B)} \mathbb{k}_X \quad (4.7)$$

by Theorem 4.10. This provides another proof of Corollary 4.7, which states that when $\mathbb{k}$ is a field and $\mathbb{k}B$ is unital, then the $\mathbb{k}_X$ with $X \in \Lambda(B)$ are the simple $\mathbb{k}B$ -modules.

It will be convenient to have a basis for the radical of a left regular band algebra.

Proposition 4.13. *Let B be a left regular band and $\mathbb{k}$ a commutative ring with unit. Let J be the kernel of the homomorphism $\sigma: \mathbb{k}B \rightarrow \mathbb{k}\Lambda(B)$ induced by $\sigma(b) = Bb$ for $b \in B$. Fix, for each $X \in \Lambda(B)$, an element $e_X \in B$ with $Be_X = X$. Then the non-zero elements of the form $b - e_{Bb}$ form a basis for J over $\mathbb{k}$.*

Proof. The linear independence of the non-zero elements of the form $b - e_{Bb}$ is immediate from the linear independence of B . Trivially, $\sigma(b - e_{Bb}) = Bb - Bb = 0$. So it remains to show that these elements span J as a $\mathbb{k}$ -module. So suppose that

$$x = \sum_{b \in B} c_b b = \sum_{X \in \Lambda(B)} \sum_{Bb=X} c_b b$$

belongs to J . Then we have

$$0 = \sum_{X \in \Lambda(B)} \sum_{Bb=X} c_b X$$

and hence for each $X \in \Lambda(B)$, one has that

$$\sum_{Bb=X} c_b = 0.$$

It follows that

$$x = \sum_{X \in \Lambda(B)} \sum_{Bb=X} c_b (b - e_{Bb})$$

as required. $\square$

A consequence of Proposition 4.13 that we shall use in the next section is the following.

Corollary 4.14. *Let B be a left regular band and $\mathbb{k}$ a field. Let J be the kernel of the natural homomorphism $\sigma: \mathbb{k}B \rightarrow \mathbb{k}\Lambda(B)$ induced by $b \mapsto Bb$ for $b \in B$. Then J is the Jacobson radical of $\mathbb{k}B^1$.*

Proof. Let R be the kernel of the natural homomorphism $\rho: \mathbb{k}B^1 \longrightarrow \mathbb{k}\Lambda(B^1)$ induced by $b \mapsto B^1b$ for $b \in B^1$; it's the radical of $\mathbb{k}B^1$ by Corollary 4.12. As 1 is the unique generator of the left ideal B^1 and $Ba = Bb$ if and only if $B^1a = B^1b$ for $a, b \in B$, it follows from Proposition 4.13 that J and R have the same basis and hence coincide. $\square$

4.4. Existence of identity elements in left regular band algebras.

Let B be a finite left regular band. It will be important to characterize when $\mathbb{k}B$ is a unital ring. The answer turns out to be independent of the ground ring $\mathbb{k}$ and depends, instead, on the topology of B and its contractions as posets. Following our earlier convention, we say that B is *connected* if $\Delta(B_{\geq X})$ (or, equivalently, the Hasse diagram of $B_{\geq X}$) is connected for each $X \in \Lambda(B)$. Of course, monoids are connected since all contractions of a monoid are monoids and hence have contractible order complexes. It turns out that $\mathbb{k}B$ is unital if and only if B is connected.

The proof is surprisingly indirect in that some representation theory is used. It would be nice to have a completely elementary proof of the existence of an identity. Later we shall see that if B is a connected left regular band, then, in fact, $\Delta(B_{\geq X})$ is an acyclic topological space for all $X \in \Lambda(B)$.

Theorem 4.15. *The following conditions are equivalent for a finite left regular band B .*

- (1) B is connected.
- (2) $\mathbb{Z}B$ is unital.
- (3) $\mathbb{k}B$ is unital for each commutative ring with unit $\mathbb{k}$.

Proof. Clearly the second item implies the third as $\mathbb{k}B \cong \mathbb{k} \otimes_{\mathbb{Z}} \mathbb{Z}B$ has identity $1 \otimes 1$. Suppose that $\Delta(B_{\geq X})$ is not connected. Denote by $\pi_0(Y)$ the set of connected components of a topological space Y . There is a surjective homomorphism $B_{\geq X} \longrightarrow \pi_0(\Delta(B_{\geq X}))$ where $\pi_0(\Delta(B_{\geq X}))$ is given the left zero multiplication (in fact, this is the maximal left zero image of $B_{\geq X}$; see [73]). Then we have surjective homomorphisms

$$\mathbb{k}B \longrightarrow \mathbb{k}B_{\geq X} \longrightarrow \mathbb{k}\pi_0(\Delta(B_{\geq X}))$$

and hence $\mathbb{k}B$ is not unital (as $\mathbb{k}\pi_0(\Delta(B_{\geq X}))$ is not by the discussion at the beginning of Section 4.3). This proves that the third item implies the first.

Suppose now that B is connected. If $\mathbb{Z}B$ is unital, then its identity must be $\eta = \sum_{X \in \Lambda(B)} \eta_X$, where the η_X are as defined as per (4.4), since η is a right identity by Theorem 4.2. It clearly suffices to show that η is, in fact, a left identity for $\mathbb{Q}B$.

Consider $\mathbb{Q}B^1$ and let $\hat{1} = B^1 \in \Lambda(B^1)$. Then $\eta_{\hat{1}} = 1 - \eta$, together with the η_X with $X \in \Lambda(B)$, form a complete set of orthogonal primitive idempotents for $\mathbb{Q}B^1$ by Theorem 4.2. Note that $\mathbb{Q}B$ is an ideal in the unital $\mathbb{Q}$ -algebra $\mathbb{Q}B^1$. Our goal is to show that $(1 - \eta)\mathbb{Q}B = 0$, as this implies that $x = \eta x$ for all $x \in \mathbb{Q}B$, as required. Let J be the kernel of the surjective homomorphism $\sigma: \mathbb{Q}B \longrightarrow \mathbb{Q}\Lambda(B)$ induced by $\sigma(b) = Bb$ for $b \in B$. The

algebra $\mathbb{Q}\Lambda(B)$ is unital by Theorem 4.10 and hence $\sigma(\eta)$ is the identity of $\mathbb{Q}\Lambda(B)$ because surjective homomorphism map right identities to right identities. Therefore, $\sigma(x - \eta x) = 0$ for all $x \in \mathbb{Q}B$ and so $(1 - \eta)\mathbb{Q}B \subseteq J$, whence $(1 - \eta)\mathbb{Q}B = (1 - \eta)J$. So it suffices to prove that $(1 - \eta)J = 0$.

As the ideal J is nilpotent by Theorem 4.11, it suffices to prove that $(1 - \eta)J = (1 - \eta)J^2$ or, equivalently, $(1 - \eta)J \subseteq J^2$. Thus, to complete the proof of the theorem, we are reduced to proving that $(1 - \eta)[J/J^2] = 0$. Recall that J is the Jacobson radical of $\mathbb{Q}B^1$ by Corollary 4.14 and $1 - \eta$ is the primitive idempotent corresponding to the simple module $\mathbb{Q}_{\hat{1}}$, which is annihilated by B and fixed by 1. Therefore, we have

$$\begin{aligned} (1 - \eta)[J/J^2] &\cong \text{Hom}_{\mathbb{Q}B^1/J}([\mathbb{Q}B^1/J](1 - \eta), J/J^2) \\ &= \text{Hom}_{\mathbb{Q}B^1/J}(\mathbb{Q}_{\hat{1}}, J/J^2) \\ &\cong \text{Hom}_{\mathbb{Q}B^1/J}(J/J^2, \mathbb{Q}_{\hat{1}}) \\ &\cong \text{Hom}_{\mathbb{Q}B^1}(J, \mathbb{Q}_{\hat{1}}) \end{aligned}$$

where the penultimate isomorphism uses that $\mathbb{Q}B^1/J$ is a split semisimple $\mathbb{Q}$ -algebra and the last isomorphism uses that $J^2 = \text{rad}(J)$ and that $\mathbb{Q}_{\hat{1}}$ is simple. In summary, to prove that η is an identity for $\mathbb{Q}B$, it remains to show that there are no non-zero $\mathbb{Q}B^1$ -module homomorphisms from J to $\mathbb{Q}_{\hat{1}}$.

So let $\varphi: J \rightarrow \mathbb{Q}_{\hat{1}}$ be a $\mathbb{Q}B^1$ -module homomorphism and fix, for each $X \in \Lambda(B)$, an element $e_X \in B$ with $X = Be_X$. By Proposition 4.13, J has $\mathbb{Q}$ -vector space basis consisting of elements of the form $b - e_X$ where $Bb = X$ and $b \neq e_X$. Thus to prove that φ is the zero mapping, it suffices to show that if $x, y \in B$ with $Bx = By = X$, then $\varphi(x - y) = 0$.

Let $\Gamma(X)$ be the connected graph from Lemma 3.18. Assume first that $x, y \in L_X$ are adjacent vertices. Then there exists $b \in B$ with $bx = x$ and $by = y$. Therefore, we have that

$$\varphi(x - y) = \varphi(b(x - y)) = b\varphi(x - y) = 0.$$

If x, y are not adjacent, then there is a sequence $x = x_0, x_1, \dots, x_n = y$ with x_i, x_{i+1} adjacent vertices of $\Gamma(X)$ for $i = 0, \dots, n-1$. But then we have the telescoping sum

$$x - y = \sum_{i=0}^{n-1} (x_i - x_{i+1})$$

and so

$$\varphi(x - y) = \sum_{i=0}^{n-1} \varphi(x_i - x_{i+1}) = 0$$

where the last equality holds by the previous case. We conclude that $\varphi = 0$, as required. This concludes the proof that $\mathbb{Z}B$ has an identity. $\square$

Most of our examples of left regular bands that are not monoids are strong elimination systems and hence are connected left regular bands.

Corollary 4.16. *Let $(E, \mathcal{L})$ be a strong elimination system and $\mathbb{k}$ a field. Then $\mathbb{k}\mathcal{L}$ is a unital $\mathbb{k}$ -algebra.*

Proof. This follows from Theorem 4.15 because $\mathcal{L}$ is a connected left regular band by Proposition 3.19. $\square$

4.5. Cartan invariants. Let A be a unital finite dimensional $\mathbb{k}$ -algebra over a field $\mathbb{k}$. Let $P_1, \dots, P_r$ be a complete set of representatives of the isomorphism classes of projective indecomposable A -modules and $S_1, \dots, S_r$ be a complete set of representatives of the isomorphism classes of simple A -modules. The *Cartan matrix* of A is the $r \times r$ -matrix C with C_{ij} the number of composition factors of P_j isomorphic to S_i . If A is a split $\mathbb{k}$ -algebra, then

$$C_{ij} = \dim_{\mathbb{k}} \operatorname{Hom}_A(P_i, P_j) = \dim_{\mathbb{k}} e_i A e_j$$

where e_i, e_j are primitive idempotents with $P_i \cong Ae_i$ and $P_j \cong Ae_j$. Thus one also has in this case that C_{ij} is the number of composition factors of the injective indecomposable module I_i isomorphic to S_j .

Saliola computed the Cartan matrix of the algebra of a left regular band monoid in terms of the Möbius function of its support lattice. We provide here the straightforward generalization for the algebra of a connected left regular band. If B is a connected left regular band, then both the projective indecomposable $\mathbb{k}B$ -modules and the simple $\mathbb{k}B$ -modules are indexed by $\Lambda(B)$ and so it is natural to index the Cartan matrix by this set. More precisely, if $X, Y \in \Lambda(B)$, then $C_{X,Y} = \dim_{\mathbb{k}} \operatorname{Hom}(\mathbb{k}L_X, \mathbb{k}L_Y)$.

Theorem 4.17. *Let B be a connected left regular band with support semilattice $\Lambda(B)$ and let $\mathbb{k}$ be a field. Fix $e_X \in B$ with $Be_X = X$ for all $X \in \Lambda(B)$. Then the Cartan matrix of $\mathbb{k}B$ is given by*

$$C_{X,Y} = \sum_{Z \leq X} |e_Z B \cap L_Y| \cdot \mu(Z, X) \quad (4.8)$$

for $X, Y \in \Lambda(B)$ where μ is the Möbius function of $\Lambda(B)$. In particular, $C_{X,Y}$ is unipotent lower triangular with respect to any linear extension of the partial order on $\Lambda(B)$.

Proof. Let $\rho: B \rightarrow M_n(\mathbb{k})$ be the matrix representation afforded by $\mathbb{k}L_Y$ with respect to the basis L_Y and let χ be the character of ρ , that is, $\chi(b)$ is the trace of $\rho(b)$. Note that $\chi(b) = |bB \cap L_Y|$ because b is idempotent. Recall that if $Z \in \Lambda(B)$, then the irreducible representation χ_Z corresponding to the simple module $\mathbb{k}_Z$ is given by

$$\chi_Z(a) = \begin{cases} 1, & \text{if } Ba \geq Z \\ 0, & \text{else} \end{cases}$$

for $a \in B$.

Choosing a basis for $\mathbb{k}L_Y$ adapted to a composition series, we conclude that

$$\chi = \sum_{Z \in \Lambda(B)} C_{Z,Y} \cdot \chi_Z$$

and hence

$$|e_X B \cap L_Y| = \chi(e_X) = \sum_{Z \in \Lambda(B)} C_{Z,Y} \cdot \chi_Z(e_X) = \sum_{Z \leq X} C_{Z,Y}.$$

An application of Rota's Möbius inversion theorem yields (4.8).

For the final statement, observe that $e_Z B \cap L_Y = \emptyset$ if $Z \not\geq Y$. Thus $C_{X,Y} = 0$ unless $X \geq Y$ and $C_{X,X} = |e_X B \cap L_X| \cdot \mu(X, X) = 1$. This completes the proof. $\square$

5. PROJECTIVE RESOLUTIONS AND GLOBAL DIMENSION

In this section we use various simplicial and CW complexes associated to left regular bands to compute projective resolutions of their simple modules. As a consequence, we obtain a more direct proof of the results of our previous paper on the global dimension of left regular bands [75], which went through classifying spaces of small categories. Also by allowing left regular bands that are not monoids, the theory can include a number of new and interesting examples.

5.1. Actions of left regular bands on CW posets. It turns out that it is more convenient to work with actions of left regular bands on CW posets than on regular CW complexes as the former are more combinatorial in nature. Thanks to the functor from CW posets to regular CW complexes, we can then obtain actions on the associated CW complexes and hence on their cellular chain complexes.

If B is a left regular band acting on a set X and $x \in X$, then

$$\text{Stab}(x) = \{b \in B \mid bx = x\}$$

denotes the *stabilizer* of x . We say that the action of B on X is *unitary* if $BX = X$. Note that this is equivalent to $\text{Stab}(x) \neq \emptyset$ for all $x \in X$. Indeed, if $BX = X$ and $x = by$ with $y \in X$, then $x = bx$ and hence $\text{Stab}(x) \neq \emptyset$. Conversely, if $b \in \text{Stab}(x)$, then $x = bx \in BX$. Notice that if B is a monoid, then the action of B on X is unitary if and only if the identity acts via the identity mapping. Clearly $BX = X$ if the identity of B acts by the identity mapping. If $\text{Stab}(x) \neq \emptyset$, say $bx = x$, then $1x = 1(bx) = (1b)x = bx = x$. Therefore, the identity acts via the identity mapping. Notice that the action of B on the left of itself is unitary because $b \in \text{Stab}(b)$ for all $b \in B$. The importance of unitary actions stems from the following proposition.

Proposition 5.1. *Let B be a connected left regular band and $\mathbb{k}$ a commutative ring with unit. Suppose that B has a unitary action on a set X . Then $\mathbb{k}X$ is a unitary $\mathbb{k}B$ -module.*

Proof. Let η be the identity of $\mathbb{k}B$ and let $x \in X$. Suppose that $b \in B$ with $bx = x$. Then $\eta x = \eta(bx) = (\eta b)x = bx = x$ and so η fixes the basis X of $\mathbb{k}X$. It follows that $\mathbb{k}X$ is a unitary $\mathbb{k}B$ -module. $\square$

The following proposition will be useful several times.

Proposition 5.2. *Let B be a connected left regular band acting by cellular maps on a poset P . Let $\sigma_0 < \sigma_1 < \cdots < \sigma_q$ form a simplex σ of $\Delta(P)$. Then $\text{Stab}(\sigma) = \text{Stab}(\sigma_q)$. In particular, the action of B on P is unitary if and only if all of the simplices of $\Delta(P)$ have non-empty stabilizers under the action of B on $\Delta(P)$ by simplicial maps (i.e., the action of B on the set of simplices of $\Delta(P)$ is unitary).*

Proof. If $b \in B$ and $b\sigma_q = \sigma_q$, then Proposition 3.8 implies that b stabilizes σ . Conversely, if b stabilizes σ , then from $b\sigma_0 < b\sigma_1 < \cdots < b\sigma_q$, we deduce that $b\sigma_q = \sigma_q$. $\square$

It is crucial that B acts on itself by cellular mappings

Proposition 5.3. *Let B be a left regular band. Then the action of B on itself by left multiplication is unitary and by cellular maps.*

Proof. By Lemma 2.6 the action is order preserving and we have already observed that it is unitary. Let $a, b \in B$ and suppose that $b' \leq ab$. Then $bb' \leq b$ and $a(bb') = (ab)b' = b'$. Thus left multiplication by a is a cellular mapping. $\square$

Fix for this discussion a commutative ring with unit $\mathbb{k}$. Let us remind the reader of some basics of simplicial homology. See [81, Chapter 1.5] for details. Let K be a (finite) simplicial complex. If $\sigma = \{v_0, \dots, v_q\}$ is a q -simplex, then an *orientation* of σ is an equivalence class of total orderings of the vertices of σ , where two orderings are equivalent if they differ by an even permutation. Each q -simplex admits exactly two orientations if $q > 0$ (and one if $q = 0$). We write $[v_{i_0}, \dots, v_{i_q}]$ for the class of the ordering $v_{i_0}, \dots, v_{i_q}$. An *oriented q -simplex* consists of a q -simplex together with an orientation.

To construct the *simplicial chain complex* $C_\bullet(K; \mathbb{k})$ of K with coefficients in $\mathbb{k}$, we recall that the q^{th} -chain module $C_q(K; \mathbb{k})$ is the quotient of the free $\mathbb{k}$ -module on the set of oriented simplices by the relations identifying an oriented simplex σ with $-\sigma'$ where σ' has the same underlying simplex as σ but with the opposite orientation (for $q > 0$). Of course, by fixing an orientation of each simplex, it is easy to see that $C_q(K; \mathbb{k})$ is a free $\mathbb{k}$ -module with a basis in bijection with the set of q -simplices.

The boundary operator $d_q: C_q(K; \mathbb{k}) \rightarrow C_{q-1}(K; \mathbb{k})$ for $q \geq 1$ is given by

$$d_q([v_0, \dots, v_q]) = \sum_{i=0}^q (-1)^i [v_0, \dots, \widehat{v_i}, \dots, v_q]$$

where $\widehat{v_i}$ means omit v_i . The augmentation $\varepsilon: C_0(K; \mathbb{k}) \rightarrow \mathbb{k}$ is given by $\varepsilon(v) = 1$ for any 0-simplex v (where we recall that an oriented 0-simplex is the same thing as a 0-simplex). The (reduced) simplicial homology $\widetilde{H}_\bullet(K; \mathbb{k})$ of K with coefficients in $\mathbb{k}$ is then the homology of the (augmented) simplicial chain complex. The reduced cohomology of K with coefficients in $\mathbb{k}$ is defined dually.

If $f: K \rightarrow L$ is a *simplicial map*, then the induced chain map

$$f_*: C_\bullet(K; \mathbb{k}) \rightarrow C_\bullet(L; \mathbb{k})$$

is defined on an oriented q -simplex by

$$f_*([v_0, \dots, v_q]) = \begin{cases} [f(v_0), \dots, f(v_q)], & \text{if } v_0, \dots, v_q \text{ are distinct} \\ 0, & \text{else.} \end{cases}$$

The simplicial chain complex construction is functorial on the category of simplicial complexes and simplicial maps.

If L is a subsimplicial complex of a simplicial complex K , then $C_\bullet(L; \mathbb{k})$ is a subchain complex of $C_\bullet(K; \mathbb{k})$. The *relative simplicial chain complex* $C_\bullet(K, L; \mathbb{k})$ is the quotient chain complex.

The next proposition will be used throughout the text, often without comment.

Proposition 5.4. *Let B be a connected left regular band acting on a simplicial complex K by simplicial maps such that the stabilizer of each simplex is non-empty. Let L be a subcomplex invariant under B . Then the simplicial chain complex $C_\bullet(K; \mathbb{k})$ and the relative simplicial chain complex $C_\bullet(K, L; \mathbb{k})$ are chain complexes of unitary $\mathbb{k}B$ -modules. Thus the relative homology $H_q(K, L; \mathbb{k})$ is a unitary $\mathbb{k}B$ -module for all $q \geq 0$.*

Proof. First note that $C_\bullet(K; \mathbb{k})$ is a chain complex of $\mathbb{k}B$ -modules by functoriality of simplicial homology under simplicial maps. Let us verify that the $\mathbb{k}B$ -module $C_q(K; \mathbb{k})$ is unitary. Indeed, let $\sigma = [v_0, \dots, v_q]$ be an oriented q -simplex and suppose that $b \in B$ stabilizes the underlying q -simplex $\{v_0, \dots, v_q\}$. Then $\{bv_0, \dots, bv_q\} = \{v_0, \dots, v_q\}$ and hence b fixes each of $v_0, \dots, v_q$, being an idempotent. Thus $b_*([v_0, \dots, v_q]) = [bv_0, \dots, bv_q] = [v_0, \dots, v_q]$ and so $b\sigma = \sigma$. Hence if η is the identity of $\mathbb{k}B$, then we have that $\eta\sigma = \eta(b\sigma) = (\eta b)\sigma = b\sigma = \sigma$. Therefore, $C_q(K; \mathbb{k})$ is a unitary $\mathbb{k}B$ -module for each $q \geq 0$ and so $C_\bullet(K; \mathbb{k})$ is a chain complex of unitary $\mathbb{k}B$ -modules. We conclude that the subchain complex $C_\bullet(L; \mathbb{k})$ consists of unitary $\mathbb{k}B$ -modules, as does the quotient complex $C_\bullet(K, L; \mathbb{k})$. It follows that the relative homology consists of unitary $\mathbb{k}B$ -modules. $\square$

With Proposition 5.4 in hand, we can now establish that if B is a connected left regular band, then each simplicial complex $\Delta(B_{\geq X})$ with $X \in \Lambda(B)$ is acyclic.

Theorem 5.5. *Let B be a connected left regular band. Then $\Delta(B_{\geq X})$ is acyclic for all $X \in \Lambda(B)$.*

Proof. As $B_{\geq X}$ is a connected left regular band in its own right, for all $X \in \Lambda(B)$, it suffices to prove that $\Delta(B)$ is acyclic. (One can then replace B by $B_{\geq X}$ to obtain the general case.) Note that $\Delta(B)$ is a subcomplex of $\Delta(B^1)$. Therefore, the augmented simplicial chain complex C of $\Delta(B)$ (over $\mathbb{Z}$) can be viewed as a subcomplex of the augmented simplicial chain complex C' of $\Delta(B^1)$ by identifying q -chains on $\Delta(B)$ with q -chains on $\Delta(B^1)$ that

are supported on oriented simplices of $\Delta(B)$. The complex C' is acyclic (even contractible) because $\Delta(B^1)$ is contractible, having cone point 1.

The complex C' is, in fact, a complex of $\mathbb{Z}B^1$ -modules by Proposition 5.4. Thus each element $x \in \mathbb{Z}B^1$ induces a chain map $C' \rightarrow C'$. Moreover, since $b\Delta(B^1) \subseteq \Delta(B)$ for all $b \in B$, it follows that if $x \in \mathbb{Z}B$ then the image of the chain map $C' \rightarrow C'$ induced by x is contained in C . In particular, let $\eta \in \mathbb{Z}B$ be the identity element guaranteed by Theorem 4.15. Then η induces an idempotent chain map $C' \rightarrow C'$ with image contained in C and, moreover, it fixes C by Proposition 5.4. Thus the chain complex C is a retract of the acyclic chain complex C' . This implies that C is acyclic (by the functoriality of homology). In fact, since C' is a contractible chain complex over $\mathbb{Z}$, so is its retract C . This completes the proof. $\square$

We do not know whether B connected implies $B_{\geq X}$ is contractible for all $X \in \Lambda(B)$, although this happens to be the case in all our examples. Notice that if we replace $\mathbb{Z}$ in the above proof with an arbitrary commutative ring with unit $\mathbb{k}$, we would get that $\tilde{H}_q(\Delta(B); \mathbb{k}) = 0$ for all $q \geq 0$ without appealing to the universal coefficient theorem.

Remark 5.6. The proof of Theorem 5.5 can easily be adapted to show that if S is any semigroup (possibly infinite) such that $\mathbb{Z}S$ has an identity, then the order complexes of both its posets of principal right ideals and principal left ideals are acyclic.

The cellular chain complex with coefficients in $\mathbb{k}$ for a CW poset P can be described as follows. For $q \geq 0$, put

$$P^q = \{\sigma \in P \mid \dim \sigma \leq q\}.$$

It is the face poset of the q -skeleton of $\Sigma(P)$. One has $C_q(\Sigma(P); \mathbb{k}) = H_q(\Delta(P^q), \Delta(P^{q-1}); \mathbb{k})$, where the left hand side is the cellular chain module and the right hand side is the relative simplicial homology module. The boundary map is the composition

$$\begin{aligned} H_q(\Delta(P^q), \Delta(P^{q-1}); \mathbb{k}) &\xrightarrow{\partial} H_{q-1}(\Delta(P^{q-1}); \mathbb{k}) \\ &\longrightarrow H_{q-1}(\Delta(P^{q-1}), \Delta(P^{q-2}); \mathbb{k}) \end{aligned}$$

where ∂ is the boundary map from the long exact sequence in relative simplicial homology. Recall that if τ is a q -cell of P , then

$$H_q(\Delta(P_{\leq \tau}), \Delta(P_{< \tau}); \mathbb{k}) \cong \mathbb{k}.$$

An orientation for τ is a choice of a basis element $[\tau]$ for the rank one free $\mathbb{k}$ -module $H_q(\Delta(P_{\leq \tau}), \Delta(P_{< \tau}); \mathbb{k})$. We also write $[\tau]$ for its image under the canonical embedding

$$H_q(\Delta(P_{\leq \tau}), \Delta(P_{< \tau}); \mathbb{k}) \longrightarrow H_q(\Delta(P^q), \Delta(P^{q-1}); \mathbb{k}) = C_q(\Sigma(P); \mathbb{k}).$$

The elements $[\tau]$ with τ a q -cell of P form a basis for $C_q(\Sigma(P); \mathbb{k})$. See [77] or [81, Chapter 4.39], for instance, for background on cellular homology.

We now assume that B is a connected left regular band with a unitary action on a CW poset P . There is an induced action $B \curvearrowright \Sigma(P)$ by regular cellular maps by functoriality of Σ . Notice that $\mathbb{k}$ has the structure of a trivial $\mathbb{k}B$ -module. The augmented cellular chain complex

$$C_\bullet(\Sigma(P); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}$$

for $\Sigma(P)$ with coefficients in $\mathbb{k}$ then becomes a chain complex of unitary $\mathbb{k}B$ -modules (finitely generated over $\mathbb{k}$) by functoriality of cellular homology with respect to cellular maps and Propositions 5.2 and 5.4. Alternatively, a cellular map $f: P \rightarrow P$ obviously sends P^q into P^q and so the induced simplicial map sends $\Delta(P^q)$ into $\Delta(P^q)$. If P is acyclic, then

$$C_\bullet(\Sigma(P); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}$$

is a resolution of the trivial module $\mathbb{k}$. As mentioned above, $C_q(\Sigma(P); \mathbb{k})$ has a basis consisting of the oriented q -cells $[\sigma]$ of P . Moreover, if $b \in B$, then $b[\sigma] = 0$ if $\dim b\sigma < \dim \sigma$ and otherwise $b[\sigma] \in \mathbb{k}[b\sigma]$ because B acts by regular cellular mappings.

Let us say that an action $B \curvearrowright P$ of a left regular band B on a poset P is *semi-free* if $\text{Stab}(\tau)$ has a minimum element (or equivalently a zero element) for each $\tau \in P$. In particular, each stabilizer is assumed non-empty and hence the action is unitary. Notice that in this case if $Q \subseteq P$ is a B -invariant lower set, then the action of B on Q is also by cellular maps and semi-free. For example, the action of B on the left of itself (cf. Proposition 5.3) is semi-free: b is the minimum element of $\text{Stab}(b)$ for $b \in B$. We say that an action of B on a simplicial complex K is semi-free if the induced action on the face poset $\mathcal{P}(K)$ is *semi-free*.

Lemma 5.7. *Suppose that the action $B \curvearrowright P$ of a left regular band B on a poset P is semi-free. Then the action of B on $\mathcal{P}(\Delta(P))$ is also semi-free, i.e., the action of B on $\Delta(P)$ is semi-free.*

Proof. This follows from Proposition 5.2, which states that the stabilizer of a simplex coincides with the stabilizer of its maximum element. $\square$

In particular, the action of B on $\Delta(B)$ is semi-free. Note that if B has an identity, then $\Delta(B)$ is contractible because the identity 1 is a cone point. More generally, if B is connected, then $\Delta(B)$ is acyclic by Theorem 5.5.

If B acts on a poset P and $a, b \in B$ with $Ba = Bb$, then the subposets aP and bP are isomorphic. Indeed, the left action of b on P induces a mapping $aP \rightarrow bP$ whose inverse is given by the left action of a because $ab = a$ and $ba = b$. If $A \subseteq B$, we put $AP = \bigcup_{a \in A} aP$; note that AP is a lower set. If A is a left ideal, then AP is B -invariant.

5.2. Projective resolutions. Our goal is to show that if a connected left regular band B has a semi-free action on an acyclic CW poset P , then the augmented cellular chain complex $C_\bullet(\Sigma(P); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}$ is a projective resolution of the trivial $\mathbb{k}B$ -module $\mathbb{k}$. In particular, this applies to $P =$

$\mathcal{P}(\Delta(B))$ and so the augmented simplicial chain complex $C_\bullet(\Delta(B); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}$ is a projective resolution of the trivial $\mathbb{k}B$ -module $\mathbb{k}$. We shall give some other examples as well.

We begin with a lemma about recognizing Schützenberger representations. If $e \in B$ and $L_e = \sigma^{-1}(Be)$ is the $\mathcal{L}$ -class of e , then B acts on L_e by partial maps (by restricting the regular action of B on the left of itself). That is, if $b \in B$ and $a \in L_e$, then

$$b \cdot a = \begin{cases} ba, & \text{if } Bb \supseteq Be \\ \text{undefined,} & \text{else.} \end{cases}$$

The Schützenberger representation defined in Section 4 is then the linearization of this action by partial maps.

Lemma 5.8. *Let B be a left regular band and let $Be \in \Lambda(B)$. Suppose that B acts on a finite non-empty set Y by partial transformations such that:*

- (1) *by is defined, for $y \in Y$, if and only if $Bb \supseteq Be$;*
- (2) *$aY \cap bY \neq \emptyset$ if and only if $a = b$ for all $a, b \in L_e$;*
- (3) *$BeY = L_eY = Y$.*

Then Y is B -equivariantly isomorphic to a disjoint union of $|eY|$ copies of the action of B on L_e . More precisely, Bx provides an isomorphic copy of L_e for each $x \in eY$ and distinct elements of eY provide disjoint copies.

Proof. Fix $y \in eY$. First we observe that B acts transitively on By . Indeed, if $Bb \supseteq Be$, then $By \supseteq Bby \supseteq Bey = By$ and so $By = Bby$. Next, we define a map $F: By \rightarrow L_e$ by $F(by) = be$ for $Bb \supseteq Be$. To see that this is well defined, observe that if $ay = by$ with $Ba \supseteq Be$, then $ae = bey$ and $ae, be \in L_e$. Thus by assumption (2), we conclude $ae = be$ and so F is well defined. It is injective because $ae = F(ay) = F(by) = be$ implies $ay = aey = bey = by$. It is surjective because $b = be = F(by)$ for $b \in L_e$. If $a, b \in B$ with $Bb \supseteq Be$, then $a(by)$ is defined if and only if $Ba \supseteq Be$, which is if and only if $a \cdot be$ is defined, in which case $F(a(by)) = F((ab)y) = (ab)e = a(be) = a \cdot F(by)$. This proves that F is B -equivariant.

Next observe that if $y' \in eY$ and $By' \cap By \neq \emptyset$, then $By' = By$ by transitivity. But if $by = y'$, then since $b \in L_e$, we have $eb = e$ and so $y = ey = eby = ey' = y'$. Thus, from $BeY = Y$, we deduce that Y is a disjoint union of $|eY|$ isomorphic copies of the action of B on L_e , one for each $y \in eY$, as required. $\square$

Recall that a *principal series* for a finite semigroup S is an unrefinable chain $I_0 \subset I_1 \subset \cdots \subset I_n = S$ of two-sided ideals. Set $I_{-1} = \emptyset$ for convenience. It is known [46, Theorem 2.40] that $I_m \setminus I_{m-1}$ is a $\mathcal{J}$ -class for $0 \leq m \leq n$ and each $\mathcal{J}$ -class occurs in this way exactly once; moreover, I_0 is always the minimal ideal of S . For a left regular band B , ideals are the same thing as left ideals and so a principal series is just a maximal chain

$$\widehat{0} = L_0 \subset L_1 \subset \cdots \subset L_n = B \quad (5.1)$$

of left ideals of B . The $\mathcal{L}$ -classes each appear exactly once as $L_m \setminus L_{m-1}$ (where $L_{-1} = \emptyset$).

Lemma 5.9. *Let R be a unital ring and let M be an R -module. Suppose that M has a filtration*

$$0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_n = M$$

with M_i/M_{i-1} a projective module for $i = 1, \dots, n$. Then we have

$$M \cong \bigoplus_{i=1}^n M_i/M_{i-1}$$

and hence M is projective.

Proof. We proceed via induction on n . If $n = 1$, there is nothing to prove. In general, since the sequence

$$0 \longrightarrow M_{n-1} \longrightarrow M_n \longrightarrow M_n/M_{n-1} \longrightarrow 0$$

is exact and M_n/M_{n-1} is projective, we have $M_n \cong M_{n-1} \oplus M_n/M_{n-1}$. The result now follows by induction applied to the filtration

$$0 = M_0 \subseteq M_1 \subseteq \cdots \subseteq M_{n-1}$$

of M_{n-1} . □

The following theorem is one of the principal results of this text.

Theorem 5.10. *Let B be a connected left regular band and $\mathbb{k}$ a commutative ring with unit. Suppose that $B \curvearrowright P$ is a semi-free action of B on an acyclic CW poset P . Then the augmented cellular chain complex $C_\bullet(\Sigma(P); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}$ is a projective resolution of the trivial $\mathbb{k}B$ -module.*

More precisely, if we fix e_X with $Be_X = X$ for all $X \in \Lambda(B)$, then

$$C_q(\Sigma(P); \mathbb{k}) \cong \bigoplus_{X \in \Lambda(B)} n_X \cdot \mathbb{k}L_X$$

where $\mathbb{k}L_X$ is the Schützenberger representation associated to X and n_X is the number of q -cells σ with e_X as the minimum element of $\text{Stab}(\sigma)$.

Proof. The augmented cellular chain complex is a resolution of $\mathbb{k}$ because P is acyclic. Fix a principal series for B as in (5.1). Let P_m be the subposet $L_m P$ and note that $L_n P = BP = P$ since the action is unitary. We take $P_{-1} = \emptyset$. Note that $\sigma \in P_m$ if and only if $\text{Stab}(\sigma) \cap L_m \neq \emptyset$ and so $\sigma \in P_m \setminus P_{m-1}$ if and only if the minimum element of $\text{Stab}(\sigma)$ is in $L_m \setminus L_{m-1}$.

Recall that the relative chain complex $C_\bullet(\Sigma(P_m), \Sigma(P_{m-1}); \mathbb{k})$ is defined to be $C_\bullet(\Sigma(P_m); \mathbb{k})/C_\bullet(\Sigma(P_{m-1}); \mathbb{k})$. The filtration

$$\Sigma(P_0) \subseteq \Sigma(P_1) \subseteq \cdots \subseteq \Sigma(P_n) = \Sigma(P)$$

of $\Sigma(P)$ by B -invariant subcomplexes induces a filtration of chain complexes of unitary $\mathbb{k}B$ -modules

$$0 = C_\bullet(\Sigma(P_{-1}); \mathbb{k}) \subseteq C_\bullet(\Sigma(P_0); \mathbb{k}) \subseteq \cdots \subseteq C_\bullet(\Sigma(P_n); \mathbb{k}) = C_\bullet(\Sigma(P); \mathbb{k})$$

with subquotients the relative chain complexes $C_\bullet(\Sigma(P_m), \Sigma(P_{m-1}); \mathbb{k})$. So, to show that $C_q(\Sigma(P); \mathbb{k})$ is projective, it suffices to prove that the $\mathbb{k}B$ -modules $C_q(\Sigma(P_m), \Sigma(P_{m-1}); \mathbb{k})$ are projective for all $m, q \geq 0$. It will then follow that

$$C_q(\Sigma(P); \mathbb{k}) \cong \bigoplus_{m=0}^n C_q(\Sigma(P_m), \Sigma(P_{m-1}); \mathbb{k}) \quad (5.2)$$

is projective by Lemma 5.9.

If P_m has the same set of q -cells as P_{m-1} , then there is nothing to prove. So assume that $P_m \setminus P_{m-1}$ contains some q -cells. Let Y be the set of q -cells of P_m that do not belong to P_{m-1} . As P_m and P_{m-1} are B -invariant, we have that B acts by partial maps on Y by restricting the action of B on P to Y . That is, if $b \in B$ and $\tau \in Y$, then

$$b \cdot \tau = \begin{cases} b\tau, & \text{if } b\tau \in Y \\ \text{undefined,} & \text{if } b\tau \in P_{m-1}. \end{cases}$$

Suppose that $L_m \setminus L_{m-1} = \sigma^{-1}(X)$ with $X \in \Lambda(B)$, that is, $L_m \setminus L_{m-1}$ is the $\mathcal{L}$ -class of e_X . We claim that the conditions of Lemma 5.8 hold with $e = e_X$.

If $\tau \in Y$, then $\tau = a\tau$ for some $a \in L_m \setminus L_{m-1} = L_e$. Thus the third condition is satisfied. We continue to assume that $a\tau = \tau$ with $a \in L_e$. If $Bb \supseteq Be = Ba$, then $cb = a$ for some $c \in B$. But then $\tau = a\tau = (cb)\tau$ and so $b\tau$ is defined, i.e., $b\tau \in P_m \setminus P_{m-1}$. On the other hand, if $Bb \not\supseteq Be$, then $ba \in L_{m-1}$ and so $b\tau = ba\tau \in P_{m-1}$. Thus b is undefined at τ for the action on Y . This establishes the first condition of Lemma 5.8.

For the second condition, suppose that $a, b \in L_e = L_m \setminus L_{m-1}$ and $\tau \in aY \cap bY$. Then $a, b \in \text{Stab}(\tau)$. Let c be the minimum element of $\text{Stab}(\tau)$ (guaranteed by the definition of a semi-free action). Then $c \leq a, b$. We recall that in a left regular band, $x < y$ implies $\sigma(x) < \sigma(y)$ by Proposition 3.6. Therefore, if $a \neq b$, then $c < a, b$ and so $c \in L_m \setminus L_e = L_{m-1}$. But then $\tau = c\tau \in P_{m-1}$, contradicting $\tau \in Y$. Thus $a = b$. Note that $\sigma \in eY$ if and only if σ is a q -cell and e is the minimum element of $\text{Stab}(\sigma)$. We conclude that the module $\mathbb{k}Y$ is isomorphic to $n_X \cdot \mathbb{k}L_X$ by Lemma 5.8.

Now we prove that we can find orientations for the q -cells of Y such that the action of B (restricted to Y) is orientation preserving. For each $\tau \in eY$ fix an orientation $[\tau]$ of τ . Recall that an orientation for τ is a choice of a basis element $[\tau]$ for $H_q(\Delta(P_{\leq \tau}), \Delta(P_{< \tau}); \mathbb{k}) \cong \mathbb{k}$. Now, for $a \in L_m \setminus L_{m-1}$, we have that the action of a on P gives rise to a poset isomorphism of $P_{\leq \tau}$ and $P_{\leq a\tau}$ and hence a homeomorphism of pairs

$$(\Delta(P_{\leq \tau}), \Delta(P_{< \tau})) \longrightarrow (\Delta(P_{\leq a\tau}), \Delta(P_{< a\tau}))$$

(with inverse given by the action of e). Thus we can set $[a\tau] = a[\tau]$ as an orientation for $a\tau$. By Lemma 5.8, for each $\tau' \in Y$, there exist unique $a \in L_e$ and $\tau \in eY$ such that $\tau' = a\tau$. Thus our scheme provides a well-defined orientation for each element of Y . Then if $b \in B$ with $Bb \supseteq Be$, $a \in L_e$

and $\tau \in eY$, we have $b[a\tau] = ba[\tau] = [ba\tau]$ where the last equality follows because $ba \in L_m \setminus L_{m-1}$. Thus b sends the chosen orientation of $a\tau$ to the chosen orientation of $ba\tau$. If $Bb \not\supseteq Be$, then $bY \subseteq P_{m-1}$ and so b annihilates $C_q(\Sigma(P_m), \Sigma(P_{m-1}); \mathbb{k})$. It now follows that $C_q(\Sigma(P_m), \Sigma(P_{m-1}); \mathbb{k}) \cong \mathbb{k}Y \cong n_X \cdot \mathbb{k}L_X$ where the first isomorphism sends $[\tau]$ to τ . In light of (5.2), this completes the proof. $\square$

The following special case is of central importance. We recall that if B is a connected left regular band, then $\Delta(B)$ is acyclic by Theorem 5.5.

Theorem 5.11. *Let B be a connected left regular band and $\mathbb{k}$ a commutative ring with unit. Then the augmented simplicial chain complex*

$$C_\bullet(\Delta(B); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}$$

is a projective resolution of the trivial $\mathbb{k}B$ -module.

More precisely, if we fix e_X with $Be_X = X$ for all $X \in \Lambda(B)$, then

$$C_q(\Delta(B); \mathbb{k}) \cong \bigoplus_{X \in \Lambda(B)} n_X \cdot \mathbb{k}L_X$$

where $\mathbb{k}L_X$ is the Schützenberger representation associated to X and n_X is the number of q -simplices in $\Delta(e_X B)$ with e_X as a vertex.

Notice that $C_0(\Delta(B); \mathbb{k}) = \mathbb{k}B$ and hence the resolution in Theorem 5.11 is never minimal if B is non-trivial.

Theorem 5.10, in conjunction with Corollary 4.9 and Theorem 5.5, yields the following corollary.

Corollary 5.12. *Let B be a connected left regular band and $\mathbb{k}$ a commutative ring with unit. Suppose that $X \in \Lambda(B)$. If $B_{\geq X} \curvearrowright P$ is a semi-free action on an acyclic CW poset complex P , then the augmented cellular chain complex $C_\bullet(\Sigma(P); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}_X$ is a finite projective resolution of $\mathbb{k}_X$ by finitely generated $\mathbb{k}B$ -modules. In particular, $C_\bullet(\Delta(B_{\geq X}); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}_X$ is a finite projective resolution of $\mathbb{k}_X$ by finitely generated $\mathbb{k}B$ -modules.*

Fix, for each $Y \in \Lambda(B_{\geq X})$, an element e_Y with $Y = Be_Y$. The number of occurrences of the Schützenberger representation $\mathbb{k}L_Y$ as a direct summand in $C_q(\Sigma(P); \mathbb{k})$ is the number of q -cells τ of P such that e_Y is the minimum element of $\text{Stab}(\tau)$. In particular, the number of occurrences of $\mathbb{k}L_Y$ as a direct summand in $C_q(\Delta(B_{\geq X}); \mathbb{k})$ is the number of q -simplices of $\Delta(e_Y B_{\geq X})$ with e_Y as a vertex.

An important special case is when $B_{\geq X}$ is a CW poset.

Corollary 5.13. *Let B be a connected left regular band and $\mathbb{k}$ a commutative ring with unit. Suppose that $X \in \Lambda(B)$ and $B_{\geq X}$ is a CW poset. Then the augmented cellular chain complex $C_\bullet(\Sigma(B_{\geq X}); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}_X$ is a finite projective resolution of $\mathbb{k}_X$ by finitely generated $\mathbb{k}B$ -modules.*

Moreover, we have the direct sum decomposition

$$C_q(\Sigma(B_{\geq X}); \mathbb{k}) \cong \bigoplus_{\text{rk}[X,Y]=q} \mathbb{k}L_Y \quad (5.3)$$

where we note that $\Lambda(B)_{\geq X}$ is a graded poset by Proposition 3.6 and Corollary 3.7 and where $\mathbb{k}L_Y$ denotes the Schützenberger representation associated to $Y \in \Lambda(B)$.

Proof. The CW poset $B_{\geq X}$ is acyclic by Theorem 5.5. Corollary 5.12 establishes everything except the decomposition (5.3). Fix $e_Y \in B$ with $Y = Be_Y$. If $Y \geq X$, then e_Y is the unique cell of $B_{\geq X}$ with e_Y as the minimum element of its stabilizer and the dimension of this cell in $B_{\geq X}$ is $\text{rk}[X, Y]$ by Proposition 3.6 and Corollary 3.7. Applying again Corollary 5.12 completes the proof of (5.3). $\square$

Remark 5.14. The minimal element of the stabilizer of $b \in B_{\geq X}$ is b and so it easily follows from how we orient the cells in the proof of Theorem 5.10 that the isomorphism in (5.3) takes the oriented cell $[b]$ to b for $b \in L_Y$ and $Y \geq X$.

5.3. Ext and global dimension. We now compute $\text{Ext}_{\mathbb{k}B}^n(\mathbb{k}_X, \mathbb{k}_Y)$ for $X, Y \in \Lambda(B)$. Restricting to the case that B is a monoid, we obtain an alternate proof of the results of [75]. First we need a lemma.

Lemma 5.15. *Let $Y \in \Lambda(B)$ and let V be a $\mathbb{k}B$ -module. Fix e_Y with $Be_Y = Y$. Then $\text{Hom}_{\mathbb{k}B}(V, \mathbb{k}_Y)$ can be identified as a $\mathbb{k}$ -module with the set of $\mathbb{k}$ -linear mappings $f: e_Y V \rightarrow \mathbb{k}$ that vanish on $\mathbb{k}[\partial e_Y B]V$ (where the latter is interpreted as 0 if $\partial e_Y B = \emptyset$).*

Proof. Let $f \in \text{Hom}_{\mathbb{k}B}(V, \mathbb{k}_Y)$ and $v \in V$. Then $f(e_Y v) = e_Y f(v) = f(v)$ and so f is uniquely determined by its deletion to $e_Y V$. Moreover, if $b \in \partial e_Y B$, then $f(bv) = bf(v) = 0$. Therefore, it remains to show that if $g: e_Y V \rightarrow \mathbb{k}$ vanishes on $\mathbb{k}[\partial e_Y B]V$ and is $\mathbb{k}$ -linear, then defining $f(v) = g(e_Y v)$ results in a $\mathbb{k}B$ -module homomorphism $f: V \rightarrow \mathbb{k}_Y$. Let $b \in B$. If $Bb \not\supseteq Y$, then $e_Y b \in \partial e_Y B$ and so $f(bv) = g(e_Y bv) = 0 = bf(v)$. On the other hand, if $Bb \supseteq Y$, then $e_Y b = e_Y$ and so $f(bv) = g(e_Y bv) = g(e_Y v) = bf(v)$. This completes the proof of the lemma. $\square$

The next theorem is a fundamental result of this paper. Throughout this text we follow the standard convention that

$$\tilde{H}^{-1}(X; \mathbb{k}) = \begin{cases} \mathbb{k}, & \text{if } X = \emptyset \\ 0, & \text{else.} \end{cases}$$

Theorem 5.16. *Let B be a connected left regular band and $\mathbb{k}$ a commutative ring with unit. Let $X, Y \in \Lambda(B)$. Fix e_Y with $Be_Y = Y$. Suppose that*

$B_{\geq X} \curvearrowright P$ is a semi-free action of $B_{\geq X}$ on an acyclic CW poset P . Then

$$\begin{aligned} \text{Ext}_{\mathbb{k}B}^n(\mathbb{k}_X, \mathbb{k}_Y) &= \begin{cases} H^n(\Sigma(e_Y P), \Sigma((\partial e_Y B_{\geq X})P); \mathbb{k}), & \text{if } Y \geq X \\ 0, & \text{else.} \end{cases} \\ &= \begin{cases} \mathbb{k}, & \text{if } X = Y, n = 0 \\ \tilde{H}^{n-1}(\Sigma((\partial e_Y B_{\geq X})P); \mathbb{k}), & \text{if } Y > X, n \geq 1 \\ 0, & \text{else.} \end{cases} \end{aligned}$$

Proof. We compute $\text{Ext}_{\mathbb{k}B}^n(\mathbb{k}_X, \mathbb{k}_Y)$ using the projective resolution

$$C_\bullet(\Sigma(P); \mathbb{k}) \xrightarrow{\varepsilon} \mathbb{k}_X$$

from Corollary 5.12. Thus $\text{Ext}_{\mathbb{k}B}^n(\mathbb{k}_X, \mathbb{k}_Y) = H^n(\text{Hom}_{\mathbb{k}B}(C_\bullet(\Sigma(P); \mathbb{k}), \mathbb{k}_Y))$.

First suppose that $Y \not\geq X$. Then $e_Y C_q(\Sigma(P); \mathbb{k}) = 0$ and so we have $\text{Hom}_{\mathbb{k}B}(C_q(\Sigma(P); \mathbb{k}), \mathbb{k}_Y) = 0$ for all $q \geq 0$ by Lemma 5.15. It follows that $\text{Ext}_{\mathbb{k}B}^n(\mathbb{k}_X, \mathbb{k}_Y) = 0$ for all $n \geq 0$ in this case.

Next suppose that $Y \geq X$. Then observe that

$$\begin{aligned} e_Y C_q(\Sigma(P); \mathbb{k}) &= C_q(\Sigma(e_Y P); \mathbb{k}) \\ \mathbb{k}[\partial e_Y B] C_q(\Sigma(P); \mathbb{k}) &= C_q(\Sigma((\partial e_Y B_{\geq X})P); \mathbb{k}). \end{aligned}$$

Thus Lemma 5.15 identifies $\text{Hom}_{\mathbb{k}B}(C_\bullet(\Sigma(P); \mathbb{k}), \mathbb{k}_Y)$ with the relative cochain complex $C^\bullet(\Sigma(e_Y P), \Sigma(\partial e_Y B_{\geq X} P); \mathbb{k})$ and so

$$\text{Ext}_{\mathbb{k}B}^n(\mathbb{k}_X, \mathbb{k}_Y) = H^n(\Sigma(e_Y P), \partial \Sigma(e_Y B_{\geq X} P); \mathbb{k})$$

for all $n \geq 0$ in this case. The first equality is now proved.

For the second equality, we recall that $e_Y P$ is acyclic by Proposition 3.8. In the case $X = Y$, we have $\partial e_Y B_{\geq X} = \emptyset$ and so

$$H^n(\Sigma(e_Y P), \Sigma(\partial e_Y B_{\geq X} P); \mathbb{k}) = H^n(\Sigma(e_Y P); \mathbb{k})$$

which is $\mathbb{k}$ in dimension zero and zero in all other dimensions.

If $Y > X$, then $\partial e_Y B_{\geq X} \neq \emptyset$. The long exact sequence for reduced relative cohomology, together with the acyclicity of $e_Y P$, then yields

$$H^n(\Sigma(e_Y P), \Sigma((\partial e_Y B_{\geq X})P); \mathbb{k}) \cong \tilde{H}^{n-1}(\Sigma((\partial e_Y B_{\geq X})P); \mathbb{k})$$

for all $n \geq 1$. We have $H^0(\Sigma(e_Y P), \Sigma((\partial e_Y B_{\geq X})P); \mathbb{k}) = 0$ because $\Sigma(e_Y P)$ is connected and $\Sigma((\partial e_Y B_{\geq X})P)$ is non-empty. $\square$

Specializing to $K = \Delta(B_{\geq X})$ and B a monoid recovers the main result of [75, Theorem 4.1], which was proved using classifying spaces of small categories, Quillen's Theorem A [90] and homological trickery.

Theorem 5.17. *Let B be a connected left regular band and $\mathbb{k}$ a commutative ring with unit. Let $X, Y \in \Lambda(B)$. Fix e_Y with $Be_Y = Y$. Then*

$$\begin{aligned} \text{Ext}_{\mathbb{k}B}^n(\mathbb{k}_X, \mathbb{k}_Y) &= \begin{cases} H^n(\Delta(e_Y B_{\geq X}), \Delta(\partial e_Y B_{\geq X}); \mathbb{k}), & \text{if } Y \geq X \\ 0, & \text{else.} \end{cases} \\ &= \begin{cases} \mathbb{k}, & \text{if } X = Y, n = 0 \\ \tilde{H}^{n-1}(\Delta(\partial e_Y B_{\geq X}); \mathbb{k}), & \text{if } Y > X, n \geq 1 \\ 0, & \text{else.} \end{cases} \end{aligned}$$

Proof. Take $P = \Delta(B_{\geq X})$ in Theorem 5.16 and use that $\Delta(B_{\geq X})$ is acyclic by Theorem 5.5. $\square$

As a corollary, we recover a description of the quiver of a left regular band monoid algebra [75, 96], now under the weaker assumption of connectedness of the left regular band.

Theorem 5.18. *Let B be a connected left regular band and $\mathbb{k}$ a field. Then the quiver of $\mathbb{k}B$ has vertex set $\Lambda(B)$. There are no arrows $X \rightarrow Y$ unless $X < Y$, in which case the number of arrows $X \rightarrow Y$ is one fewer than the number of connected components of $\Delta(\partial e_Y B_{\geq X})$ where $Y = Be_Y$.*

Proof. This follows from Theorem 5.17 as $\dim \tilde{H}^0(K)$ is one fewer than the number of connected components of K for a simplicial complex K . $\square$

Corollary 5.19. *B be a connected left regular band and $\mathbb{k}$ a commutative ring with unit. Let $X, Y \in \Lambda(B)$. Fix e_Y with $Be_Y = Y$. Let $-\infty$ be the minimum element of $\Lambda(S(B)) = \Lambda(B) \cup \{-\infty\}$. Then*

$$\begin{aligned} (1) \quad \text{Ext}_{\mathbb{k}S(B)}^n(\mathbb{k}_X, \mathbb{k}_Y) &= \begin{cases} \mathbb{k}, & \text{if } X = Y, n = 0 \\ \tilde{H}^{n-1}(\Delta(\partial e_Y B_{\geq X}); \mathbb{k}), & \text{if } Y > X, n \geq 1 \\ 0, & \text{else;} \end{cases} \\ (2) \quad \text{Ext}_{\mathbb{k}S(B)}^n(\mathbb{k}_{-\infty}, \mathbb{k}_Y) &= \begin{cases} \tilde{H}^{n-2}(\Delta(\partial e_Y B); \mathbb{k}), & \text{if } n \geq 1 \\ 0, & \text{else;} \end{cases} \\ (3) \quad \text{Ext}_{\mathbb{k}S(B)}^n(\mathbb{k}_{-\infty}, \mathbb{k}_{-\infty}) &= \begin{cases} \mathbb{k}, & \text{if } n = 0; \\ 0, & \text{else;} \end{cases} \\ (4) \quad \text{Ext}_{\mathbb{k}S(B)}^n(\mathbb{k}_X, \mathbb{k}_{-\infty}) &= 0 \text{ for all } X \in \Lambda(B). \end{aligned}$$

Proof. This follows from Theorem 5.17 and the topological fact (which can be easily proved via the Meyer-Vietoris sequence) that $\tilde{H}^{q+1}(S(X); \mathbb{k}) \cong \tilde{H}^q(X; \mathbb{k})$. $\square$

Our next corollary concerns joins.

Corollary 5.20. *Let B' be a left regular band, B be a connected left regular band and $\mathbb{k}$ a field.*

(1) If $X, Y \in \Lambda(B)$, then

$$\mathrm{Ext}_{\mathbb{k}(B' * B)}^n(\mathbb{k}_X, \mathbb{k}_Y) = \begin{cases} \mathbb{k}, & \text{if } X = Y, n = 0 \\ \tilde{H}^{n-1}(\Delta(\partial e_Y B_{\geq X}); \mathbb{k}), & \text{if } Y > X, n \geq 1 \\ 0, & \text{else.} \end{cases}$$

(2) If $X, Y \in \Lambda(B')$, then

$$\mathrm{Ext}_{\mathbb{k}(B' * B)}^n(\mathbb{k}_X, \mathbb{k}_Y) = \begin{cases} \mathbb{k}, & \text{if } X = Y, n = 0 \\ \tilde{H}^{n-1}(\Delta(\partial e_Y B'_{\geq X}); \mathbb{k}), & \text{if } Y > X, n \geq 1 \\ 0, & \text{else.} \end{cases}$$

(3) If $X \in \Lambda(B')$ and $Y \in \Lambda(B)$, then

$$\mathrm{Ext}_{\mathbb{k}(B' * B)}^n(\mathbb{k}_X, \mathbb{k}_Y) = \bigoplus_{p+q=n-2} \tilde{H}^p(\Delta(B'_{\geq X}); \mathbb{k}) \otimes_{\mathbb{k}} \tilde{H}^q(\Delta(\partial e_Y B); \mathbb{k})$$

if $n \geq 1$ and otherwise is zero.

(4) If $X \in \Lambda(B)$ and $Y \in \Lambda(B')$, then $\mathrm{Ext}_{\mathbb{k}(B' * B)}^n(\mathbb{k}_X, \mathbb{k}_Y) = 0$.

Proof. This follows from Theorem 5.17, the observation that

$$\partial e_Y(B' * B)_{\geq X} = B'_{\geq X} * \partial e_Y B'$$

for $X \in \Lambda(B')$ and $Y \in \Lambda(B)$ and the fact that

$$\tilde{H}^{n+1}(X * Y; \mathbb{k}) \cong \bigoplus_{p+q=n} \tilde{H}^p(X; \mathbb{k}) \otimes_{\mathbb{k}} \tilde{H}^q(Y; \mathbb{k})$$

(cf. [22, Equation (9.12)]). Here again we use the usual conventions for cohomology in negative degrees. $\square$

Theorem 5.17 allows us to compute the global dimension of $\mathbb{k}B$ when $\mathbb{k}$ is a field. This result was first obtained in [75, Theorem 4.4] for left regular band monoids.

Corollary 5.21. *If $\mathbb{k}$ is a field and B is a connected left regular band, then the global dimension of $\mathbb{k}B$ is the largest n such that the reduced cohomology $\tilde{H}^{n-1}(\Delta(\partial e_Y B_{\geq X}); \mathbb{k}) \neq 0$ for some $X, Y \in \Lambda(B)$ with $Y \geq X$.*

Corollary 5.21 was used in [75] to compute the global dimension of almost all the examples of left regular band monoids that we have been considering in this paper. We recall quickly some of the results. First we give an upper bound in terms of Leray numbers, which improves upon the upper bound that [85] would provide in the case of a left regular band.

Corollary 5.22. *Let B be a finite connected left regular band and $\mathbb{k}$ a field. Then $\mathrm{gl. dim} \mathbb{k}B$ is bounded above by the $\mathbb{k}$ -Leray number $L_{\mathbb{k}}(\Delta(B))$ of the order complex of B .*

Proof. This is immediate from Corollary 5.21 as $\Delta(\partial e_Y B_{\geq X})$ is an induced subcomplex of $\Delta(B)$ for $X < Y$ in $\Lambda(B)$ and $Y = Be_Y$. $\square$

A graph Γ is *chordal* if it contains no induced cycle on four or more vertices. It is well known (see [75, Proposition 4.8] for a proof, but we claim no originality) that a simplicial complex has $\mathbb{k}$ -Leray number zero if and only if it is a simplex and has $\mathbb{k}$ -Leray number at most one if and only if it is the clique complex of a chordal graph.

A left regular band monoid is called *right hereditary* if its Hasse diagram is a rooted tree. For instance, the free left regular band and the left regular bands associated by Brown to matroids [34] and by Björner [23] to interval greedoids are right hereditary. Further examples, coming from Karnofsky-Rhodes expansions of semilattices, are given in [75]. In [75, Theorem 4.9], we verified that if B is a right hereditary left regular band, then $\Delta(B)$ is the clique complex of a chordal graph and hence has Leray number zero. From this and Corollary 5.22, we deduce the following result, which is [75, Theorem 4.9].

Corollary 5.23. *Let B be a right hereditary left regular band monoid. Then $\mathbb{k}B$ is hereditary for any field $\mathbb{k}$.*

Theorem 5.17 and Proposition 3.14 imply a tight connection between the homology of induced subcomplexes of clique complexes and the representation theory of free partially commutative left regular bands [75, Theorem 4.16].

Corollary 5.24. *Let $\Gamma = (V, E)$ be a finite simple graph, $\mathbb{k}$ a field and $B(\Gamma)$ the free partially commutative left regular band associated to Γ . Identifying $\Lambda(B(\Gamma))$ with the power set of V , we have*

$$\mathrm{Ext}_{\mathbb{k}B(\Gamma)}^n(\mathbb{k}_X, \mathbb{k}_Y) \cong \tilde{H}^{n-1}(\mathrm{Clique}(\Gamma[X \setminus Y]; \mathbb{k}))$$

for $X \supsetneq Y$ where $\Gamma[X \setminus Y]$ is the induced subgraph on the vertex set $X \setminus Y$.

Inspection of Corollary 5.24 shows that the $\mathbb{k}$ -Leray number the clique complex $\mathrm{Clique}(\Gamma)$ is exactly the global dimension of $\mathbb{k}B(\Gamma)$. This result first appeared as [75, Corollary 4.17] and gives an interpretation of the Leray number in terms of non-commutative algebra. The Leray number is usually given a commutative algebra interpretation as the Castelnuovo-Mumford regularity of the Stanley-Reisner ring of $\mathrm{Clique}(\Gamma)$.

Corollary 5.25. *Let Γ be a finite simple graph and $\mathbb{k}$ a field. Denoting by $B(\Gamma)$ the free partially commutative left regular band associated to Γ , we have that the global dimension of $\mathbb{k}B(\Gamma)$ is precisely the $\mathbb{k}$ -Leray number of the clique complex of Γ .*

Next we turn to the global dimension of connected CW left regular bands. This is a new result, which encompasses several results from [75] (where only monoids were considered).

Corollary 5.26. *Let B be a connected left regular band and $X \in \Lambda(B)$ such that $B_{\geq X}$ is a CW poset. (Note that $\Lambda(B)_{\geq X}$ is then a graded poset by*

Proposition 3.6 and Corollary 3.7.) Then, for $X, Y \in \Lambda(X)$, we have that

$$\dim_{\mathbb{k}} \operatorname{Ext}_{\mathbb{k}B}^q(\mathbb{k}_X, \mathbb{k}_Y) = \begin{cases} 1, & \text{if } X \leq Y \text{ and } \operatorname{rk}[X, Y] = q \\ 0, & \text{else.} \end{cases} \quad (5.4)$$

In particular, if B is a connected CW left regular band, then (5.4) holds for all $X, Y \in \Lambda(B)$ and consequently

$$\operatorname{gl. dim} \mathbb{k}B = \dim \Sigma(B)$$

holds.

Proof. Suppose that $X \leq Y$. As $\|\Delta(\partial e_Y B_{\geq X})\|$ is a sphere of dimension $\dim \Delta(e_Y B_{\geq X}) - 1 = \operatorname{rk}[X, Y] - 1$ (where $S^{-1} = \emptyset$), we deduce (5.4) from Theorem 5.17. Since the longest chain in $\Lambda(B)$ is of length $\dim \Sigma(B)$, the statement on global dimension for connected CW left regular bands is immediate from (5.4). $\square$

Corollary 5.26 covers the cases of face monoids of central hyperplane arrangements [75, 98] and face monoids of complex hyperplane arrangements [75]. The principal new examples in this paper are the semigroups of covectors of COMs including the face semigroup associated to a T -convex set of topes, the face semigroup of an affine hyperplane arrangement, the semigroup associated to an affine oriented matroid and the face semigroup of a CAT(0) cube complex. Let us record a few special cases of Corollary 5.26 in the following corollary.

Corollary 5.27. *Let $\mathbb{k}$ be a field.*

- (1) *If $(E, \mathcal{L})$ is a COM, then $\operatorname{gl. dim} \mathbb{k}\mathcal{L} = d$ where d is the dimension of the regular CW complex with face poset $\mathcal{L}$.*
- (2) *If $\mathcal{A}$ is an essential central hyperplane arrangement in $\mathbb{R}^d$ with face monoid $\mathcal{F}(\mathcal{A})$, then $\operatorname{gl. dim} \mathbb{k}\mathcal{F}(\mathcal{A}) = d$.*
- (3) *Suppose that $\mathcal{A}$ is an essential affine hyperplane arrangement in $\mathbb{R}^d$. Then $\operatorname{gl. dim} \mathbb{k}\mathcal{F}(\mathcal{A}) = d$.*
- (4) *If $\mathcal{A}$ is an essential complex hyperplane arrangement in $\mathbb{C}^d$, then $\operatorname{gl. dim} \mathbb{k}\mathcal{F}(\mathcal{A}) = 2d$.*
- (5) *If $(E, \mathcal{L})$ is an oriented matroid, then $\operatorname{gl. dim} \mathbb{k}\mathcal{L}$ is the rank of $(E, \mathcal{L})$.*
- (6) *If $(E, \mathcal{L}, g)$ is an affine oriented matroid with the rank of $(E, \mathcal{L})$ equal to $d + 1$, then $\operatorname{gl. dim} \mathbb{k}\mathcal{L}^+(g) = d$.*
- (7) *if K is a d -dimensional finite CAT(0) cube complex with face semigroup $\mathcal{F}(K)$, then $\operatorname{gl. dim} \mathbb{k}\mathcal{F}(K) = d$.*

5.3.1. Geometric left regular bands. We give in this subsection further semi-free actions of left regular bands on acyclic simplicial complexes. In [75] a left regular band B was defined to be *geometric* if the left stabilizer of each element of B is commutative. It then follows that the stabilizer of any subset of B is commutative and hence, if non-empty, has a minimum element. Note that if B embeds in the left regular band $\{0, +, -\}^n$ of covectors for

some $n \geq 0$, then B is geometric. Therefore, geometric left regular bands include free partially commutative left regular bands and COMs and strong eliminations systems such as central and affine hyperplane face semigroups, (affine) oriented matroids, face semigroups of T -convex sets of tope and face semigroups of CAT(0) cube complexes. The class of semigroups embeddable into $\{0, +, -\}^n$ is characterized in our paper [73]. The main examples in this paper that are not geometric left regular bands are face monoids of complex hyperplane arrangements and monoids of covectors of oriented interval greedoids.

Assume that B is a connected geometric left regular band and let $X \in \Lambda(B)$. Let Δ_X be the simplicial complex with vertex set the corresponding $\mathcal{L}$ -class L_X and where a subset $\tau \subseteq L_X$ forms a simplex if the elements of τ have a common upper bound. Because B is geometric τ , in fact, has a least upper bound in B . As L_X consists of the minimal elements of $B_{\geq X}$, it follows from the dual of Rota's cross-cut theorem (Theorem 3.1) that Δ_X is homotopy equivalent to $\Delta(B_{\geq X})$ and hence is acyclic by Theorem 5.5. Notice that if B is a monoid, then Δ_X is just the full simplex with vertex set L_X .

There is a natural semi-free action of $B_{\geq X} \curvearrowright \Delta_X$ by simplicial maps. Indeed, if $\tau = \{b_0, \dots, b_q\}$ has upper bound b and $a \in B_{\geq X}$, then $a\tau = \{ab_0, \dots, ab_q\}$ has upper bound ab . Also, if b is an upper bound for τ then $b \in \text{Stab}(\tau)$. Since the stabilizer of each simplex is a non-empty commutative left regular band, it has a minimum element and hence the action is semi-free. We have essentially established the following theorem.

Theorem 5.28. *Let B be a connected geometric left regular band and $X \in \Lambda(B)$. Let Δ_X be the simplicial complex constructed above. Then the augmented simplicial chain complex $C_\bullet(\Delta_X; \mathbb{k}) \rightarrow \mathbb{k}_X$ is a finite projective resolution by finitely generated $\mathbb{k}B$ -modules. Moreover, $C_0(\Delta_X; \mathbb{k}) \rightarrow \mathbb{k}_X$ is the projective cover of $\mathbb{k}_X$. In particular, the global dimension of B is bounded above by one less than the largest size of an $\mathcal{L}$ -class of B .*

Proof. Since $B_{\geq X} \curvearrowright \Delta_X$ is semi-free and Δ_X is acyclic, the first statement follows from Theorem 5.10. The second statement follows because $C_0(\Delta_X; \mathbb{k}) \cong \mathbb{k}L_X$. The final statement is a consequence of the inequality $\dim \Delta_X \leq |L_X| - 1$. $\square$

5.4. Minimal projective resolutions. Assume for this section that $\mathbb{k}$ is a field. Let A be a finite dimensional $\mathbb{k}$ -algebra and let M be a finite dimensional A -module. A projective resolution $P_\bullet \rightarrow M$ is said to be *minimal* if, for each $q > 0$, one has $d(P_q) \subseteq \text{rad}(P_{q-1})$ or, equivalently, $d: P_{q-1} \rightarrow d(P_{q-1})$ is a projective cover. Every finite dimensional module admits a minimal resolution, which is unique up to isomorphism (cf. Proposition 5.29 below). The following well-known proposition is stated in the context of group algebras in [39, Proposition 3.2.3], but the proof given there is valid for any finite dimensional algebra.

Proposition 5.29. *Let A be a finite dimensional $\mathbb{k}$ -algebra, M a finite dimensional A -module and $P_\bullet \rightarrow M$ a projective resolution. Then the following are equivalent.*

- (1) $P_\bullet \rightarrow M$ is the minimal resolution of M .
- (2) $\text{Hom}_A(P_q, S) \cong \text{Ext}_A^q(M, S)$ for any $q \geq 0$ and simple A -module S .
- (3) The coboundary map $d^*: \text{Hom}_A(P_q, S) \rightarrow \text{Hom}_A(P_{q+1}, S)$ is the zero map for all simple A -modules S and $q \geq 0$.
- (4) If $Q_\bullet \rightarrow M$ is a projective resolution of M , then the chain map $Q_\bullet \rightarrow P_\bullet$ lifting the identity map on M is surjective in each degree $q \geq 0$.
- (5) If $Q_\bullet \rightarrow M$ is a projective resolution of M , then the chain map $P_\bullet \rightarrow Q_\bullet$ lifting the identity map on M is injective in each degree $q \geq 0$.

The last two items explain why minimal resolutions are called “minimal.” The second item is a convenient way to test if a resolution is minimal. In the case that A is split over $\mathbb{k}$, it can be reformulated as saying that the number of occurrences of the projective cover of a simple module S as a direct summand in P_q is $\dim_{\mathbb{k}} \text{Ext}_A^q(M, S)$. Indeed, $\text{Hom}_A(P_q, S) = \text{Hom}_{A/\text{rad}(A)}(P_q/\text{rad}(P_q), S)$ and so the claim follows from Schur’s lemma because S is absolutely irreducible. We shall make use of this formulation of (2).

Theorem 5.30. *Let B be a connected left regular band, $X \in \Lambda(B)$ and $\mathbb{k}$ be a field. Suppose that $B_{\geq X}$ is a CW poset. Then the augmented cellular chain complex $C_\bullet(\Sigma(B_{\geq X}); \mathbb{k}) \rightarrow \mathbb{k}_X$ is the minimal projective resolution of $\mathbb{k}_X$.*

Proof. It follows from Corollary 5.13 that $C_\bullet(\Sigma(B_{\geq X}); \mathbb{k}) \rightarrow \mathbb{k}_X$ is a projective resolution and that the number of occurrences of the Schützenberger representation $\mathbb{k}L_Y$ associated to Y in $C_q(\Sigma(B_{\geq X}); \mathbb{k})$ is 1 if $Y \geq X$ and $\text{rk}[X, Y] = q$, and otherwise is 0. On the other hand, $\dim_{\mathbb{k}} \text{Ext}_{\mathbb{k}B}^q(\mathbb{k}_X, \mathbb{k}_Y) = 1$ if $Y \geq X$ and $\text{rk}[X, Y] = q$, and otherwise is 0 by Corollary 5.26. We conclude that $C_\bullet(\Sigma(B_{\geq X}); \mathbb{k}) \rightarrow \mathbb{k}_X$ is minimal by Proposition 5.29 $\square$

As a corollary, we obtain minimal projective resolutions for all the simple modules in the case of a connected CW left regular band.

Corollary 5.31. *Let B be a connected CW left regular band and let $\mathbb{k}$ be a field. Then the augmented cellular chain complex $C_\bullet(\Sigma(B_{\geq X}); \mathbb{k}) \rightarrow \mathbb{k}_X$ is the minimal projective resolution of X for all $X \in \Lambda(B)$.*

In particular, the corollary applies to the case where B is the face monoid of a real or complex hyperplane arrangement, the monoid of covectors of an oriented matroid or oriented interval greedoid, the face semigroup of an affine hyperplane arrangement or CAT(0) cube complex, the semigroup of covectors associated to an affine oriented matroid or, more generally, the semigroup of covectors of a COM.

6. QUIVER PRESENTATIONS

Let us fix for the discussion a finite connected left regular band B and a field $\mathbb{k}$. In this section we will compute a quiver presentation for $\mathbb{k}B$ under some very strong hypotheses on B that will be satisfied by connected CW left regular bands.

6.1. A general result. Recall that the quiver of $\mathbb{k}B$ is denoted $Q(\mathbb{k}B)$. It has vertex set $\Lambda(B)$. If $X < Y$, then the number of directed edges from X to Y is one less than the number of connected components of $\partial e_Y B_{\geq X}$ by Theorem 5.17; there are no other arrows. In particular, $Q(\mathbb{k}B)$ is acyclic and is independent of $\mathbb{k}$. Fix a complete set of primitive orthogonal idempotents $\{\eta_X \mid X \in \Lambda(B)\}$ constructed as per Section 4.3.

Let us assume the following four properties of B .

- (B1) $Q(\mathbb{k}B)$ has no multiple edges.
- (B2) $Q(\mathbb{k}B)$ is graded in the sense that all paths between two given vertices $X < Y$ have the same length. Set $\text{rk}[X, Y]$ to be the common length of all paths from X to Y .
- (B3)

$$\dim_{\mathbb{k}} \text{Ext}_{\mathbb{k}B}^2(\mathbb{k}_X, \mathbb{k}_Y) = \begin{cases} 1, & \text{if } \text{rk}[X, Y] = 2 \\ 0, & \text{else.} \end{cases}$$

- (B4) There is a $\mathbb{k}B$ -module homomorphism $\partial: \mathbb{k}B \rightarrow \mathbb{k}B$ such that:
 - (a) $\partial^2 = 0$;
 - (b) $\partial(\eta_X)\eta_X = 0$, for all $X \in \Lambda(B)$;
 - (c) $\partial(\eta_Y)\eta_X \neq 0$ if $X \rightarrow Y$ is an arrow of $Q(\mathbb{k}B)$.

We remark that under these assumptions, $Q(\mathbb{k}B)$ is the Hasse diagram of a partial order $\preceq$ on $\Lambda(B)$ that is coarser than the original ordering. More precisely, we put $X \preceq Y$ if there is a path from X to Y in $Q(\mathbb{k}B)$ and note that $X \preceq Y$ implies $X \leq Y$ and that $(\Lambda(B), \preceq)$ is a graded poset. As usual, let us write $X \prec Y$ if $X \preceq Y$ and $X \neq Y$.

Theorem 6.1. *Suppose that B is a connected left regular band satisfying (B1)–(B4). Then $\mathbb{k}B$ has a quiver presentation $(Q(\mathbb{k}B), I)$ where I is given by the following minimal system of relations R . For each $X < Y$ with $\text{rk}[X, Y] = 2$, we have a relation*

$$r_{X,Y} = \sum_{X \prec Z \prec Y} (X \rightarrow Z \rightarrow Y)$$

in R . In particular, I is a homogeneous ideal and $\mathbb{k}B \cong \mathbb{k}Q(\mathbb{k}B)/I$ is a quadratic algebra.

Proof. Set $Q = Q(\mathbb{k}B)$ and suppose that there is an edge $X \rightarrow Y$ in Q . Then, by (B2), there are no paths of length greater than one in $Q(\mathbb{k}B)$ from

X to Y and so $\eta_Y \text{rad}^2(\mathbb{k}B)\eta_X = 0$. Therefore, we have by (B1) that

$$\begin{aligned} 1 &= \dim_{\mathbb{k}} \text{Ext}_{\mathbb{k}B}^1(\mathbb{k}_X, \mathbb{k}_Y) = \dim_{\mathbb{k}} \eta_Y [\text{rad}(\mathbb{k}B) / \text{rad}^2(\mathbb{k}B)] \eta_X \\ &= \dim_{\mathbb{k}} \eta_Y \mathbb{k}B \eta_X. \end{aligned}$$

Observe that, by (B4), $0 \neq \partial(\eta_Y)\eta_X = \eta_Y \partial(\eta_Y)\eta_X$ and so $\{\partial(\eta_Y)\eta_X\}$ is a basis for $\eta_Y \mathbb{k}B \eta_X \cong \eta_Y [\text{rad}(\mathbb{k}B) / \text{rad}^2(\mathbb{k}B)] \eta_X$. Thus we have a surjective homomorphism $\varphi: \mathbb{k}Q \rightarrow \mathbb{k}B$ given by $\varphi(\varepsilon_X) = \eta_X$ and $\varphi(X \rightarrow Y) = \partial(\eta_Y)\eta_X$ with $\ker \varphi$ admissible by the general theory of split basic finite dimensional algebras [6, 15].

Next we show that $r_{X,Y} \in \ker \varphi$ if $\text{rk}[X, Y] = 2$. Note that

$$\varphi(r_{X,Y}) = \sum_{X \prec Z \prec Y} \partial(\eta_Y)\eta_Z \partial(\eta_Z)\eta_X = \sum_{X \prec Z \prec Y} \eta_Y \partial(\eta_Y)\eta_Z \partial(\eta_Z)\eta_X. \quad (6.1)$$

Notice that if $Z \in \Lambda(B)$ and $X \prec Z \prec Y$ is false, then

$$\eta_Y \partial(\eta_Y)\eta_Z \partial(\eta_Z)\eta_X = 0.$$

Indeed, if $Z \notin \{X, Y\}$, then there is no path in Q from X to Y through Z by definition of $\preceq$ and so $\eta_Y \mathbb{k}B \eta_Z \mathbb{k}B \eta_X = 0$. If $Z \in \{X, Y\}$ we use that $\partial(\eta_X)\eta_X = 0 = \partial(\eta_Y)\eta_Y$ by (B4). Thus the right hand side of (6.1) is equal to

$$\begin{aligned} \sum_{Z \in \Lambda(B)} \eta_Y \partial(\eta_Y)\eta_Z \partial(\eta_Z)\eta_X &= \sum_{Z \in \Lambda(B)} \partial[\partial(\eta_Y)\eta_Z]\eta_X \\ &= \partial \left[\partial(\eta_Y) \cdot \sum_{Z \in \Lambda(B)} \eta_Z \right] \eta_X \\ &= \partial^2(\eta_Y)\eta_X \\ &= 0 \end{aligned}$$

using that $\partial^2 = 0$ and that $\sum_{Z \in \Lambda(B)} \eta_Z = 1$. This proves that $\varphi(r_{X,Y}) = 0$.

To complete the proof, let R' be a minimal system of relations for $I = \ker \varphi$. We claim that

$$R' = \{c_{X,Y} \cdot r_{X,Y} \mid \text{rk}[X, Y] = 2\}$$

where each $c_{X,Y} \in \mathbb{k} \setminus \{0\}$. It will then follow that the $r_{X,Y}$ also form a minimal system of relations for I .

By Theorem 4.1 and (B3) we have that $R' = \{s_{X,Y} \mid \text{rk}[X, Y] = 2\}$ where $s_{X,Y}$ is a non-zero linear combination of paths from X to Y in Q , necessarily of length 2 by (B2). It follows that if $\text{rk}[X, Y] = 2$, then

$$r_{X,Y} = \sum_{\text{rk}[U,V]=2} p_{U,V} s_{U,V} q_{U,V}$$

where $p_{U,V} \in \varepsilon_Y \mathbb{k}Q \varepsilon_V$ and $q_{U,V} \in \varepsilon_U \mathbb{k}Q \varepsilon_X$. But since each path in $r_{X,Y}$ has length 2 and each path in $s_{U,V}$ has length 2, we conclude that $p_{U,V} s_{U,V} q_{U,V} = 0$ if $(U, V) \neq (X, Y)$ and that $r_{X,Y} = k_{X,Y} \cdot s_{X,Y}$ for some $k_{X,Y} \in \mathbb{k} \setminus \{0\}$. This establishes the claim and completes the proof. $\square$

6.2. A quiver presentation for CW left regular bands. One of the main results of this text is the next theorem, which computes a quiver presentation of the algebra of a connected CW left regular band, generalizing a result of the second author for face monoids of central hyperplane arrangements [98].

Theorem 6.2. *Suppose that B is a connected CW left regular band.*

- (a) *The quiver $Q = Q(\mathbb{k}B)$ of $\mathbb{k}B$ is the Hasse diagram of $\Lambda(B)$.*
- (b) *$\Lambda(B)$ is graded.*
- (c) *$\mathbb{k}B$ has a quiver presentation (Q, I) where I has minimal system of relations*

$$r_{X,Y} = \sum_{X < Z < Y} (X \longrightarrow Z \longrightarrow Y)$$

ranging over all rank 2 intervals $[X, Y]$ in $\Lambda(B)$.

Proof. The fact that $\Lambda(B)$ is graded is immediate from Proposition 3.6 and Corollary 3.7 because B is graded, being a CW poset.

By taking $q = 1$ in (5.4) from Corollary 5.26, we see that there is an edge $X \longrightarrow Y$ in Q if and only if $X < Y$ is a cover relation in $\Lambda(B)$ and that there are no multiple edges. This establishes (a).

To prove the remainder of the theorem, it suffices to show that Theorem 6.1 applies to B . It is immediate from the previous discussion that (B1)–(B2) are satisfied. Property (B3) is a consequence of (5.4) from Corollary 5.26. So it remains to verify that (B4) holds.

Choose for each $X \in \Lambda(B)$ an idempotent e_X with $Be_X = X$ and let $\{\eta_X \mid X \in \Lambda(B)\}$ be the complete set of orthogonal primitive idempotents for $\mathbb{k}B$ from Section 4.3. Let, as usual, $\mathbb{k}L_X$ denote the Schützenberger representation associated to the $\mathcal{L}$ -class of e_X . Then as a left $\mathbb{k}B$ -module we have

$$\mathbb{k}B \cong \bigoplus_{X \in \Lambda(B)} \mathbb{k}L_X \cong \bigoplus_{q \geq 0} C_q(\Sigma(B); \mathbb{k})$$

where the last isomorphism follows from (5.3) of Corollary 5.13 with $X = \hat{0}$. Thus we can define a $\mathbb{k}$ -module homomorphism $\partial: \mathbb{k}B \longrightarrow \mathbb{k}B$ by transporting the differential

$$d: \bigoplus_{q \geq 0} C_q(\Sigma(B); \mathbb{k}) \longrightarrow \bigoplus_{q \geq 0} C_q(\Sigma(B); \mathbb{k})$$

under the isomorphism between $\mathbb{k}B$ and $\bigoplus_{q \geq 0} C_q(\Sigma(B); \mathbb{k})$. Clearly, we have $\partial^2 = 0$.

Let us make the definition of ∂ more explicit. We recall from Corollary 4.4 that $\{b\eta_X \mid \sigma(b) = X\}$ is a basis for $\mathbb{k}B$ and that $\mathbb{k}L_X \cong \mathbb{k}B\eta_X$. Identifying q -cells of B with basis elements of $C_q(\Sigma(B); \mathbb{k})$, we have

$$d(b) = \sum [b : a]a$$

where the sum runs over all maximal proper faces a of b and the $[b : a] = \pm 1$ are the incidence numbers associated to the boundary maps of $C_\bullet(\Sigma(B); \mathbb{k})$ with respect to the orientation chosen in the proof of Theorem 5.10 (see also Remark 5.14). See [77, Chapter IX, §7] for details on incidence numbers for regular cell complexes. We then have, for $b \in B$, that

$$\partial(b\eta_{\sigma(b)}) = \begin{cases} \sum [b : a]a\eta_{\sigma(a)}, & \text{if } b \in L \\ 0, & \text{else} \end{cases}$$

where the sum runs over all maximal proper faces a of b .

To complete the proof, first note that

$$\partial(\eta_X)\eta_X = \partial(e_X\eta_X)\eta_X = \sum [e_X : a]a\eta_{\sigma(a)}\eta_X = 0$$

because each maximal proper face a of e_X satisfies $\sigma(a) < X$ and $b\eta_X = 0$ if $\sigma(b) \not\geq X$ by Theorem 4.2. On the other hand, if Y covers X in $\Lambda(B)$, then $\partial e_Y B_{\geq X} \cong S^0$ because $B_{\geq X}$ is a CW poset and e_Y is a 1-cell of $B_{\geq X}$. Therefore, e_Y has exactly two maximal proper faces a_1, a_2 with support X . Thus

$$\begin{aligned} \partial(\eta_Y)\eta_X &= \partial(e_Y\eta_Y)\eta_X = \sum [e_Y : a]a\eta_{\sigma(a)}\eta_X \\ &= [e_Y : a_1]a_1\eta_X + [e_Y : a_2]a_2\eta_X \neq 0. \end{aligned}$$

An application of Theorem 6.1 completes the proof of the theorem. $\square$

Example 6.3 (Ladders). Let L_n be the ladder left regular band from Example 3.17. Theorem 6.2 shows that the quiver of $\mathbb{k}L_n$ is a directed path with $n+1$ vertices (that is, the Dynkin quiver A_{n+1} oriented as a directed path and that the relations declare all paths of length two to be zero). In other words, $\mathbb{k}L_n$ is the quadratic monomial algebra $\mathbb{k}A_{n+1}/\text{rad}^2(\mathbb{k}A_{n+1})$.

We shall later use Theorem 6.2 to show that the algebra of the face semigroup of a CAT(0) cube complex is isomorphic to the incidence algebra of its intersection semilattice.

An interesting question is whether $\mathbb{Z}B$ is isomorphic to the quotient of the integral path ring of the Hasse diagram of $\Lambda(B)$ modulo the ideal generated by the sum of all paths of length 2 when B is a connected CW left regular band.

7. QUADRATIC AND KOSZUL DUALS

In this section we consider quadratic and Koszul duals for connected CW left regular band algebras.

7.1. Quadratic duals. Let P be a graded poset with Hasse diagram Q . Let I of $\mathbb{k}Q$ be the ideal generated by the elements

$$r_{\sigma, \tau} = \sum_{\sigma < \gamma < \tau} (\sigma \longrightarrow \gamma \longrightarrow \tau), \quad \text{with } \text{rk}[\sigma, \tau] = 2. \quad (7.1)$$

Motivated by Theorem 6.1, we are led to consider the following question: what is the quadratic dual of $A = \mathbb{k}Q/I$?

In our examples the poset P will also have an additional property termed strong connectedness in [78].

A simplicial complex K is called a *chamber complex* if it is pure and any two facets can be connected by a gallery. Recall that if C, C' are facets, then a *gallery* from C to C' is a sequence $C = C_0, \dots, C_n = C'$ of facets such that C_i, C_{i+1} are distinct and adjacent, where C_i, C_{i+1} are *adjacent* if they have a common codimension one face. The reader is referred to [1, Appendix A.1] for details on chamber complexes. Crucial for us is the fact that every connected triangulated manifold without boundary is a chamber complex (cf. [1, Exercise A.17]).

Notice that if P is a graded poset with $\Delta(P)$ a pure simplicial complex, then two facets $\sigma_0 < \dots < \sigma_n$ and $\tau_0 < \dots < \tau_n$ of $\Delta(P)$ are adjacent if and only if there exists $0 \leq i \leq n$ with $\sigma_j = \tau_j$ for $j \neq i$ and $\sigma_i \neq \tau_i$; see Figure 21.




FIGURE 21. Adjacent facets in an order complex

Lemma 7.1. *Let $f: P \rightarrow Q$ be a surjective cellular map of posets such that $\sigma < \sigma'$ implies $f(\sigma) < f(\sigma')$ and suppose that $\Delta(P)$ is a chamber complex. Then $\Delta(Q)$ is a chamber complex of the same dimension.*

Proof. Corollary 3.7 and its proof imply that $\Delta(Q)$ is pure, $f: \Delta(P) \rightarrow \Delta(Q)$ preserves dimensions of simplices and f maps the set of facets of $\Delta(P)$ onto the set of facets of $\Delta(Q)$. Moreover, given a pair of adjacent facets of $\Delta(P)$, either f collapses them or takes them to adjacent facets. The lemma follows easily from these observations. $\square$

The following definition is from [78].

Definition 7.2 (Strongly connected). Let P be a graded poset. Then P is said to be *strongly connected* if $\Delta([\sigma, \tau])$ is a chamber complex for each closed interval $[\sigma, \tau]$ of P .

For example, any semimodular lattice is strongly connected (see the proof of [98, Proposition 9.6]). Notice that P is strongly connected if and only if P^{op} is strongly connected.

Remark 7.3. Observe that if K is a chamber complex, then a cone $K * x$ on K with cone point x is also a chamber complex. Indeed, the map $F \mapsto F \cup \{x\}$

gives a bijection between facets of K and facets of $K * x$ which raises the dimension by 1. Moreover, this bijection preserves adjacency.

The reader should recall the definition of the *incidence algebra* $I(P; \mathbb{k})$ of a poset P from Subsection 4.3.3. The following lemma provides a quiver presentation for the incidence algebra of a strongly connected graded poset.

Lemma 7.4. *Let P be a strongly connected graded poset with Hasse diagram Q and let $\mathbb{k}$ be a commutative ring with unit. Then there is an isomorphism $I(P; \mathbb{k}) \cong \mathbb{k}Q/I$ where I is generated by all differences $p - q$ of parallel paths p, q in Q of length 2.*

Proof. First note that $I(P; \mathbb{k}) \cong \mathbb{k}Q/I'$ where I' is spanned by all differences $p - q$ of parallel paths in Q . Thus $I \subseteq I'$. We claim that $I = I'$.

Notice that paths from σ to τ in Q correspond to facets of $\Delta([\sigma, \tau])$. If $p, q: \sigma \rightarrow \tau$ are two parallel paths in Q corresponding to distinct adjacent facets in $\Delta([\sigma, \tau])$, then we can write $p = uav$ and $q = ubv$ where a, b are parallel paths of length 2; see Figure 22. Thus $p - q = u(a - b)v \in I$.




FIGURE 22. Parallel paths in Q corresponding to adjacent facets

Now suppose that $p, q: \sigma \rightarrow \tau$ are arbitrary parallel paths in Q . Then since $\Delta([\sigma, \tau])$ is a chamber complex we can find a sequence $p = p_0, \dots, p_n = q$ of paths from σ to τ such that p_i and p_{i+1} represent distinct adjacent facets of $\Delta([\sigma, \tau])$. Then $p - q = (p_0 - p_1) + (p_1 - p_2) + \dots + (p_{n-1} - p_n) \in I$. It follows that $I = I'$, as required. This completes the proof. $\square$

The following generalizes [98, Proposition 9.6] from the case of central hyperplane arrangements.

Theorem 7.5. *Let P be a strongly connected graded poset. Let Q be the Hasse diagram of P and let $A = \mathbb{k}Q/I$ where I is the ideal generated by the system of relations (7.1). Then the quadratic dual $A^\perp$ is isomorphic to the incidence algebra $I(P^{op}; \mathbb{k})$ of the opposite poset of P .*

Proof. Clearly $I_2^\perp$ is spanned by all differences $p - q$ of parallel paths of length 2 in Q^{op} . As P^{op} is a strongly connected graded poset, the result follows from Lemma 7.4. $\square$

In order to apply Theorem 7.5 to connected CW left regular bands, we need that $\Lambda(B)$ is a strongly connected graded poset.




FIGURE 23. A rank 2 interval in a thin poset

Lemma 7.6. *Let B be a connected CW left regular band. Then $\Lambda(B)$ is a strongly connected graded poset.*

Proof. Theorem 6.2 establishes that $\Lambda(B)$ is graded. Let $[X, Y]$ be a closed interval of $\Lambda(B)$. Choose an idempotent e_Y with $Be_Y = Y$. Then $[X, Y]$ is the support lattice of $e_Y B_{\geq X}$. Note that $\Delta(e_Y B_{\geq X})$ is the cone

$$\Delta(\partial e_Y B_{\geq X}) * e_Y$$

and that $\Delta(\partial e_Y B_{\geq X})$ is a chamber complex, being a triangulation of a sphere. Thus $\Delta(e_Y B_{\geq X})$ is a chamber complex by Remark 7.3. We conclude using Proposition 3.6 and Lemma 7.1 that $\Delta([X, Y])$ is a chamber complex and hence $\Lambda(B)$ is strongly connected. $\square$

Theorem 7.7. *Let B be a connected CW left regular band (and so in particular $\mathbb{k}B$ is a quadratic algebra with the quiver presentation from Theorem 6.2). Then the quadratic dual $\mathbb{k}B^!$ is isomorphic to $I(\Lambda(B)^{op}; \mathbb{k})$.*

Proof. The theorem is an immediate consequence of Theorem 6.2, Theorem 7.5 and Lemma 7.6. $\square$

We next aim to give a necessary condition for the algebra of a connected CW left regular band to be an incidence algebra. Consequently, most of our CW left regular bands are not isomorphic to incidence algebras. We will also show that if K is a CAT(0) cube complex, then the algebra of its face semigroup $\mathcal{F}(K)$ is isomorphic to the incidence algebra of its intersection semilattice $\mathcal{L}(K)$. The same is true, more generally, for any lopsided system.

A graded poset is said to be *thin* (cf. [26, Section 4.7]) if each interval $[x, y]$ of rank 2 has cardinality 4; see Figure 23. For example, if X is a simplicial complex and P is the face poset of X with an adjoined empty face, then P is thin. This is because each subposet $P_{\leq p}$ is a boolean lattice and boolean lattices are thin.

Theorem 7.8. *Let B be a connected CW left regular band and $\mathbb{k}$ a field. If $\mathbb{k}B$ is isomorphic to the incidence algebra of a poset, then $\Lambda(B)$ is thin.*

Proof. Let Q be the Hasse diagram of $\Lambda(B)$. If P is a poset, then the quiver of the incidence algebra of P is the Hasse diagram of P . Since Q is the quiver of $\mathbb{k}B$ by Theorem 6.2, we conclude that if $\mathbb{k}B$ is isomorphic to an incidence algebra, then it is isomorphic to the incidence algebra of $\Lambda(B)$. Let us assume that this is indeed the case. We show that $\Lambda(B)$ is thin.

Let $[X, Y]$ be an interval of rank 2 in $\Lambda(B)$. Let $\{\eta_X \mid X \in \Lambda(B)\}$ be a complete set of orthogonal primitive idempotent constructed as per Section 4.3 and let I be the admissible ideal from Theorem 6.2. Then $\dim \eta_Y \mathbb{k} B \eta_X = 1$ because $I(\Lambda(B); \mathbb{k})$ is the quotient of $\mathbb{k}Q$ by the ideal I' generated by differences of parallel paths and so $\dim \varepsilon_Y [\mathbb{k}Q/I'] \varepsilon_X = 1$. Thus $\varepsilon_Y I \varepsilon_X$ has codimension 1 in $\varepsilon_Y \mathbb{k}Q \varepsilon_X$. On the other hand, we claim $\dim \varepsilon_Y I \varepsilon_X = 1$. Indeed, retaining the notation of Theorem 6.2, if $r \in \varepsilon_Y I \varepsilon_X$, then

$$r = \sum_{\text{rk}[U,V]=2} p_{U,V} r_{U,V} q_{U,V}$$

where $p_{U,V} \in \varepsilon_Y \mathbb{k}Q \varepsilon_V$ and $q_{U,V} \in \varepsilon_U \mathbb{k}Q \varepsilon_X$. But since each path in r has length 2 and each path in $r_{U,V}$ has length 2, it follows that $p_{U,V} r_{U,V} q_{U,V} = 0$ if $(U, V) \neq (X, Y)$ and that $r = k_{X,Y} \cdot r_{X,Y}$ for some $k_{X,Y} \in \mathbb{k}$. Thus $\varepsilon_Y I \varepsilon_X = \mathbb{k} r_{X,Y}$. We conclude that $\dim \varepsilon_Y \mathbb{k}Q \varepsilon_X = 1 + \dim \varepsilon_Y I \varepsilon_X = 2$ and hence $[X, Y]$ consists of four elements. Thus $\Lambda(B)$ is thin. $\square$

Remark 7.9. The intersection lattice of the braid arrangement in $\mathbb{R}^n$ for $n \geq 3$, or of any central line arrangement in $\mathbb{R}^2$ with at least 3 lines, is not thin and so their algebras are not incidence algebras.

Corollary 7.10. *Let K be a finite $\text{CAT}(0)$ cube complex and let $\mathbb{k}$ be a field. Let $\mathcal{F}(K)$ be the face semigroup of K and $\mathcal{L}(K)$ the intersection semilattice of K . Then $\mathbb{k}\mathcal{F}(K) \cong I(\mathcal{L}(K); \mathbb{k})$.*

Proof. The semigroup $\mathcal{F}(K)$ is a right ideal in L^n where n is the number of hyperplanes of K . The support semilattice $\Lambda(\mathcal{F}(K)) \cong \mathcal{L}(K)$ is then an ideal in $\Lambda(L^n)$, which in turn is congruent to the boolean lattice $P(\mathcal{H})$ where $\mathcal{H}$ is the set of hyperplanes of K . If we fix e_X with $L^n e_X = X$ for $X \in \Lambda(L^n)$ such that $e_X \in \mathcal{F}(K)$ whenever $X \in \Lambda(\mathcal{F}(K))$ (viewed as an ideal of $\Lambda(L^n)$), then using that $\mathcal{F}(K)$ is a right ideal we conclude that the primitive idempotent η_X , constructed as per Section 4.3, belongs to $\mathbb{k}\mathcal{F}(K)$ whenever $X \in \Lambda(\mathcal{F}(K))$.

We conclude that if $f = \sum_{X \in \Lambda(\mathcal{F}(K))} \eta_X$, then $\mathbb{k}\mathcal{F}(K) = f \mathbb{k} L^n f$. If A is an incidence algebra and f is a sum of a subset of a complete set of orthogonal primitive idempotents, then $f A f$ is the incidence algebra of the subposet corresponding to all elements of the poset whose corresponding primitive idempotent appears in f . Thus it suffices to show that $\mathbb{k} L^n$ is isomorphic to an incidence algebra.

But $\mathbb{k}[M \times N] \cong \mathbb{k}M \otimes \mathbb{k}N$ for any monoids M, N and the class of incidence algebras of posets is closed under tensor product because $I(P; \mathbb{k}) \otimes I(P'; \mathbb{k}) \cong I(P \times P'; \mathbb{k})$. So it suffices to show that $\mathbb{k}L$ is the incidence algebra of a poset. But the quiver of L is $\hat{0} \rightarrow \hat{1}$ and hence $\mathbb{k}L$ is the incidence algebra of the two element chain. The result follows. $\square$

Remark 7.11. The proof of Corollary 7.10 applies to any right ideal in L^n that is a connected left regular band and, in particular, for lopsided systems.

7.2. Koszul duals. Let B be a connected CW left regular band. Our goal in this section is to prove that $\mathbb{k}B$ is a Koszul algebra. The Koszul dual is then $I(\Lambda(B)^{op}; \mathbb{k})$ and $\text{Ext}(\mathbb{k}B) \cong I(\Lambda(B); \mathbb{k})$. We follow closely [98, Section 9].

Fix a complete set of orthogonal primitive idempotents $\{\eta_X \mid X \in \Lambda(B)\}$ constructed as per Section 4.3. Then the grading of $\mathbb{k}B$ coming from the quiver presentation in Theorem 6.2 can be defined ‘intrinsically’ by

$$\mathbb{k}B = \bigoplus_{n \geq 0} \left[\bigoplus_{\text{rk}[X,Y]=n} \eta_Y \mathbb{k}B \eta_X \right]$$

(or one can just check directly that this is a grading with $(\mathbb{k}B)_0 \cong \mathbb{k}^{\Lambda(B)}$).

Theorem 7.12. *Let B be a connected CW left regular band. Then $\mathbb{k}B$ is a Koszul algebra and the Koszul dual $\mathbb{k}B^!$ is isomorphic to $I(\Lambda(L)^{op}; \mathbb{k})$. Hence $\text{Ext}(\mathbb{k}B) \cong I(\Lambda(L); \mathbb{k})$.*

Proof. To prove that $\mathbb{k}B$ is Koszul with respect to the above grading it suffices to show that each simple module $\mathbb{k}_X$ (viewed as a graded module concentrated in degree 0) has a linear resolution.

The augmented cellular chain complex $C_\bullet(\Sigma(L_{\geq X}); \mathbb{k}) \rightarrow \mathbb{k}_X$ is the minimal projective resolution by Theorem 5.30. We need to show that it is a linear resolution. As usual let $\mathbb{k}L_Y$ be the Schützenberger representation associated to $Y \in \Lambda(B)$. Then

$$C_q(\Sigma(B_{\geq X}); \mathbb{k}) \cong \bigoplus_{\text{rk}[X,Y]=q} \mathbb{k}L_Y \cong \bigoplus_{\text{rk}[X,Y]=q} \mathbb{k}B\eta_Y \quad (7.2)$$

by Corollary 5.13. As before, we identify $C_q(\Sigma(B_{\geq X}); \mathbb{k})$ with

$$\bigoplus_{\text{rk}[X,Y]=q} \mathbb{k}B\eta_Y$$

via the isomorphism sending an oriented cell $[b]$ (as per the proof of Theorem 5.10) to the element $b\eta_{\sigma(b)}$.

Next we observe that if $\text{rk}[X, Y] = q$, then we can define a grading

$$\mathbb{k}B\eta_Y = \bigoplus_{i \geq 0} \left[\bigoplus_{\text{rk}[X,W]=i} \eta_W \mathbb{k}B\eta_Y \right].$$

This of course puts a grading on the right hand side of (7.2). If $W \not\geq Y$, and in particular if $\text{rk}[X, W] < q$, then $\eta_W \mathbb{k}B\eta_Y = 0$. Also the degree q component of $\mathbb{k}B\eta_Y$ is $\eta_Y \mathbb{k}B\eta_Y = \mathbb{k}\eta_Y$. Thus $\mathbb{k}B\eta_Y$ is generated in degree q and hence so is the right hand side of (7.2). It remains to show that d is a degree 0 map. Indeed, we have

$$d(\eta_W a \eta_Y) = \eta_W d(a \eta_Y) \in \bigoplus_{\text{rk}[X,Y']=q-1} \eta_W \mathbb{k}B\eta_{Y'}$$

and so d has degree 0. We may now conclude that $C_\bullet(\Sigma(B_{\geq X}); \mathbb{k}) \longrightarrow \mathbb{k}_X$ is a linear resolution. $\square$

In particular, this result applies to the case where B is the face monoid of a real or complex hyperplane arrangement, the monoid of covectors of an oriented matroid or oriented interval greedoid, the face semigroup of an affine hyperplane arrangement or CAT(0) cube complex or the semigroup of covectors of an affine oriented matroid or, more generally, a COM.

Example 7.13 (Ladders). This is a continuation of Examples 3.17 and 6.3. Theorem 7.12 implies that $\mathbb{k}L_n$ is a Koszul algebra whose dual is the incidence algebra of a chain of $n + 1$ elements. But this incidence algebra is precisely the path algebra of the Dynkin quiver A_{n+1} oriented as a directed path. Of course, it is well known that the Koszul dual of the path algebra of the Dynkin quiver A_{n+1} oriented as a directed path is the monomial algebra obtained by factoring out the paths of length two.

Polo and Woodcock [87, 119] independently showed that the incidence algebra $I(P; \mathbb{k})$ of a graded poset P is a Koszul algebra with respect to the natural grading if and only if each open interval in P is $\mathbb{k}$ -Cohen-Macaulay. We thus obtain the following theorem, which will be applied to enumerate cells of connected CW left regular bands.

Theorem 7.14. *Let B be a connected CW left regular band. Then each open interval of $\Lambda(B)$ is a Cohen-Macaulay poset. In particular, if B is a monoid, then the proper part $\Lambda(B) \setminus \{\widehat{0}, \widehat{1}\}$ of $\Lambda(B)$ is Cohen-Macaulay.*

Proof. By the theorem of Polo and Woodcock [87, 119] and Theorem 7.12, we have that each open interval of $\Lambda(B)$ is $\mathbb{k}$ -Cohen-Macaulay for every field $\mathbb{k}$ and hence is Cohen-Macaulay by the universal coefficient theorem. $\square$

8. INJECTIVE ENVELOPES FOR HYPERPLANE ARRANGEMENTS, ORIENTED MATROIDS, CAT(0) CUBE COMPLEXES AND COMS

In this section we give a geometric construction of the injective envelopes of the simple modules for $\mathbb{k}B$ when B has the property that bB is isomorphic to the face monoid of a hyperplane arrangement or, more generally, to the monoid of covectors of an oriented matroid for all $b \in B$. This includes left regular bands associated to central and affine hyperplane arrangements, (possibly affine) oriented matroids, CAT(0) cube complexes and, more generally, COMs.

As the injective envelope of a simple left module is the vector space dual of the right projective cover of the corresponding simple right module, we proceed by constructing the right projective cover of each simple module. We first do this for hyperplane arrangements and then we give the more general proof for oriented matroids. Although this has some redundancy, there may be readers who are more familiar with hyperplane arrangements than oriented matroids and so we would like that case to be self-contained.

Let us fix a field $\mathbb{k}$ for this section. We begin with some generalities on right modules for left regular bands. If B is a left regular band and $e \in B$, then we define $\varphi_e: B \rightarrow eB$ by $\varphi_e(b) = eb$. Recall that $eB = eBe$ is a left regular band monoid.

Proposition 8.1. *The mapping $\varphi_e: B \rightarrow eB$ is a surjective homomorphism.*

Proof. We have $\varphi_e(ab) = eab = eae b = \varphi_e(a)\varphi_e(b)$ by the left regular band axioms. Clearly, φ is surjective. $\square$

Any right $\mathbb{k}eB$ -module can be lifted to a $\mathbb{k}B$ -module via φ_e . This applies in particular to $\mathbb{k}eB$ itself.

Proposition 8.2. *Let B be a left regular band and $e \in B$. There is an isomorphism of right $\mathbb{k}B$ -modules $e\mathbb{k}B \rightarrow \mathbb{k}eB$ induced by the identity map on eB where we give $\mathbb{k}eB$ the right $\mathbb{k}B$ -module structure induced via φ_e and we view $e\mathbb{k}B$ as a right ideal of $\mathbb{k}B$.*

Proof. If $a \in eB$ and $b \in B$, then $ab = eab = eae b = aeb = a\varphi_e(b)$ and the result follows. $\square$

An immediate consequence of Proposition 8.2 is the following corollary.

Corollary 8.3. *Let B be a connected left regular band and $X \in \Lambda(B)$ with $X = Be_X$. Then the simple right $\mathbb{k}B$ -module $\mathbb{k}_X$ is obtained by lifting $\mathbb{k}e_X B / \mathbb{k}[\partial e_X B]$ from $\mathbb{k}e_X B$ via φ_e . Moreover, if P_X is a projective indecomposable right $\mathbb{k}e_X B$ -module with $P_X / \text{rad}(P_X) \cong \mathbb{k}e_X B / \mathbb{k}[\partial e_X B]$, then P_X is a direct summand in $e_X \mathbb{k}B$ as a right $\mathbb{k}B$ -module and is the right projective cover of $\mathbb{k}_X$ as a $\mathbb{k}B$ -module.*

Thus, for constructing right projective covers of simple modules, we are reduced to considering left regular band monoids B and $\mathbb{k}_{\hat{1}}$ where $\hat{1} = B$ is the maximum element of the lattice $\Lambda(B)$. The next lemma provides necessary and sufficient conditions for a quotient by a right ideal to give a right projective cover of $\mathbb{k}_{\hat{1}}$.

Lemma 8.4. *Let B be a left regular band monoid and denote by $\hat{1}$ the maximum of $\Lambda(B)$. Suppose that $R \subsetneq B$ is a proper right ideal. Then the natural homomorphism $\psi: \mathbb{k}B / \mathbb{k}R \rightarrow \mathbb{k}_{\hat{1}}$ induced by $1 \mapsto 1$ and $b \mapsto 0$ for $b \neq 1$ is a right projective cover if and only if*

- (1) $BR = \partial B$;
- (2) R is a connected left regular band.

Proof. Suppose first that ψ is a right projective cover. Then the exact sequence

$$0 \longrightarrow \mathbb{k}R \longrightarrow \mathbb{k}B \longrightarrow \mathbb{k}B / \mathbb{k}R \longrightarrow 0$$

splits because $\mathbb{k}B / \mathbb{k}R$ is projective. It follows that $\mathbb{k}R$ is projective and hence of the form $\eta \mathbb{k}B$ for some idempotent $\eta \in \mathbb{k}B$, which must then necessarily

be in $\mathbb{k}R$. But then η is a left identity for $\mathbb{k}R$, which is therefore a unital algebra as all left regular band algebras have a right identity by Theorem 4.2. Thus R is connected by Theorem 4.15. Suppose that $BR \neq \partial B$ and let $b \in \partial B \setminus BR$. Notice that if $Ba \supseteq Bb$, then $a \in B \setminus R$. Thus we can define a non-zero $\mathbb{k}B$ -module homomorphism $\alpha: \mathbb{k}B/\mathbb{k}R \rightarrow \mathbb{k}_{Bb}$ by

$$\begin{cases} \alpha(a + \mathbb{k}R) = 1, & \text{if } Ba \supseteq Bb \\ 0, & \text{else.} \end{cases}$$

This contradicts that $\mathbb{k}B/\mathbb{k}R$ has simple top $\mathbb{k}_{\hat{1}}$. Thus $BR = \partial B$.

We begin the proof of the converse by verifying that $\mathbb{k}B/\mathbb{k}R$ is indecomposable. This requires only assumption (1). Clearly, ψ is a surjective module homomorphism with kernel $\mathbb{k}\partial B/\mathbb{k}R$. We first show that $\ker \psi = \text{rad}(\mathbb{k}B/\mathbb{k}R)$. The inclusion $\text{rad}(\mathbb{k}B/\mathbb{k}R) \subseteq \ker \psi$ follows because $\mathbb{k}_{\hat{1}}$ is simple. We show the reverse inclusion by verifying that

$$\mathbb{k}\partial B/\mathbb{k}R \subseteq (\mathbb{k}B/\mathbb{k}R) \cdot \text{rad}(\mathbb{k}B).$$

Let $x \in \partial B$. By hypothesis, we can find $r \in R$ with $x \in Br$. Then $Bx = Brx$ and so $x - rx \in \text{rad}(\mathbb{k}B)$ by Proposition 4.13. But then $(1 + \mathbb{k}R)(x - rx) = x - rx + \mathbb{k}R = x + \mathbb{k}R$ because $rx \in R$. This completes the proof that $\text{rad}(\mathbb{k}B/\mathbb{k}R) = \ker \psi$. We deduce that $\mathbb{k}B/\mathbb{k}R$ is an indecomposable module (since a module over a finite dimensional algebra with simple top is indecomposable).

Now we prove that assumption (2) yields projectivity. Indeed, since R is connected we deduce that $\mathbb{k}R$ has an identity η by Theorem 4.15. As $\mathbb{k}R$ is a right ideal, clearly $\eta\mathbb{k}B \subseteq \mathbb{k}R$. As the reverse inclusion is obvious, we deduce that $\mathbb{k}B = \eta\mathbb{k}B \oplus (1 - \eta)\mathbb{k}B = \mathbb{k}R \oplus (1 - \eta)\mathbb{k}B$ and so $\mathbb{k}B/\mathbb{k}R \cong (1 - \eta)\mathbb{k}B$ is a projective right $\mathbb{k}B$ -module. $\square$

Remark 8.5. Let A be a set with at least two element and $F(A)$ the free left regular band monoid on A . As the elements of A are in distinct maximal $\mathcal{L}$ -classes of $F(A)$, the only proper right ideal R of $F(A)$ with $F(A)R = \partial F(A)$ is $R = \partial F(A)$. But the Hasse diagram of $\partial F(A)$ is a forest with $|A|$ connected components. It follows from Lemma 8.4 that the right projective cover of $\mathbb{k}_{\hat{1}}$ is not of the form $\mathbb{k}F(A)/\mathbb{k}R$ for any right ideal R of $F(A)$.

Our goal is to show that Lemma 8.4 can be applied to the algebras of a number of the CW left regular bands we have been considering.

8.1. Hyperplane arrangements. Let us begin with the case of the face monoid of a central arrangement. The idea behind what follows was inspired by considering the line shelling associated to a zonotope [122], but we will work with the dual picture. Let $\mathcal{A} = \{H_1, \dots, H_n\}$ be an essential central hyperplane in $V = \mathbb{R}^d$. Let us assume that H_i is defined by a form $f_i \in V^*$. Let $\theta: \mathbb{R}^d \rightarrow \mathcal{F}(\mathcal{A})$ be given by

$$\theta(x) = (\text{sgn}(f_1(x)), \dots, \text{sgn}(f_n(x))).$$

We say that a hyperplane H in V is *in general position* or *generic* with respect to $\mathcal{A}$ if H contains no element of the intersection lattice $\mathcal{L}(\mathcal{A}) \cong \Lambda(\mathcal{F}(\mathcal{A}))$ except the origin. Equivalently, if $f \in V^*$ is a form defining H , then $f(W) = \{0\}$ implies $W = \{0\}$ for all $W \in \mathcal{L}(\mathcal{A})$.

Proposition 8.6. *There exist generic hyperplanes with respect to $\mathcal{A}$.*

Proof. Suppose that $\mathcal{L}(\mathcal{A}) \setminus \{\{0\}\} = \{W_1, \dots, W_r\}$. Then the annihilator $W_i^\perp$ is a proper subspace of V^* . As a vector space over an infinite field is never a finite union of proper subspaces, we conclude that there is a form $f \in V^* \setminus \bigcup_{i=1}^r W_i^\perp$. The hyperplane H defined by $f = 0$ is generic with respect to $\mathcal{A}$. $\square$

Fix now a hyperplane H generic with respect to $\mathcal{A}$ defined by a form $f \in V^*$. We retain all the previous notation. Let $\hat{\mathcal{A}} = \mathcal{A} \cup \{H\}$ and let $\hat{\theta}: \mathbb{R}^d \rightarrow \mathcal{F}(\hat{\mathcal{A}})$ be given by

$$\hat{\theta}(x) = (\text{sgn}(f_1(x)), \dots, \text{sgn}(f_n(x)), \text{sgn}(f(x))).$$

The natural projection $\pi: L^{n+1} \rightarrow L^n$ defined by

$$\pi(x_1, \dots, x_{n+1}) = (x_1, \dots, x_n)$$

restricts to a surjective homomorphism $\pi: \mathcal{F}(\hat{\mathcal{A}}) \rightarrow \mathcal{F}(\mathcal{A})$ as $\pi(\hat{\theta}(x)) = \theta(x)$ for $x \in \mathbb{R}^d$. Denote by $\mathcal{F}^+(\hat{\mathcal{A}})$ the set of elements $(x_1, \dots, x_{n+1}) \in \mathcal{F}(\hat{\mathcal{A}})$ such that $x_{n+1} = +$. Note that $\mathcal{F}^+(\hat{\mathcal{A}})$ is a right ideal in $\mathcal{F}(\hat{\mathcal{A}})$ and is isomorphic to the face semigroup of the $(d-1)$ -dimensional affine arrangement obtained by intersecting the hyperplanes of $\mathcal{A}$ with the affine hyperplane defined by $f = 1$. Thus $\mathcal{F}^+(\hat{\mathcal{A}})$ is a connected CW left regular band by Proposition 3.14 (it is, in fact, an affine oriented matroid).

We define the *visual hemisphere* $R(H)$ of $\mathcal{A}$ with respect to H to be $\pi(\mathcal{F}^+(\hat{\mathcal{A}}))$. This is a right ideal of $\mathcal{F}(\mathcal{A})$. The reason for the name is that if one performs a line shelling of the zonotope $Z(\mathcal{A}) \subseteq V^*$ polar to $\mathcal{A}$ with respect to the line spanned by f in V^* , then the faces of $Z(\mathcal{A})$ corresponding to the elements of $R(H)$ make up the visual hemisphere of $Z(\mathcal{A})$ with respect to this shelling [122]. Note that $R(H)$ consists of those faces F containing a point x with $f(x) > 0$ by construction and hence is a realizable COMs.

Theorem 8.7. *Let $\mathcal{A}$ be an essential central hyperplane arrangement in $\mathbb{R}^d$, let $\mathbb{k}$ be a field and let H be a hyperplane in $\mathbb{R}^d$ which is generic with respect to $\mathcal{A}$. Let $\hat{1}$ be the maximum element of the intersection lattice $\mathcal{L}(\mathcal{A}) \cong \Lambda(\mathcal{F}(\mathcal{A}))$. Then the natural homomorphism*

$$\psi: \mathbb{k}\mathcal{F}(\mathcal{A})/\mathbb{k}R(H) \rightarrow \mathbb{k}_{\hat{1}}$$

is the right projective cover where $R(H)$ is the visual hemisphere of $\mathcal{A}$ with respect to H .

Proof. We verify that the right ideal $R(H)$ of $\mathcal{F}(\mathcal{A})$ satisfies the conditions of Lemma 8.4. We retain the previous notation. First we observe that

$R(H)$ is proper. Indeed, suppose that $0 \in R(H)$. Then $0 = \pi(\widehat{\theta}(x)) = \theta(x)$ with $\widehat{\theta}(x) \in \mathcal{F}^+(\widehat{\mathcal{A}})$. But as $\mathcal{A}$ is essential, $0 = \theta(x)$ implies $x = 0$. But then $\widehat{\theta}(0) = 0 \notin \mathcal{F}^+(\widehat{\mathcal{A}})$. This contradiction shows that $0 \notin R(H)$. Next we observe that $\pi: \mathcal{F}^+(\widehat{\mathcal{A}}) \rightarrow R(H)$ is an isomorphism because all elements of $\mathcal{F}^+(\widehat{\mathcal{A}})$ have $+$ as the last coordinate. Therefore, $R(H)$ is a connected left regular band, being isomorphic to the face semigroup of an affine hyperplane arrangement (or affine oriented matroid). It remains to verify that $\mathcal{F}(\mathcal{A})R(H) = \partial\mathcal{F}(\mathcal{A})$.

Let $F \in \mathcal{F}(\mathcal{A})$ with $F \neq 0$. We continue to denote by f the form defining the hyperplane H . As H is generic with respect to $\mathcal{A}$, it follows that the form f does not vanish on the span of F (which belongs to $\mathcal{L}(\mathcal{A})$) and hence does not vanish on F . Therefore, we can find $x \in F$ with $f(x) \neq 0$. Note that $F = \theta(x)$. If $f(x) > 0$, then $\widehat{\theta}(x) \in \mathcal{F}^+(\widehat{\mathcal{A}})$ and so $F = \theta(x) = \pi(\widehat{\theta}(x)) \in R(H)$. If $f(x) < 0$, then $f(-x) = -f(x) > 0$ and so $\widehat{\theta}(-x) \in \mathcal{F}^+(\widehat{\mathcal{A}})$. We conclude that $\theta(-x) = \pi(\widehat{\theta}(-x)) \in R(H)$. But $x, -x$ belong to the same elements of the intersection lattice $\mathcal{L}(\mathcal{A})$ and so $F = \theta(x) = \theta(x)\theta(-x)$. Thus $F \in \mathcal{F}(\mathcal{A})R(H)$. An application of Lemma 8.4 completes the proof of the theorem. $\square$

Combining Theorem 8.7 with Corollary 8.3 allows us to compute right projective covers and injective envelopes for all simple modules for a number of examples including face semigroups of central and affine hyperplane arrangements, convex sets of topes in central hyperplane arrangements, CAT(0) cube complexes and realizable COMs.

Corollary 8.8. *Let B be a connected left regular band and $\mathbb{k}$ a field. Let $X \in \Lambda(B)$ and assume that $X = Be_X$. Suppose that $\tau: e_X B \rightarrow \mathcal{F}(\mathcal{A})$ is an isomorphism where $\mathcal{A}$ is an essential central hyperplane arrangement and that H is a generic hyperplane with respect to $\mathcal{A}$ with associated visual hemisphere $R(H)$.*

- (1) *The right projective cover of $\mathbb{k}_X$ is the natural homomorphism*

$$\psi: \mathbb{k}e_X B / \mathbb{k}\tau^{-1}(R(H)) \rightarrow \mathbb{k}_X$$

induced by $e_X \mapsto 1$ and $b \mapsto 0$ for $b \in \partial e_X B$.

- (2) *The injective envelope of $\mathbb{k}_X$ is the left $\mathbb{k}B$ -module I_X consisting of all mappings $f: e_X B \rightarrow \mathbb{k}$ vanishing on $\tau^{-1}(R(H))$ with pointwise vector space operations and the natural left action of B given by $(bf)(a) = f(ab)$ for $b \in B$ and $a \in e_X B$. The simple socle of I_X is the subspace of those maps vanishing on $\partial e_X B$; it is isomorphic to $\mathbb{k}_X$.*

Recall that if $\mathcal{A}$ is a hyperplane arrangement (central or affine) and $G \in \mathcal{F}(\mathcal{A})$, then $G\mathcal{F}(\mathcal{A})$ is isomorphic to the face monoid of a central hyperplane arrangement which is a deletion of $\mathcal{A}$ (one deletes all the hyperplanes except those containing G). Thus Corollary 8.8 can be applied to compute the injective envelope of any simple module for a hyperplane face semigroup or,

more generally, for the semigroup associated to a T -convex set of topes for a hyperplane arrangement or a realizable COM.

Let us consider some examples.

Example 8.9 (Boolean arrangement). As an example, if $\mathcal{A}$ is the boolean arrangement in $\mathbb{R}^n$, then $F(\mathcal{A}) = L^n$. The hyperplane defined by

$$f(x_1, \dots, x_n) = x_1 + \dots + x_n$$

is generic with respect to $\mathcal{A}$ and $R(H)$ consists of those sign vectors with at least one positive coordinate. Therefore, $\mathbb{k}F(\mathcal{A})/\mathbb{k}R(H)$ has basis the cosets of elements of $\{0, -\}^n$ and hence has dimension 2^n . Consequently, the injective envelope of $\mathbb{k}_{\hat{1}}$ has dimension 2^n .

The zonotope associated to the boolean arrangement is the hypercube. Thus if K is a CAT(0) cube complex, X is an element of the intersection semilattice of K and C_X is a cube crossed exactly by the hyperplanes in X , then the injective envelope of $\mathbb{k}_X$ consists of all mappings from faces of C_X to $\mathbb{k}$ that vanish on those faces of C_X that lie in the positive half-space associated to some element of X where we fix an orientation of the hyperplanes of X .

Example 8.10 (Braid arrangement). Recall that the braid arrangement in $V = \mathbb{R}^n$ is defined by the set of hyperplanes $H_{ij} = \{x \in \mathbb{R}^n \mid x_i = x_j\}$ with $1 \leq i < j \leq n$. This is not an essential arrangement: the intersection of the hyperplanes is the line ℓ given by the equation $x_1 = x_2 = \dots = x_n$. Let $\mathcal{A}$ be the essential arrangement in V/ℓ induced by the braid arrangement. It is well known, and not too difficult to show (cf. [17]), that the elements of $\mathcal{F}(\mathcal{A})$ are in bijection with ordered set partitions of $[n] = \{1, \dots, n\}$. An ordered set partition $(P_1, \dots, P_r)$ corresponds to the face in $\mathbb{R}^n$ defined by the equations $x_i = x_j$ for i, j belonging to the same block and the inequalities $x_i < x_j$ if the block containing i is to the left of the block containing j . For example, when $n = 4$, the ordered set partition $(\{1, 3\}, \{2, 4\})$ corresponds to the face in $\mathbb{R}^n$ defined by $x_1 = x_3 < x_2 = x_4$. The product of ordered partitions is given by the simple rule:

$$(P_1, \dots, P_r)(Q_1, \dots, Q_s) = (P_1 \cap Q_1, \dots, P_1 \cap Q_s, \dots, P_r \cap Q_1, \dots, P_r \cap Q_s)^\wedge$$

where $(B_1, \dots, B_k)^\wedge$ is the result of removing all occurrences of the empty set from $(B_1, \dots, B_k)$. The support lattice $\Lambda(\mathcal{F}(\mathcal{A}))$ can be identified with the lattice of set partitions of $[n]$ ordered by refinement and the support map $\sigma: \mathcal{F}(\mathcal{A}) \rightarrow \Lambda(\mathcal{F}(\mathcal{A}))$ is given by $\sigma((P_1, \dots, P_r)) = \{P_1, \dots, P_r\}$.

We can identify the dual space of V/ℓ with the space of functionals on V which vanish on ℓ . It is not difficult to check that the functional

$$f(x) = (n-1)x_n - (x_1 + \dots + x_{n-1})$$

vanishes on ℓ and defines a generic hyperplane H with respect to $\mathcal{A}$. Clearly, f is positive at some point of the face corresponding to an ordered set partition $(P_1, \dots, P_r)$ if and only if $n \notin P_1$. Thus $\mathbb{k}\mathcal{F}(\mathcal{A})/\mathbb{k}R(H)$ has basis

the cosets of faces corresponding to ordered partitions $(P_1, \dots, P_r)$ with $n \in P_1$.

One can define an equivalence relation on ordered set partitions of $[n]$ by identifying cyclic conjugates, that is,

$$(P_1, P_2, \dots, P_r) \sim (P_2, P_3, \dots, P_r, P_1) \sim \dots \sim (P_r, P_1, \dots, P_{r-1}).$$

An equivalence class of ordered set partitions is called a *necklace of partitions of $[n]$* . Clearly, each necklace contains exactly one representative $(P_1, \dots, P_r)$ with $n \in P_1$. Thus the dimension of the injective envelope of $\mathbb{k}_{\hat{1}}$ is the number of necklaces of partitions of $[n]$. On the other hand, the results of the second author [98] (or Theorem 9.7 below) implies that if $X \in \Lambda(\mathcal{F}(\mathcal{A}))$, then the multiplicity of $\mathbb{k}_X$ as a composition factor of the injective envelope of $\mathbb{k}_{\hat{1}}$ is given by $|\mu(X, \hat{1})|$ where μ is the Möbius function of the partition lattice. As a consequence we obtain the following combinatorial identity, which can also be verified directly via a straightforward computation with the Möbius function of the partition lattice.

Proposition 8.11. The number of necklaces of partitions of $[n]$ is

$$\sum_{X \in \Pi_n} |\mu(X, \hat{1})|$$

where Π_n is the lattice of set partitions of $[n]$, μ is the Möbius function of Π_n and $\hat{1}$ is the partition into one block.

8.2. Oriented matroids. To adapt the proof of Theorem 8.7 to oriented matroids, we need an analogue of adjoining a generic hyperplane. Let $(E, \mathcal{L})$ be an oriented matroid and let $Z: \mathcal{L} \rightarrow \Lambda(\mathcal{L})$ be the map taking a sign vector in L^E to its zero set. A *cocircuit* of $\mathcal{L}$ is a maximal non-identity element of $\mathcal{L}$ (with respect to the natural partial order on $\mathcal{L}$) or, equivalently, a non-zero element $x \in \mathcal{L}$ with $Z(x)$ maximal with respect to inclusion. Note that y is a cocircuit if and only if $-y$ is one. It is a general fact about oriented matroids that they are generated as a monoid by their cocircuits (cf. the dual of [26, Proposition 3.7.2]).

An oriented matroid $(\tilde{E}, \tilde{\mathcal{L}})$ is a *single element extension* of an oriented matroid $(E, \mathcal{L})$ if $\tilde{E} = E \cup \{p\}$ for some $p \notin E$ and $\tilde{\mathcal{L}}$ is the image of $\mathcal{L}$ under the projection $\pi: L^{\tilde{E}} \rightarrow L^E$ given by $\pi(x) = x|_E$. Let $\mathcal{C}^*$ denote the set of cocircuits of $\mathcal{L}$. Then there is a unique mapping $\sigma: \mathcal{C}^* \rightarrow L$ satisfying $\sigma(-y) = -\sigma(y)$ for all $y \in \mathcal{C}^*$ and some additional axioms such that the set $\tilde{\mathcal{C}}^*$ of cocircuits of $\tilde{\mathcal{L}}$ is given by

$$\begin{aligned} \tilde{\mathcal{C}}^* = & \{(y, \sigma(y)) \mid y \in \mathcal{C}^*\} \cup \\ & \{(y_1 y_2, 0) \mid y_1, y_2 \in \mathcal{C}^*, \sigma(y_1) = -\sigma(y_2) \neq 0, \\ & S(y_1, y_2) = \emptyset, \rho(y_1 y_2) = 2\} \end{aligned} \quad (8.1)$$

where ρ is obtained from the rank function on the geometric lattice $\Lambda(\mathcal{L})$ by $\rho(x) = \text{rank}(E) - \text{rank}(Z(x))$. See [26, Proposition 7.1.4 and Theorem 7.1.8]

for details. We will write $\mathcal{L}_\sigma$ for the single element extension of $\mathcal{L}$ associated to a mapping $\sigma: \mathcal{C}^* \rightarrow L$ satisfying the equivalent conditions of [26, Theorem 7.1.8] for defining a single element extension $\mathcal{L}$.

A single element extension $\mathcal{L}_\sigma$ of $\mathcal{L}$ is said to be *in general position* or *generic* if $\sigma(y) \neq 0$ for all $y \in \mathcal{C}^*$. Such extensions exist (cf. [26, Proposition 7.2.2]). One method to construct them is via lexicographic extensions. Here is the setup. Let E be the ground set of $\mathcal{L}$, let $I = \{e_1, \dots, e_k\}$ be a linearly ordered subset of E and let $\alpha_1, \dots, \alpha_k \in \{+, -\}$. Define a mapping $\sigma: \mathcal{C}^* \rightarrow L$ by

$$\sigma(y) = \begin{cases} \alpha_i y_{e_i}, & \text{if } i \text{ is minimal with } y_{e_i} \neq 0 \\ 0, & \text{if } y_{e_i} = 0 \text{ for all } 1 \leq i \leq k. \end{cases}$$

Then σ defines a single element extension $\mathcal{L}_\sigma$ called a *lexicographic extension* of $\mathcal{L}$. See [26, Proposition 7.2.4]. If $I = E$, then $\mathcal{L}_\sigma$ is obviously generic.

For example, if $\mathcal{L}$ is the oriented matroid associated to a central hyperplane arrangement and f_i is the form defining the hyperplane $e_i \in I$, for $1 \leq i \leq k$, then the lexicographic extension associated to I and the α_i is obtained by adding the hyperplane $f = 0$ to the original arrangement where

$$f = \varepsilon \alpha_1 f_1 + \varepsilon^2 \alpha_2 f_2 + \dots + \varepsilon^k \alpha_k f_k$$

for some small $\varepsilon > 0$. See the discussion preceding [26, Proposition 7.2.4].

Let $(E, \mathcal{L})$ be an oriented matroid and let $(E \cup \{p\}, \mathcal{L}_\sigma)$ be a generic single element extension with corresponding map $\sigma: \mathcal{C}^* \rightarrow L$ where $\mathcal{C}^*$ denotes the set of cocircuits of $\mathcal{L}$. Then there is a corresponding affine oriented matroid $(E \cup \{p\}, \mathcal{L}_\sigma, p)$ with associated left regular band $\mathcal{L}_\sigma^+(p) = \{x \in \mathcal{L}_\sigma \mid x_p = +\}$. Let $\pi: \mathcal{L}_\sigma \rightarrow \mathcal{L}$ be the projection $\pi(x) = x|_E$. We define $R(\sigma) = \pi(\mathcal{L}_\sigma^+(p))$ to be the *visual hemisphere* associated to σ . Since $\mathcal{L}_\sigma^+(p)$ is a right ideal of $\mathcal{L}_\sigma$ and π is surjective, we conclude that $R(\sigma)$ is a right ideal of $\mathcal{L}$.

Theorem 8.12. *Let $(E, \mathcal{L})$ be an oriented matroid, let $\mathbb{k}$ be a field and let $(E \cup \{p\}, \mathcal{L}_\sigma)$ be a generic single element extension with corresponding map $\sigma: \mathcal{C}^* \rightarrow L$ where $\mathcal{C}^*$ denotes the set of cocircuits of $\mathcal{L}$. Let $\hat{1} = \mathcal{L}$ be the maximum element of $\Lambda(\mathcal{L})$. Then the natural homomorphism*

$$\psi: \mathbb{k}\mathcal{L}/\mathbb{k}R(\sigma) \rightarrow \mathbb{k}_{\hat{1}}$$

is the right projective cover where $R(\sigma)$ is the visual hemisphere associated to σ .

Proof. We check that the right ideal $R(\sigma)$ of $\mathcal{L}$ satisfies the conditions of Lemma 8.4. Let us maintain the previous notation. The set of cocircuits of $\mathcal{L}_\sigma$ will be denoted by $\tilde{\mathcal{C}}^*$. Notice that $\tilde{\mathcal{C}}^*$ contains no element of the form $(0, z)$ with $z \in \{+, -\}$ by (8.1) because $0 \notin \mathcal{C}^*$. Thus $\mathcal{L}_\sigma$ contains no such element because $\tilde{\mathcal{C}}^*$ generates $\mathcal{L}_\sigma$. Therefore, $(0, +) \notin \mathcal{L}_\sigma^+(p)$ and so $R(\sigma)$ is proper.

Next observe that $\pi: \mathcal{L}_\sigma^+(p) \rightarrow R(\sigma)$ is an isomorphism since all elements of $\mathcal{L}_\sigma^+(p)$ have $+$ as the last coordinate. Therefore, $R(\sigma)$ is a connected left

regular band as affine oriented matroids are such. It remains to verify that $\mathcal{L}R(\sigma) = \partial\mathcal{L}$.

Let $y \in \mathcal{L} \setminus \{0\}$. Then $y = y_1 \cdots y_m$ where $y_1, \dots, y_m$ are cocircuits. Put $\tilde{y}_i = (y_i, \sigma(y_i)) \in \tilde{\mathcal{C}}^*$ (using (8.1)). Note that $\pi(\tilde{y}_i) = y_i$. Since $\mathcal{L}_\sigma$ is generic, either $\sigma(y_1) = +$ or $\sigma(y_1) = -$. Assume first that $\sigma(y_1) = +$. Then $\tilde{y}_1 = (y_1, +)$ and so $\tilde{y} = \tilde{y}_1 \cdots \tilde{y}_m \in \mathcal{L}_\sigma^+(p)$ and hence $y = y_1 \cdots y_m = \pi(\tilde{y}) \in R(\sigma)$. Next assume that $\sigma(y_1) = -$. Then $-\tilde{y}_1 = (-y_1, +) \in \mathcal{L}_\sigma^+(p)$ and $\pi(-\tilde{y}_1) = -y_1$. Therefore, $(-y_1)y_2 \cdots y_m \in R(\sigma)$ and hence $y = y_1 \cdots y_m = y_1(-y_1)y_2 \cdots y_m \in \mathcal{L}R(\sigma)$ using that $x(-x) = x$ in L^E .

The theorem now follows via an application of Lemma 8.4. $\square$

One advantage of Theorem 8.12 over Theorem 8.7 for the case of central hyperplane arrangement is that the former is combinatorial: one can effectively construct a generic lexicographic single element extension. Writing explicitly a generic hyperplane is more complicated.

Theorem 8.12, together with Corollary 8.3, allows us to compute right projective covers and injective envelopes for all simple modules for left regular bands associated to COMs including oriented matroids and affine oriented matroids. We summarize in the following corollary.

Corollary 8.13. *Let B be a connected left regular band and $\mathbb{k}$ a field. Let $X \in \Lambda(B)$ with $X = Be_X$. Suppose that $\tau: e_X B \rightarrow \mathcal{L}$ is an isomorphism where $\mathcal{L}$ is the monoid of covectors of an oriented matroid and that $\mathcal{L}_\sigma$ is a generic single element extension of $\mathcal{L}$ with associated visual hemisphere $R(\sigma)$.*

- (1) *The right projective cover of $\mathbb{k}_X$ is the natural homomorphism*

$$\psi: \mathbb{k}e_X B / \mathbb{k}\tau^{-1}(R(\sigma)) \rightarrow \mathbb{k}_X$$

induced by $e_X \mapsto 1$ and $b \mapsto 0$ for $b \in \partial e_X B$.

- (2) *The injective envelope of $\mathbb{k}_X$ is the left $\mathbb{k}B$ -module I_X consisting of all mappings $f: e_X B \rightarrow \mathbb{k}$ vanishing on $\tau^{-1}(R(\sigma))$ with pointwise vector space operations and the natural left action of B given by $(bf)(a) = f(ab)$ for $b \in B$ and $a \in e_X B$. The simple socle of I_X (isomorphic to $\mathbb{k}_X$) is the subspace of those maps vanishing on $\partial e_X B$.*

Recall that if $(E, \mathcal{L})$ is an oriented matroid and $x \in \mathcal{L}$, then $x\mathcal{L}$ is isomorphic to the monoid of covectors for the deletion $(E \setminus A, \mathcal{L} \setminus A)$ where $A = E \setminus Z(x)$. Thus Corollary 8.13 can be used to compute the injective envelopes of all simple modules for the semigroup of an oriented matroid, affine oriented matroid or T -convex set of topes. More generally, $x\mathcal{L}$ is isomorphic to the monoid of covectors of an oriented matroid for any COM $(E, \mathcal{L})$ (cf. [13]) and so Corollary 8.13 can be applied to compute injective envelopes for algebras of COMs.

It is an open problem to give an explicit construction of injective indecomposables for more general classes of left regular band algebras.

9. ENUMERATION OF CELLS FOR CW LEFT REGULAR BANDS

In this section we prove analogues of Zaslavsky's theorem [26, Theorem 4.6.1], and other enumerative results from hyperplane/oriented matroid theory, for connected CW left regular bands. For CAT(0) cube complexes, our results recover those of Dress *et. al* [51], which were originally stated in the language of median graphs. In the case of complex hyperplane arrangements, our result specializes to [23]. When comparing our results to those in [26], one should bear in mind that people in hyperplane and oriented matroid theory work with the dual cell complex to the monoid of covectors and they do not include the identity as corresponding to a face.

9.1. Flag vectors. If X is a (finite) d -dimensional regular CW complex, then f_k denotes the number of k -dimensional cells of X for $0 \leq k \leq d$. The sequence

$$f(X) = (f_0, \dots, f_d)$$

is called the f -vector of X . The Euler characteristic of X is

$$\chi(X) = \sum_{k=0}^d (-1)^k f_k.$$

It is well known that

$$\chi(X) = \sum_{k=0}^d (-1)^k \beta_k(X)$$

where $\beta_k(X)$ is the dimension of $H_k(X; \mathbb{Q})$.

More generally, for $J = \{j_1, \dots, j_k\}$ with $0 \leq j_1 < j_2 < \dots < j_k \leq d$, let f_J be the number of chains of cells $e_1 < e_2 < \dots < e_k$ in the face poset $\mathcal{P}(X)$ such that $\dim e_i = j_i$ for $1 \leq i \leq k$. The array $\tilde{f}(X) = (f_J)_{J \subseteq \{0, \dots, d\}}$ is called the *flag vector* (or, *Fine f -vector*) of X . In this section, we will compute $f(\Sigma(B))$ and $\tilde{f}(\Sigma(B))$ for the regular cell complex $\Sigma(B)$ associated to a connected CW left regular band B in terms of $\Lambda(B)$. This generalizes well-known results for hyperplane arrangements and oriented matroids [26, Section 4.6].

We begin with a lemma about cellular maps between posets satisfying properties analogous to that of the map from a connected CW left regular band to its support semilattice.

Lemma 9.1. *Let $f: P \rightarrow Q$ be a cellular mapping of finite posets such that:*

- (1) Q has a minimum $\hat{0}$;
- (2) f preserves strict inequalities;
- (3) $f^{-1}(Q_{\geq q})$ is an acyclic CW poset for all $q \in Q$.

Then

$$|f^{-1}(q)| = \sum_{q' \geq q} (-1)^{\dim q' - \dim q} \mu_Q(q, q')$$

where μ_Q is the Möbius function of Q .

If, in addition, each open interval of Q is a Cohen-Macaulay poset, then

$$|f^{-1}(q)| = \sum_{q' \geq q} |\mu_Q(q, q')|$$

holds.

Proof. Note that $P = f^{-1}(Q_{\geq \hat{0}})$ is a CW poset and hence is graded. Therefore, Q is graded by Corollary 3.7. As the Euler characteristic of any acyclic CW complex is 1 and f preserves dimensions (because it preserves strict inequalities), we obtain

$$1 = \chi(f^{-1}(Q_{\geq q})) = \sum_{p \in f^{-1}(Q_{\geq q})} (-1)^{\dim p - \dim q} = \sum_{q' \geq q} (-1)^{\dim q' - \dim q} |f^{-1}(q')|.$$

Consequently, we have

$$(-1)^{\dim q} = \sum_{q' \geq q} (-1)^{\dim q'} |f^{-1}(q')|$$

and so by Möbius inversion we obtain

$$(-1)^{\dim q} |f^{-1}(q)| = \sum_{q' \geq q} (-1)^{\dim q'} \mu_Q(q, q').$$

It follows that

$$|f^{-1}(q)| = \sum_{q' \geq q} (-1)^{\dim q' - \dim q} \mu_Q(q, q')$$

as required.

The final statement follows because $(-1)^{\dim q' - \dim q} \mu_Q(q, q') \geq 0$ holds if $\Delta((q, q'))$ is Cohen-Macaulay (cf. [108, Theorem 10]). $\square$

Recall that, for a poset Q with minimum $\hat{0}$, one puts $\mu_Q(x) = \mu_Q(\hat{0}, x)$.

Corollary 9.2. *With $f: P \rightarrow Q$ as in Lemma 9.1, one has*

$$|f^{-1}(\hat{0})| = \sum_{q \geq \hat{0}} (-1)^{\dim q} \mu_Q(\hat{0}, q) = \sum_{q \geq \hat{0}} (-1)^{\dim q} \mu_Q(q).$$

In addition, if each open interval $(\hat{0}, q)$ of Q is Cohen-Macaulay, then

$$|f^{-1}(\hat{0})| = \sum_{q \in Q} |\mu_Q(q)|$$

holds.

Applying this result to the case of a connected CW left regular band, we obtain the following result, generalizing the Las Vergnas-Zaslavsky Theorem [26, Theorem 4.6.1] for hyperplane arrangements and oriented matroids.

Theorem 9.3. *Let B be a connected CW left regular band with support homomorphism $\sigma: B \rightarrow \Lambda(B)$. Let $X \in \Lambda(B)$. Then*

$$|\sigma^{-1}(X)| = \sum_{Y \geq X} |\mu_{\Lambda(B)}(X, Y)|$$

where $\mu_{\Lambda(B)}$ is the Möbius function of $\Lambda(B)$. In particular, the size of the minimal ideal of B is

$$f_0(\Sigma(B)) = \sum_{X \in \Lambda(B)} |\mu_{\Lambda(B)}(X)|.$$

More generally,

$$f_k(\Sigma(B)) = \sum_{\dim A=k, B \geq A} |\mu_{\Lambda(B)}(A, B)|$$

holds.

Proof. The support homomorphism $\sigma: B \rightarrow \Lambda(B)$ is a cellular mapping preserving strict inequalities by Proposition 3.6. The semilattice $\Lambda(B)$ has a minimum element and $\sigma^{-1}(\Lambda(B)_{\geq X}) = B_{\geq X}$ is an acyclic CW poset for $X \in \Lambda(B)$ by Theorem 5.5 and the definition of a CW left regular band. Moreover, each open interval in $\Lambda(B)$ is Cohen-Macaulay by Theorem 7.14. Lemma 9.1 now yields the result. $\square$

Notice in particular, that the cardinalities of B and all its $\mathcal{L}$ -classes depend only on $\Lambda(B)$. This can also be deduced from the fact that $\mathbb{k}B$ depends only on $\Lambda(B)$ up to isomorphism.

We now specialize Theorem 9.3 to the case of a CAT(0) cube complex in order to recover a result of Dress *et al.* [51, Remark of Section 3], which is stated there in the language of median graphs. To translate between our results and theirs, one should use the equivalence between CAT(0) cube complexes and median graphs and the bijection between hyperplanes and what these authors call splits, cf. [94].

Theorem 9.4. *Let K be a finite CAT(0) cube complex with set of hyperplanes $\mathcal{H}$. Let Γ be the graph with vertex set $\mathcal{H}$ and where two hyperplanes are connected if they intersect. Then we have*

$$f_k(K) = \sum_{i \geq k} \binom{i}{k} f_{i-1}(\text{Cliq}(\Gamma))$$

where $f_{-1}(\text{Cliq}(\Gamma)) = 1$ by convention. In particular, $f_0(K)$ is one more than the number of non-empty simplices in $\text{Cliq}(\Gamma)$.

Proof. If K is a CAT(0) cube complex, then $\Lambda(\mathcal{F}(K))$ is isomorphic to the face poset, together with an empty face, of the nerve of the collection $\mathcal{H}$ by Proposition 3.25 and Theorem 3.27. Note, though, that the empty simplex has rank 0 and hence the rank of a simplex in $\Lambda(\mathcal{F}(K))$ is its number of vertices, rather than its dimension. Also, by the Helly property for CAT(0)

cube complexes [42], the nerve of $\mathcal{H}$ is a flag complex and hence isomorphic to the clique complex of Γ . Note that the interval $[X, Y]$ is isomorphic to a boolean algebra for each $X \leq Y$ in $\Lambda(\mathcal{F}(K))$. Therefore, $\mu_{\Lambda(\mathcal{F}(K))}(X, Y) = (-1)^{\text{rk}[X, Y]}$ and so $|\mu_{\Lambda(\mathcal{F}(K))}(X, Y)| = 1$.

If $i \geq k$, then there are $f_{i-1}(\text{Cliq}(\Gamma))$ elements $Y \in \Lambda(\mathcal{F}(K))$ with i vertices. Identifying Y with an i -element clique in Γ , we see that Y has $\binom{i}{k}$ subcliques with k elements and hence there are $\binom{i}{k}$ choices of $X \in \Lambda(\mathcal{F}(K))$ with X having rank k and $X \leq Y$. Denoting the number of vertices of $X \in \Lambda(\mathcal{F}(K))$ by $|X|$, it follows from Theorem 9.3 that

$$\begin{aligned} f_k(K) &= \sum_{|X|=k, Y \geq X} |\mu_{\Lambda(\mathcal{F}(K))}(X, Y)| \\ &= \sum_{i \geq k} \sum_{|Y|=i} \left(\sum_{|X|=k, X \leq Y} |\mu_{\Lambda(\mathcal{F}(K))}(X, Y)| \right) \\ &= \sum_{i \geq k} \binom{i}{k} f_{i-1}(\text{Cliq}(\Gamma)) \end{aligned}$$

using that $|\mu_{\Lambda(\mathcal{F}(K))}(X, Y)| = 1$. This completes the proof. $\square$

For example, if T is a tree (i.e., a one-dimensional CAT(0) cube complex), then the hyperplanes of T are the midpoints of edges. No two hyperplanes intersect. Theorem 9.4 then recovers, in this case, the well-known fact that the number of vertices of a tree is one more than the number of edges.

The next proposition generalizes [26, Proposition 4.6.2]. We use here that the support map $\sigma: B \rightarrow \Lambda(B)$ takes chains to chains of the same dimension.

Proposition 9.5. *Let B be a connected CW left regular band and suppose that $X_1 < X_2 < \dots < X_k$ is a chain in $\Lambda(B)$. Then*

$$|\sigma^{-1}(X_1, X_2, \dots, X_k)| = \prod_{i=1}^k \sum_{X_i \leq Y \leq X_{i+1}} |\mu_{\Lambda(B)}(X_i, Y)|$$

where $X_k \leq Y \leq X_{k+1}$ is interpreted as $X_k \leq Y$.

Proof. We proceed by induction, the case $k = 1$ being handled by Theorem 9.3. Assume it is true for $k - 1$. By Lemma 3.3, to choose a preimage of $(X_1, X_2, \dots, X_k)$ is to choose a preimage x_k of X_k in B and then a preimage of $(X_1, X_2, \dots, X_{k-1})$ in $x_k B$ under the deletion $\sigma: x_k B \rightarrow \Lambda(x_k B) = \Lambda(B)_{\leq X_k}$. Note that $x_k B$ is a connected CW left regular band. By Theorem 9.3 there are $\sum_{X_k \leq Y} |\mu_{\Lambda(B)}(X_k, Y)|$ ways to choose x_k and by induction there are then

$$\prod_{i=1}^{k-1} \sum_{X_i \leq Y \leq X_{i+1}} |\mu_{\Lambda(B)}(X_i, Y)|$$

ways to choose a preimage of $(X_1, \dots, X_{k-1})$ in $x_k B$. Thus

$$|\sigma^{-1}(X_1, X_2, \dots, X_k)| = \prod_{i=1}^k \sum_{X_i \leq Y \leq X_{i+1}} |\mu_{\Lambda(B)}(X_i, Y)|$$

completing the induction. $\square$

We can now prove the following analogue for connected CW left regular bands of a result of Bayer and Sturmfels [26, Corollary 4.6.3].

Corollary 9.6. *The flag vector $\tilde{f}(\Sigma(B))$ for a connected CW left regular band B depends only on its support semilattice $\Lambda(B)$. More precisely, if $\dim B = d$ and $J = \{j_1, \dots, j_k\}$ with $0 \leq j_1 < j_2 < \dots < j_k \leq d$, then*

$$f_J(\Sigma(B)) = \sum \prod_{i=1}^k \sum_{X_i \leq Y \leq X_{i+1}} |\mu_{\Lambda(B)}(X_i, Y)|$$

where the first sum runs over all chains $(X_1, \dots, X_k)$ in $\Lambda(B)$ with $\dim X_i = j_i$ and where $X_k \leq Y \leq X_{k+1}$ is interpreted as $X_k \leq Y$.

Proof. Since $\sigma: B \rightarrow \Lambda(B)$ preserves dimensions of flags, it is clear that

$$f_J(\Sigma(B)) = \sum |\sigma^{-1}(X_1, \dots, X_k)|$$

where the sum runs over all chains $(X_1, \dots, X_k)$ in $\Lambda(B)$ with $\dim X_i = j_i$. The result now follows from Proposition 9.5. $\square$

9.2. Cartan invariants. We use the previous results of this section to compute the Cartan matrix for a connected CW left regular band. If B is any left regular band, recall that the Cartan matrix of $\mathbb{k}B$ is the mapping $C: \Lambda(B) \times \Lambda(B) \rightarrow \mathbb{N}$ with $C_{X,Y}$ the multiplicity of the simple module $\mathbb{k}_X$ as a composition factor in the projective cover $\mathbb{k}L_Y$ of $\mathbb{k}_Y$. Suppose that $X = Be_X$ for $X \in \Lambda(B)$. Then Theorem 4.17 states

$$C_{X,Y} = \sum_{Y \leq Z \leq X} |e_Z B \cap L_Y| \cdot \mu_{\Lambda(B)}(Z, X).$$

Therefore, we have by Möbius inversion that

$$|e_X B \cap L_Y| = \sum_{Y \leq Z \leq X} C_{Z,Y} \tag{9.1}$$

for $X \geq Y$.

Assume now that B is a connected CW left regular band and $Y \leq X$ in $\Lambda(B)$. Then $e_X B_{\geq Y}$ is a connected CW left regular band monoid with support lattice $[Y, X]$ and $e_X B \cap L_Y$ is the vertex set of this complex. Thus, by Theorem 9.3, we have that

$$|e_X B \cap L_Y| = \sum_{Y \leq Z \leq X} |\mu_{\Lambda(B)}(Y, Z)|. \tag{9.2}$$

A comparison of (9.1) and (9.2), together with an application of Möbius inversion, yields $C_{X,Y} = |\mu_{\Lambda(B)}(Y, X)|$ for $X \geq Y$. We have thus proved the following theorem, generalizing a result of the second author for hyperplane face monoids [98].

Theorem 9.7. *Let B be a connected CW left regular band and $\mathbb{k}$ a field. Then the Cartan matrix for $\mathbb{k}B$ is given by*

$$C_{X,Y} = \begin{cases} |\mu_{\Lambda(B)}(Y, X)|, & \text{if } X \geq Y \\ 0, & \text{else.} \end{cases}$$

Example 9.8. If K is a CAT(0) cube complex, then $\Lambda(\mathcal{F}(K))$ is isomorphic to the face poset, together with an empty face, of the nerve of the collection of hyperplanes of K and hence each interval $[Y, X]$ is isomorphic to a boolean algebra. Therefore, $\mu_{\Lambda(\mathcal{F}(K))}(Y, X) = \pm 1$ and so $|\mu_{\Lambda(\mathcal{F}(K))}(Y, X)| = 1$. Thus the Cartan matrix of $\mathbb{k}\mathcal{F}(K)$ is the zeta function of $\Lambda(\mathcal{F}(K))$. This also follows from the fact that $\mathbb{k}\mathcal{F}(K)$ is isomorphic to the incidence algebra of $\Lambda(\mathcal{F}(K))$ by Corollary 7.10.

10. COHOMOLOGICAL DIMENSION

In this section, we compute the cohomological dimension of a finite left regular band monoid. Monoid cohomology has a long history, in part because of its connections with string rewriting systems [2, 32, 47, 56, 59, 60, 65–67, 82, 84, 86, 88, 107]. See [115] for the cohomology of categories with applications.

Let M be a monoid. Then the cohomology of M with coefficients in a $\mathbb{Z}M$ -module V is defined by

$$H^n(M, V) = \text{Ext}_{\mathbb{Z}M}^n(\mathbb{Z}, V).$$

If V is a $\mathbb{k}M$ -module for some commutative ring with unit $\mathbb{k}$, then it is known that $\text{Ext}_{\mathbb{k}M}^n(\mathbb{k}, V) = H^n(M, V)$ (use the standard bar resolution). The $\mathbb{k}$ -cohomological dimension of M is

$$\text{cd}_{\mathbb{k}} M = \sup\{n \mid H^n(M, V) \neq 0, \text{ for some } \mathbb{k}M\text{-module } V\}.$$

Obviously $\text{cd}_{\mathbb{k}} M \leq \text{cd}_{\mathbb{Z}} M = \text{cd } M$. Using the projective resolution of the trivial module from Theorem 5.11, we can compute the cohomological dimension of a finite left regular band monoid over any base ring.

Theorem 10.1. *Let B be a left regular band monoid and $\mathbb{k}$ a commutative ring with unit. Then $\text{cd}_{\mathbb{k}} B$ is the largest n such that $\tilde{H}^{n-1}(\Delta(\partial e_Y B); \mathbb{k}) \neq 0$ for some $Y \in \Lambda(B)$.*

Proof. Theorem 5.11 shows that the trivial $\mathbb{k}B$ -module admits a finite projective resolution by finitely generated $\mathbb{k}B$ -modules. Thus $\text{cd}_{\mathbb{k}} B < \infty$. Suppose that $n = \text{cd}_{\mathbb{k}} B$. Then the functor $H^n(B, -)$ is right exact by consideration of the long exact sequence for cohomology. Because every module is a quotient of a free module, it then follows that $H^n(B, F) \neq 0$ for some free $\mathbb{k}B$ -module F . Since the trivial $\mathbb{k}B$ -module admits a projective resolution

that is finitely generated in each degree, $H^n(B, -)$ preserves direct limits by a result of Brown [30]. As every free module is a direct limit of finitely generated free modules and $H^n(B, -)$ commutes with finite direct sums, it follows that $H^n(B, \mathbb{k}B) \neq 0$. We conclude that n is the largest non-negative integer for which $H^n(B, \mathbb{k}B) \neq 0$.

Let $J = \ker \sigma$ where $\sigma: \mathbb{k}B \rightarrow \mathbb{k}\Lambda(B)$ is induced by the support homomorphism. Recall from Theorem 4.11 that J is nilpotent and let $m \geq 1$ be such that $J^m = 0$. Then we have a filtration

$$0 = J^m \mathbb{k}B \subseteq J^{m-1} \mathbb{k}B \subseteq \cdots \subseteq J^0 \mathbb{k}B = \mathbb{k}B.$$

It follows using the long exact sequence for cohomology that $\text{cd}_{\mathbb{k}} B$ is the largest integer $n \geq 0$ such that

$$H^n(B, J^j \mathbb{k}B / J^{j+1} \mathbb{k}B) \neq 0$$

for some $0 \leq j \leq m-1$. But each $J^j \mathbb{k}B / J^{j+1} \mathbb{k}B$ is a $\mathbb{k}\Lambda(B)$ -module. Thus $\text{cd}_{\mathbb{k}} B$ is the largest integer n such that $H^n(B, V) \neq 0$ for some $\mathbb{k}\Lambda(B)$ -module V . But, arguing as above, this is the same as the largest integer n such that $H^n(B, \mathbb{k}\Lambda(B)) \neq 0$. We conclude from (4.7) that

$$\text{cd}_{\mathbb{k}} B = \max\{n \mid H^n(B, \mathbb{k}_X) \neq 0, \text{ for some } X \in \Lambda(B)\}$$

The result now follows from Theorem 5.17. $\square$

Let us state some corollaries.

Corollary 10.2. *Let B be a left regular band monoid and $\mathbb{k}$ a commutative ring with unit. Then $\text{cd}_{\mathbb{k}} \mathbf{S}(B) = \text{cd}_{\mathbb{k}} B + 1$*

Proof. This is immediate from Theorem 10.1 and Corollary 5.19. $\square$

Note that in the next two corollaries we do *not* require $B_{\geq X}$ to be a CW poset for all $X \in \Lambda(B)$.

Corollary 10.3. *Let B be a left regular band monoid which is a CW poset. Then $\text{cd}_{\mathbb{k}} B = \text{cd } B = \dim \Sigma(B)$. This applies in particular to real and complex hyperplane face monoids, oriented matroids and oriented interval greedoids.*

Proof. The $\|\Delta(\partial e_Y B)\|$ are precisely the boundary spheres of the closed cells of $\Sigma(B)$. The result is then immediate from Theorem 10.1. $\square$

Corollary 10.4. *Let B be a left regular band which is an acyclic CW poset. Then $\text{cd}_{\mathbb{k}} B^1 = \text{cd } B^1 = \dim \Sigma(B)$. This applies in particular to covector semigroups of COMs such as affine hyperplane arrangements, $\text{CAT}(0)$ cube complexes and affine oriented matroids.*

Proof. One has $\Delta(\partial B^1) = \Delta(B)$ is acyclic and $\partial e_Y B^1 = \partial e_Y B$ for $Y \in \Lambda(B)$. The $\|\Delta(\partial e_Y B)\|$ are precisely the boundary spheres of the closed cells of $\Sigma(B)$. The result is then immediate from Theorem 10.1. $\square$

We end by giving a surprising connection between the cohomological dimension of free partially commutative left regular bands and the Leray number of flag complexes. Connections between right-angled Artin groups and flag complexes can be found in [16].

Corollary 10.5. *If Γ is a graph, then the $\mathbb{k}$ -cohomological dimension of the free partially commutative left regular band associated to Γ is the $\mathbb{k}$ -Leray number of Γ . In particular, $\text{cd}(B(\Gamma)) \leq 1$ if and only if Γ is chordal.*

Proof. This is immediate from Proposition 3.14 and Theorem 10.1. $\square$

It is well known that in M and N are monoids, then

$$\max\{\text{cd } M, \text{cd } N\} \leq \text{cd}(M \times N) \leq \text{cd}(M) + \text{cd}(N) \quad (10.1)$$

cf. [60]. In particular, if $\text{cd } N = 0$, then $\text{cd}(M \times N) = \text{cd } M$. Also, it is known that if M is a monoid containing a right zero, then $\text{cd } M = 0$. Indeed, if e is a right zero then $\mathbb{Z} \cong \mathbb{Z}Me$ and so $\mathbb{Z}$ is projective and hence has projective dimension 0. In particular, if B is a left regular band monoid then $\text{cd } B^{op} = 0$. Recall that a *regular band* is a subdirect product of a left regular band and a right regular band (defined dually).

Corollary 10.6. *For any pair of non-negative integers m, n , there is a finite regular band monoid B with $\text{cd } B = m$ and $\text{cd } B^{op} = n$.*

Proof. Recall the definition of ladders from Example 3.17. We have $\text{cd } L_m = m$ and $\text{cd } L_n = n$ by Corollary 10.3 and so if $B = L_m \times L_n^{op}$, then $\text{cd } B = \text{cd } L_m = m$ and $\text{cd } B^{op} = \text{cd } L_n = n$ by (10.1). $\square$

Of course, we can replace L_m and L_n by face monoids of essential hyperplane arrangements in $\mathbb{R}^m$ and $\mathbb{R}^n$, respectively, and still get the same result. In [60] it is shown that, for any monoid M and for any $0 \leq m, n \leq \infty$, there is a monoid N containing M as a submonoid with $\text{cd } N = m$ and $\text{cd } N^{op} = n$.

REFERENCES

- [1] P. Abramenko and K. S. Brown. *Buildings*, volume 248 of *Graduate Texts in Mathematics*. Springer, New York, 2008. Theory and applications.
- [2] W. W. Adams and M. A. Rieffel. Adjoint functors and derived functors with an application to the cohomology of semigroups. *J. Algebra*, 7:25–34, 1967.
- [3] I. Agol. The virtual Haken conjecture. With an appendix by Agol, Daniel Groves, and Jason Manning. *Doc. Math.* 18:1045–1087, 2013.
- [4] M. Aguiar and S. Mahajan. *Coxeter groups and Hopf algebras*, volume 23 of *Fields Institute Monographs*. American Mathematical Society, Providence, RI, 2006. With a foreword by Nantel Bergeron.
- [5] M. Aguiar and S. Mahajan. *Monoidal functors, species and Hopf algebras*, volume 29 of *CRM Monograph Series*. American Mathematical Society, Providence, RI, 2010. With forewords by Kenneth Brown and Stephen Chase and André Joyal.
- [6] I. Assem, D. Simson, and A. Skowroński. *Elements of the representation theory of associative algebras. Vol. 1*, volume 65 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2006. Techniques of representation theory.
- [7] C. A. Athanasiadis and P. Diaconis. Functions of random walks on hyperplane arrangements. *Adv. in Appl. Math.*, 45(3):410–437, 2010.

- [8] M. Auslander, I. Reiten, and S. O. Smalø. *Representation theory of Artin algebras*, volume 36 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1997. Corrected reprint of the 1995 original.
- [9] H. J. Bandelt. Retracts of hypercubes. *J. Graph Theory* 8(4):501–510, 1984.
- [10] H. J. Bandelt and V. Chepoi. Metric graph theory and geometry: a survey. In *Surveys on discrete and computational geometry*, Contemp. Math., 453, 49–86. Amer. Math. Soc., Providence, RI, 2008.
- [11] H. J. Bandelt, V. Chepoi, A. Dress and J. Koolen. Combinatorics of lopsided sets. *European J. Combin.*, 27:669–689, 2006.
- [12] H. J. Bandelt, V. Chepoi, A. Dress and J. Koolen. Geometry of lopsided sets. Preprint 2013.
- [13] H. J. Bandelt, V. Chepoi and K. Knauer. COMs: Complexes of oriented matroids. <http://arxiv.org/abs/1507.06111>.
- [14] A. Beilinson, V. Ginzburg, and W. Soergel. Koszul duality patterns in representation theory. *J. Amer. Math. Soc.*, 9(2):473–527, 1996.
- [15] D. J. Benson. *Representations and cohomology. I*, volume 30 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1998. Basic representation theory of finite groups and associative algebras.
- [16] M. Bestvina and N. Brady. Morse theory and finiteness properties of groups. *Invent. Math.*, 129(3):445–470, 1997.
- [17] P. Bidigare, P. Hanlon, and D. Rockmore. A combinatorial description of the spectrum for the Tsetlin library and its generalization to hyperplane arrangements. *Duke Math. J.*, 99(1):135–174, 1999.
- [18] T. P. Bidigare. *Hyperplane arrangement face algebras and their associated Markov chains*. ProQuest LLC, Ann Arbor, MI, 1997. Thesis (Ph.D.)—University of Michigan.
- [19] L. J. Billera, S. P. Holmes, and K. Vogtmann. Geometry of the space of phylogenetic trees. *Adv. in Appl. Math.*, 27(4):733–767, 2001.
- [20] L. J. Billera, S. K. Hsiao, and J. S. Provan. Enumeration in convex geometries and associated polytopal subdivisions of spheres. *Discrete Comput. Geom.* 39(1-3):123–137, 2008.
- [21] A. Björner. Posets, regular CW complexes and Bruhat order. *European J. Combin.*, 5(1):7–16, 1984.
- [22] A. Björner. Topological methods. In *Handbook of combinatorics, Vol. 1, 2*, pages 1819–1872. Elsevier, Amsterdam, 1995.
- [23] A. Björner. Random walks, arrangements, cell complexes, greedoids, and self-organizing libraries. In *Building bridges*, volume 19 of *Bolyai Soc. Math. Stud.*, pages 165–203. Springer, Berlin, 2008.
- [24] A. Björner. Note: Random-to-front shuffles on trees. *Electron. Commun. Probab.*, 14:36–41, 2009.
- [25] A. Björner, B. Korte, and L. Lovász. Homotopy properties of greedoids. *Adv. in Appl. Math.*, 6(4):447–494, 1985.
- [26] A. Björner, M. Las Vergnas, B. Sturmfels, N. White, and G. M. Ziegler. *Oriented matroids*, Second edition. Encyclopedia of Mathematics and its Applications, 46. Cambridge University Press, Cambridge, 1999.
- [27] A. Björner and G. M. Ziegler. Combinatorial stratification of complex arrangements. *J. Amer. Math. Soc.*, 5:105–149, 1992.
- [28] K. Bongartz. Algebras and quadratic forms. *J. London Math. Soc. (2)*, 28(3):461–469, 1983.
- [29] M. R. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 319. Springer-Verlag, Berlin, 1999.
- [30] K. S. Brown. Homological criteria for finiteness. *Comment. Math. Helv.*, 50:129–135, 1975.

- [31] K. S. Brown. *Buildings*. Springer-Verlag, New York, 1989.
- [32] K. S. Brown. The geometry of rewriting systems: a proof of the Anick-Groves-Squier theorem. In *Algorithms and classification in combinatorial group theory (Berkeley, CA, 1989)*, volume 23 of *Math. Sci. Res. Inst. Publ.*, pages 137–163. Springer, New York, 1992.
- [33] K. S. Brown. *Cohomology of groups*, volume 87 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994. Corrected reprint of the 1982 original.
- [34] K. S. Brown. Semigroups, rings, and Markov chains. *J. Theoret. Probab.*, 13(3):871–938, 2000.
- [35] K. S. Brown. Semigroup and ring theoretical methods in probability. In *Representations of finite dimensional algebras and related topics in Lie theory and geometry*, volume 40 of *Fields Inst. Commun.*, pages 3–26. Amer. Math. Soc., Providence, RI, 2004.
- [36] K. S. Brown and P. Diaconis. Random walks and hyperplane arrangements. *Ann. Probab.*, 26(4):1813–1854, 1998.
- [37] S. Burris and H. P. Sankappanavar. *A course in universal algebra*, volume 78 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1981.
- [38] J. C. Bustamante. The classifying space of a bound quiver. *J. Algebra*, 277(2):431–455, 2004.
- [39] J. F. Carlson, L. Townsley, L. Valeri-Elizondo, and M. Zhang. *Cohomology rings of finite groups*. Algebras and Applications, 3. Kluwer Academic Publishers, Dordrecht, 2003.
- [40] H. Cartan and S. Eilenberg. *Homological algebra*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1999. With an appendix by David A. Buchsbaum, Reprint of the 1956 original.
- [41] P. Cartier and D. Foata. *Problèmes combinatoires de commutation et réarrangements*. Lecture Notes in Mathematics, No. 85. Springer-Verlag, Berlin, 1969.
- [42] V. Chepoi. Graphs of some CAT(0) complexes. *Adv. in Appl. Math.* 24(2):125–179, 2000.
- [43] F. Chung and R. Graham. Edge flipping in graphs. *Adv. in Appl. Math.*, 48(1):37–63, 2012.
- [44] C. Cibils. Cohomology of incidence algebras and simplicial complexes. *J. Pure Appl. Algebra*, 56(3):221–232, 1989.
- [45] A. H. Clifford. Semigroups admitting relative inverses. *Ann. of Math. (2)*, 42:1037–1049, 1941.
- [46] A. H. Clifford and G. B. Preston. *The algebraic theory of semigroups. Vol. I*. Mathematical Surveys, No. 7. American Mathematical Society, Providence, R.I., 1961.
- [47] D. E. Cohen. A monoid which is right FP_∞ but not left FP_1 . *Bull. London Math. Soc.*, 24(4):340–342, 1992.
- [48] Charles W. Curtis and Irving Reiner. *Representation theory of finite groups and associative algebras*. Wiley Classics Library. John Wiley & Sons Inc., New York, 1988. Reprint of the 1962 original, A Wiley-Interscience Publication.
- [49] P. Diaconis. From shuffling cards to walking around the building: an introduction to modern Markov chain theory. In *Proceedings of the International Congress of Mathematicians, Vol. I (Berlin, 1998)*, number Extra Vol. I, pages 187–204, 1998.
- [50] V. Diekert and G. Rozenberg, editors. *The book of traces*. World Scientific Publishing Co. Inc., River Edge, NJ, 1995.
- [51] A. Dress, M. Hendy, K. Huber, K. and V. Moulton. On the number of vertices and edges of the Buneman graph. *Ann. Comb*, 1(4):329–337, 1997.
- [52] A. Dochtermann and A. Engström. Algebraic properties of edge ideals via combinatorial topology. *Electron. J. Combin.*, 16(2, Special volume in honor of Anders Björner):Research Paper 2, 24, 2009.

- [53] R. Fritsch and R. Piccinini. *Cellular structures in topology*. Cambridge Studies in Advanced Mathematics, 19. Cambridge University Press, Cambridge, 1990.
- [54] O. Ganyushkin, V. Mazorchuk, and B. Steinberg. On the irreducible representations of a finite semigroup. *Proc. Amer. Math. Soc.*, 137(11):3585–3592, 2009.
- [55] M. Gerstenhaber and S. D. Schack. Simplicial cohomology is Hochschild cohomology. *J. Pure Appl. Algebra*, 30(2):143–156, 1983.
- [56] R. Gray and S. J. Pride. Homological finiteness properties of monoids, their ideals and maximal subgroups, March 2010. <http://arxiv.org/abs/1003.3227>.
- [57] E. L. Green and R. Martínez-Villa. Koszul and Yoneda algebras. II. In *Algebras and modules, II (Geiranger, 1996)*, pages 227–244, CMS Conf. Proc., 24, Amer. Math. Soc., Providence, RI, 1998.
- [58] J. A. Green. On the structure of semigroups. *Ann. of Math. (2)*, 54:163–172, 1951.
- [59] V. S. Guba and S. J. Pride. Low-dimensional (co)homology of free Burnside monoids. *J. Pure Appl. Algebra*, 108(1):61–79, 1996.
- [60] V. S. Guba and S. J. Pride. On the left and right cohomological dimension of monoids. *Bull. London Math. Soc.*, 30(4):391–396, 1998.
- [61] F. Haglund and F. Paulin. Simplicité de groupes d’automorphismes d’espaces à courbure négative, in The Epstein birthday schrift, vol. 1 of *Geom. Topol. Monogr.*, Geom. Topol. Publ., Coventry, 1998, 181–248.
- [62] K. Igusa and D. Zacharia. On the cohomology of incidence algebras of partially ordered sets. *Comm. Algebra*, 18(3):873–887, 1990.
- [63] G. Kalai and R. Meshulam. Intersections of Leray complexes and regularity of monomial ideals. *J. Combin. Theory Ser. A*, 113(7):1586–1592, 2006.
- [64] G. Kalai and R. Meshulam. Leray numbers of projections and a topological Helly-type theorem. *J. Topol.*, 1(3):551–556, 2008.
- [65] Y. Kobayashi. The homological finiteness property FP_1 and finite generation of monoids. *Internat. J. Algebra Comput.*, 17(3):593–605, 2007.
- [66] Y. Kobayashi and F. Otto. For finitely presented monoids the homological finiteness conditions FHT and bi- FP_3 coincide. *J. Algebra*, 264(2):327–341, 2003.
- [67] Y. Lafont. A new finiteness condition for monoids presented by complete rewriting systems (after Craig C. Squier). *J. Pure Appl. Algebra*, 98(3):229–244, 1995.
- [68] J. Lawrence. Lopsided sets and orthant-intersection by convex sets. *Pacific J. Math.*, 104:155–173, 1983.
- [69] F. W. Lawvere. Qualitative distinctions between some toposes of generalized graphs. In *Categories in computer science and logic (Boulder, CO, 1987)*, volume 92 of *Contemp. Math.*, pages 261–299. Amer. Math. Soc., Providence, RI, 1989.
- [70] F. W. Lawvere. More on graphic toposes. *Cahiers Topologie Géom. Différentielle Catég.*, 32(1):5–10, 1991. International Category Theory Meeting (Bangor, 1989 and Cambridge, 1990).
- [71] F. W. Lawvere. Linearization of graphic toposes via Coxeter groups. *J. Pure Appl. Algebra*, 168(2-3):425–436, 2002. Category theory 1999 (Coimbra).
- [72] A. T. Lundell and S. Weingram. *The topology of CW complexes*. The University Series in Higher Mathematics. Springer, 1969.
- [73] S. W. Margolis, F. V. Saliola and B. Steinberg. Semigroups embeddable in hyperplane face monoids. *Semigroup Forum*, 89(1):236–248, 2014.
- [74] S. W. Margolis, F. V. Saliola and B. Steinberg. Poset topology and homological invariants of algebras arising in algebraic combinatorics. FPSAC 2014, Chicago, DMTCS proc. **AT**, 2014, 71–82.
- [75] S. W. Margolis, F. V. Saliola and B. Steinberg. Combinatorial topology and the global dimension of algebras arising in combinatorics. *Eur. J. Math.*, to appear.
- [76] R. Martínez-Villa. Introduction to Koszul algebras. *Rev. Un. Mat. Argentina*, 48(2):67–95, 2007.

- [77] W. S. Massey. *A basic course in algebraic topology*. Graduate Texts in Mathematics, 127. Springer-Verlag, New York, 1991.
- [78] P. McMullen, and E. Schulte. *Abstract regular polytopes*. Encyclopedia of Mathematics and its Applications, 92. Cambridge University Press, Cambridge, 2002.
- [79] S. Moran. *Shattering-extremal systems*. Master's Thesis, Saarland University, 2012.
- [80] W. D. Munn. The Jacobson radical of a band ring. *Math. Proc. Cambridge Philos. Soc.*, 105(2):277–283, 1989.
- [81] J. R. Munkres, James R. *Elements of algebraic topology*. Addison-Wesley Publishing Company, Menlo Park, CA.
- [82] W. R. Nico. On the cohomology of finite semigroups. *J. Algebra*, 11:598–612, 1969.
- [83] W. R. Nico. Homological dimension in semigroup algebras. *J. Algebra*, 18:404–413, 1971.
- [84] W. R. Nico. A counterexample in the cohomology of monoids. *Semigroup Forum*, 4:93–94, 1972.
- [85] W. R. Nico. An improved upper bound for global dimension of semigroup algebras. *Proc. Amer. Math. Soc.*, 35:34–36, 1972.
- [86] M. Nunes. Cohomological results in monoid and category theory via classifying spaces. *J. Pure Appl. Algebra*, 101(2):213–244, 1995.
- [87] P. Polo, Patrick. On Cohen-Macaulay posets, Koszul algebras and certain modules associated to Schubert varieties. *Bull. London Math. Soc.* 27(5):425–434, 1995.
- [88] S. J. Pride. Homological finiteness conditions for groups, monoids, and algebras. *Comm. Algebra*, 34(10):3525–3536, 2006.
- [89] M. S. Putcha. Complex representations of finite monoids. II. Highest weight categories and quivers. *J. Algebra*, 205(1):53–76, 1998.
- [90] D. Quillen. Higher algebraic K-theory. I. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.
- [91] V. Reiner, F. Saliola and V. Welker. Spectra of symmetrized shuffling operators. *Mem. Amer. Math. Soc.*, 228 (1072), 2014.
- [92] J. Rhodes and B. Steinberg. *The q-theory of finite semigroups*. Springer Monographs in Mathematics. Springer, New York, 2009.
- [93] J. Richter-Gebert and G. M. Ziegler. Zonotopal tilings and the Bohne-Dress theorem. In *Jerusalem combinatorics '93*, pages 211–232, Contemp. Math., 178, Amer. Math. Soc., Providence, RI, 1994.
- [94] M. Roller. Poc-sets, median algebras and group actions. An extended study of Dunwoody's construction and Sageev's theorem. Preprint, 1998.
- [95] L. H. Rowen. *Ring theory. Vol. I*. Pure and Applied Mathematics, 127. Academic Press, Inc., Boston, MA, 1988.
- [96] F. V. Saliola. The quiver of the semigroup algebra of a left regular band. *Internat. J. Algebra Comput.*, 17(8):1593–1610, 2007.
- [97] F. V. Saliola. On the quiver of the descent algebra. *J. Algebra*, 320(11):3866–3894, 2008.
- [98] F. V. Saliola. The face semigroup algebra of a hyperplane arrangement. *Canad. J. Math.*, 61(4):904–929, 2009.
- [99] F. V. Saliola. The Loewy length of the descent algebra of type D . *Algebr. Represent. Theory*, 13(2):243–254, 2010.
- [100] F. Saliola and H. Thomas, Oriented interval greedoids. *Discrete Comput. Geom.*, 47(1):64–105, 2012.
- [101] M. Schocker. The module structure of the Solomon-Tits algebra of the symmetric group. *J. Algebra*, 301(2):554–586, 2006.
- [102] M.-P. Schützenberger. Sur certain treillis gauches. *C.R. Acad. Sci. Paris*, 224:776–778, 1947.

- [103] G. Segal. Classifying spaces and spectral sequences. *Inst. Hautes Études Sci. Publ. Math.*, 34:105–112, 1968.
- [104] L. Solomon. The Burnside algebra of a finite group. *J. Combinatorial Theory*, 2:603–615, 1967.
- [105] L. Solomon. A Mackey formula in the group ring of a Coxeter group. *J. Algebra*, 41(2):255–264, 1976.
- [106] D. Speyer and B. Sturmfels. The tropical Grassmannian. *Adv. Geom.*, 4(3):389–411, 2004.
- [107] C. C. Squier. Word problems and a homological finiteness condition for monoids. *J. Pure Appl. Algebra*, 49(1-2):201–217, 1987.
- [108] R. P. Stanley. *Cohen-Macaulay complexes*. in *Higher combinatorics (Proc. NATO Advanced Study Inst., Berlin, 1976)*. NATO Adv. Study Inst. Ser., Ser. C: Math. and Phys. Sci., 31, 51–62. Reidel, Dordrecht, 1977.
- [109] R. P. Stanley. Two poset polytopes. *Discrete Comput. Geom.*, 1:9–23, 1986.
- [110] R. P. Stanley. *Combinatorics and commutative algebra*, volume 41 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, second edition, 1996.
- [111] R. P. Stanley. An introduction to hyperplane arrangements. in *Geometric combinatorics*. IAS/Park City Math. Ser., 13, 389–496. Amer. Math. Soc., Providence, RI, 2007.
- [112] B. Steinberg. *Isomorphism of oriented matroids from a semigroup viewpoint*. <http://bensteinberg.wordpress.com/2013/08/13/equivalence-or-oriented-matroids-from-a-semigroup-viewpoint>
- [113] J. Tits. *Buildings of spherical type and finite BN-pairs*. Springer-Verlag, Berlin, 1974, Lecture Notes in Mathematics, Vol. 386.
- [114] J. Tits. Two properties of Coxeter complexes. *J. Algebra*, 41(2):265–268, 1976. Appendix to “A Mackey formula in the group ring of a Coxeter group” (*J. Algebra* 41 (1976), no. 2, 255–264) by Louis Solomon.
- [115] P. Webb. An introduction to the representations and cohomology of categories. In *Group representation theory*, pages 149–173. EPFL Press, Lausanne, 2007.
- [116] G. Wegner. d -collapsing and nerves of families of convex sets. *Arch. Math. (Basel)*, 26:317–321, 1975.
- [117] N. White (ed.). *Theory of Matroids*, Encyclopedia of Mathematics and its Applications, 26. Cambridge University Press, 1986.
- [118] D. T. Wise. *From riches to raags: 3-manifolds, right-angled Artin groups, and cubical geometry*. CBMS Regional Conference Series in Mathematics, 117. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2012.
- [119] D. Woodcock. Cohen-Macaulay complexes and Koszul rings. *J. London Math. Soc.* (2), 57(2):398–410, 1998.
- [120] R. Woodroffe. Matchings, coverings, and Castelnuovo-Mumford regularity. *J. Commut. Algebra* 6(2):287–304, 2014.
- [121] G. M. Ziegler. What is a complex matroid? *Discrete Comput. Geom.* 10:313–348, 1993.
- [122] G. M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. Theory and applications.

INDEX

- T -convex, 24
- W -permutohedron, 20
- $\mathbb{k}$ -Leray number, 31
- $\mathbb{k}$ -cohomological dimension, 118
- Ext-algebra, 65
- $\mathcal{P}$ -left regular band, 46
- $\mathcal{J}$ -equivalent, 10
- $\mathcal{L}$ -equivalent, 10
- $\mathcal{R}$ -equivalent, 10
- f -vector, 113
- q -cell, 32
- q -simplex, 29
- q -skeleton, 30, 32
- (Jacobson) radical, 58

- acyclic, 34
- adjacent, 99
- admissible, 61
- affine, 18
- affine oriented matroid, 23
- algebra
 - graded, 64
 - Koszul, 65
 - quadratic, 64
- anti-exchange, 42
- antimatroids, 42
- arrow ideal, 61
- atom, 16
- augmented intersection lattice, 28

- band, 10
 - left regular, 10
- bases, 15
- basic, 59
- blocks, 61
- boolean arrangement, 19
- bound quiver, 61
- braid arrangement, 19

- Cartan matrix, 77
- CAT(0)
 - cone, 50
 - cube complex, 49
 - metric space, 49
 - zonotopal complex, 57
- cellular, 32, 33
- central, 18
- central primitive idempotents, 61
- chamber complex, 99
- chambers, 18, 21
- chordal, 31, 91
- clique complex, 30
- closed n -cell, 31
- closed set, 37, 42
- cocircuit, 21, 110
- Cohen-Macaulay, 29
- COM, 26
 - ranking, 26
 - realizable, 25
- comparison triangle, 49
- complex hyperplane arrangement, 28
- complex of oriented matroids, 26
- composition
 - of covectors, 38
- conditional oriented matroid, 26
- cone, 30
 - CAT(0), 50
- cone point, 30
- connected, 75
- continuations, 37
- contraction, 11, 21, 40
- convex, 49
- convex geometry, 42
- covector
 - oriented interval greedoid, 38
- covectors, 13, 27
- Coxeter zonotope, 57
- cube complex, 48
 - CAT(0), 49
- CW poset, 32

- degeneration, 52
- degree, 64
- deletion, 22, 41, 42
- dependent, 16
- dimension, 32
- dual
 - Koszul, 65
 - quadratic, 65

- equivalent
 - $\mathcal{J}$ -, 10
 - $\mathcal{L}$ -, 10
 - $\mathcal{R}$ -, 10
- essential, 18, 24, 25, 28
- Euler characteristic, 113
- even polyhedron, 57
- extreme points, 42

- face monoid, 18, 28
- face semigroup, 24, 57
- face symmetry axiom, 26

- faces, 29
- facets, 29
- Fine f -vector, 113
- flag complex, 30
- flag vector, 113
- flat, 16, 36
 - greedoid, 37
- free left regular band, 14
- free partially commutative left regular band, 17
- gallery, 99
- generic, 107, 111
- geodesic, 49
 - metric space, 49
 - triangle, 49
- geometric, 92
- geometric lattice, 16
- geometric realization, 29
- global dimension, 60
- graded, 32
 - algebra, 64
 - module, 64
 - poset, 16
- graphic monoids, 10
- graphical arrangement, 19
- greedoid flat, 36, 37
- Green's relations, 10
 - $\mathcal{J}$, 10
 - $\mathcal{L}$, 10
 - $\mathcal{R}$, 10
- ground set, 15, 20, 24, 26
- half-spaces, 22, 54
- Helly property, 54
- hereditary, 15, 60
- homogeneous ideal, 64
- hyperplane, 53
- hyperplane arrangement, 18
- hyperplane face semigroup, 23
- ideal
 - homogeneous, 64
- idempotent, 10
- in general position, 107, 111
- incidence algebra, 70, 100
- indecomposable, 59
- independent subsets, 15
- induced, 29
- injective
 - envelope, 60
 - hull, 60
- interior edge, 51
- intersection
 - lattice, 19
 - augmented, 28
 - semilattice, 54
 - interval greedoid, 37
 - isomorphism
 - CW complexes, 32
- join, 13, 30
- Koszul
 - algebra, 65
 - dual, 65
- ladders, 46
- lattice
 - geometric, 16
 - semimodular, 37
- left regular band, 10
 - monoid, 10
- left zero multiplication, 65
- lexicographic extension, 111
- linear resolution, 65
- link, 29, 49
- loop, 15, 22, 37
- lopsided system, 25
- Möbius function, 71
- matroid, 15
- maximal, 58
- median graph, 54
- metric, 51
- metric space
 - CAT(0), 49
 - geodesic, 49
- mid-hyperplane, 57
- midcube, 53
- minimal, 62
 - projective resolution, 93
- modular, 58
- module
 - graded, 64
- Morita equivalent, 59
- necklace of partitions of $[n]$, 110
- nerve, 54
- open, 25
- open n -cell, 31
- order complex, 30
- orientation, 79
- oriented q -simplex, 79
- oriented interval greedoid, 39
- oriented matroid, 20
 - realizable, 21

- parallel, 15
- path algebra, 60
- permutohedron, 20
- phylogenetic tree space, 50, 51
- poset
 - CW, 32
 - graded, 16
 - ranked, 16
 - strongly connected, 99
 - thin, 101
- primitive, 59
- principal series, 83
- projective cover, 60
- projective dimension, 60
- pure, 29
- quadratic
 - algebra, 64
 - dual, 65
- quiver, 60
 - presentation, 62
- radical, 59
- rank, 15, 21, 32
- rank function, 16
- ranked
 - poset, 16
- ranking COM, 26
- realizable
 - COM, 25
 - oriented matroid, 21
 - strong elimination system, 24
- regular, 32
- regular band, 120
- regular CW complex, 31
- relative simplicial chain complex, 80
- representation
 - Schützenberger, 68
- resolution
 - linear, 65
 - minimal projective, 93
- right hereditary, 91
- Schützenberger representation, 68
- semi-free, 82
- semimodular
 - lattice, 37
 - rank function, 15
- semiprimitive, 58
- semisimple, 59
- separation set, 20, 38
- simple, 15
- simplices, 29
- simplicial chain complex, 79
- simplicial complex, 29
- simplicial map, 30, 80
- single element extension, 110
- single-element lifting, 22
- socle, 60
- spherical left regular band, 46
- split, 59
- stabilizer, 78
- strong elimination, 21
- strong elimination system, 24
 - realizable, 24
- strongly connected poset, 99
- subcomplex, 29
- support map, 11
- support semilattice, 11
- suspension, 14, 30
- system of relations, 62
- thick, 52
- thin poset, 101
- top, 59
- topes, 21, 26
- underlying matroid, 36
- unitary, 58, 78
- upper semimodular, 16
- vertex, 32
- vertices, 29
- visual hemisphere, 107, 111
- zero set, 21
- zeta function, 71
- zonotopal complex, 57
 - CAT(0), 57
- zonotopal tiling, 23
- zonotope, 18
 - polar to an arrangement, 18

(S. Margolis) DEPARTMENT OF MATHEMATICS, BAR ILAN UNIVERSITY, 52900 RAMAT GAN, ISRAEL

E-mail address, S. Margolis: `margolis@math.biu.ac.il`

(F. Saliola) DÉPARTEMENT DE MATHÉMATIQUES – LACIM, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, C.P. 8888, SUCCURSALE CENTRE-VILLE, MONTRÉAL, QUÉBEC H3C 3P8, CANADA

E-mail address, F. Saliola: `saliola.franco@uqam.ca`

(B. Steinberg) DEPARTMENT OF MATHEMATICS, CITY COLLEGE OF NEW YORK, CONVENT AVENUE AT 138TH STREET, NEW YORK, NEW YORK 10031, USA

E-mail address, B. Steinberg: `bsteinberg@ccny.cuny.edu`